



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:50 PM UTC

PDB ID : 6NBW / pdb_00006nbw
Title : Ternary Complex of Beta/Gamma-Actin with Profilin and AnCoA-NAA80
Authors : Rebowski, G.; Boczkowska, M.; Dominguez, R.
Deposited on : 2018-12-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	
2	N	235	
3	P	140	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 10938 atoms, of which 5345 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, cytoplasmic 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	367	Total	C	H	N	O	S	0	1	0
			5721	1822	2842	485	552	20			

- Molecule 2 is a protein called N-alpha-acetyltransferase 80.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	N	167	Total	C	H	N	O	S	0	0	0
			2692	852	1358	249	227	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	74	ALA	-	expression tag	UNP Q93015
N	75	GLY	-	expression tag	UNP Q93015
N	76	HIS	-	expression tag	UNP Q93015
N	77	MET	-	expression tag	UNP Q93015


- Molecule 3 is a protein called Profilin-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	P	138	Total	C	H	N	O	S	0	0	0
			2081	654	1040	179	201	7			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	P	
			43	10	12	5	13	3	

- Molecule 6 is LATRUNCULIN B (CCD ID: LAB) (formula: $C_{20}H_{29}NO_5S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	S	
			56	20	29	1	5	1	

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			14	3	8	3		
7	A	1	Total	C	H	O	0	0
			13	3	7	3		

- Molecule 8 is [(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-4-HYDROXY-3-(PHOSPHONOXY)TETRAHYDROFURAN-2-YL]METHYL (3R)-3-HYDROXY-2,2-DIMETHYL-4-OXO-4-{[3-OXO-3-(2-[(2-OXOPROPYL)THIO]ETHYL}AMINO)PROPYL]AMINO}BUTYL DIHYDROGEN DIPHOSPHATE (CCD ID: SOP) (formula: C₂₄H₄₀N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
8	N	1	Total	C	H	N	O	P	S	0	0
			88	24	36	7	17	3	1		

- Molecule 9 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
9	N	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	116	Total	O	0	0
			116	116		
10	N	57	Total	O	0	0
			57	57		
10	P	31	Total	O	0	0
			31	31		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Actin, cytoplasmic 1



- Molecule 2: N-alpha-acetyltransferase 80



- Molecule 3: Profilin-1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	104.41Å 115.59Å 132.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.10 – 2.50 26.10 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (26.10-2.50) 99.5 (26.10-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.09 (at 2.50Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.178 , 0.209 0.178 , 0.209	Depositor DCC
R_{free} test set	1403 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtriage
Anisotropy	0.817	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10938	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SOP, LAB, ATP, HIC, MES, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/2931	0.34	0/3966
2	N	0.15	0/1368	0.36	0/1854
3	P	0.12	0/1059	0.32	0/1430
All	All	0.14	0/5358	0.34	0/7250

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2879	2842	2842	22	0
2	N	1334	1358	1357	12	0
3	P	1041	1040	1040	5	0
4	A	1	0	0	0	0
5	A	31	12	12	0	0
6	A	27	29	29	2	0
7	A	12	15	16	0	0
8	N	52	36	35	0	0
9	N	12	13	13	0	0
10	A	116	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	N	57	0	0	0	0
10	P	31	0	0	0	0
All	All	5593	5345	5344	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:403:LAB:C16	6:A:403:LAB:N1	1.70	1.40
1:A:273:GLY:O	1:A:277:THR:HG23	1.88	0.73
2:N:107:ARG:NH1	2:N:111:SER:OG	2.31	0.63
1:A:372[A]:ARG:NH1	3:P:82:GLU:OE1	2.32	0.62
2:N:82:LEU:HD12	2:N:177:ARG:CZ	2.37	0.55
6:A:403:LAB:N1	6:A:403:LAB:C15	2.61	0.55
1:A:348:SER:HB3	2:N:222:THR:HG22	1.91	0.52
1:A:286:ASP:OD2	3:P:90:LYS:NZ	2.38	0.52
1:A:272:CYS:HB2	1:A:277:THR:HG22	1.94	0.49
1:A:345:ILE:HD11	2:N:227:PRO:HD2	1.95	0.49
1:A:4:ASP:O	1:A:5:ILE:HG22	2.12	0.48
1:A:351:THR:O	1:A:354:GLN:HG2	2.13	0.48
1:A:109:PRO:HB3	1:A:175:ILE:HD13	1.96	0.47
1:A:190:MET:HG2	1:A:209:VAL:HG21	1.96	0.47
1:A:352:PHE:HE2	1:A:356:TRP:CH2	2.31	0.47
2:N:301:ILE:HD12	2:N:301:ILE:O	2.16	0.46
1:A:25:ASP:HA	2:N:226:LEU:HD21	1.98	0.45
1:A:303:THR:HG22	1:A:303:THR:O	2.16	0.45
2:N:123:PHE:HA	2:N:124:PRO:C	2.41	0.45
1:A:283:MET:HE3	1:A:294:TYR:OH	2.18	0.44
3:P:25:LYS:HE2	3:P:106:ASP:O	2.17	0.44
2:N:122:ALA:O	2:N:123:PHE:HB2	2.18	0.43
1:A:318:THR:HA	1:A:327:ILE:CD1	2.49	0.43
2:N:87:VAL:HA	2:N:128:MET:HE2	2.00	0.43
1:A:321:ALA:HB3	1:A:327:ILE:HD11	2.01	0.43
1:A:70:PRO:HG3	1:A:81:ASP:HB3	2.01	0.42
1:A:180:LEU:C	1:A:180:LEU:HD23	2.44	0.42
3:P:47:VAL:O	3:P:51:VAL:HG22	2.19	0.42
2:N:227:PRO:O	2:N:228:ALA:HB3	2.19	0.42
3:P:80:ASP:HA	3:P:83:PHE:CZ	2.54	0.42
2:N:125:LEU:C	2:N:125:LEU:HD12	2.44	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:LEU:HB3	1:A:351:THR:HG22	2.02	0.41
1:A:198:TYR:CE1	1:A:248:ILE:HG22	2.55	0.41
2:N:93:LEU:HD21	2:N:140:PRO:HG2	2.02	0.40
1:A:3:ASP:O	1:A:4:ASP:CB	2.69	0.40
1:A:172:PRO:HA	1:A:175:ILE:HD12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/374 (97%)	356 (98%)	6 (2%)	1 (0%)	36	55
2	N	159/235 (68%)	153 (96%)	5 (3%)	1 (1%)	21	38
3	P	136/140 (97%)	132 (97%)	4 (3%)	0	100	100
All	All	658/749 (88%)	641 (97%)	15 (2%)	2 (0%)	36	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
2	N	123	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/316 (99%)	304 (97%)	8 (3%)	40	68
2	N	147/201 (73%)	143 (97%)	4 (3%)	39	67
3	P	113/114 (99%)	111 (98%)	2 (2%)	51	77
All	All	572/631 (91%)	558 (98%)	14 (2%)	44	70

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	116	ARG
1	A	118	LYS
1	A	132	MET
1	A	281	SER
1	A	324	THR
1	A	372[A]	ARG
1	A	372[B]	ARG
2	N	78	SER
2	N	214	GLU
2	N	263	LEU
2	N	271	GLU
3	P	53	LYS
3	P	58	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	GLN
1	A	354	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	HIC	A	73	1	10,11,12	1.53	2 (20%)	9,14,16	1.43	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	73	1	-	0/5/6/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	HIC	CD2-CG	3.07	1.41	1.36
1	A	73	HIC	CD2-NE2	2.08	1.40	1.37

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	HIC	NE2-CE1-ND1	-3.73	111.23	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	SOP	N	400	-	52,54,54	2.62	15 (28%)	75,80,80	2.11	18 (24%)
5	ATP	A	402	4	32,33,33	1.48	8 (25%)	48,52,52	1.74	10 (20%)
7	GOL	A	405	-	5,5,5	0.89	0	5,5,5	1.02	0
6	LAB	A	403	-	28,29,29	5.23	15 (53%)	29,41,41	4.27	5 (17%)
7	GOL	A	404	-	5,5,5	1.10	0	5,5,5	1.06	1 (20%)
9	MES	N	401	-	12,12,12	2.26	1 (8%)	15,16,16	1.34	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	SOP	N	400	-	-	12/52/68/68	0/3/3/3
5	ATP	A	402	4	-	2/22/38/38	0/3/3/3
7	GOL	A	405	-	-	2/4/4/4	-
6	LAB	A	403	-	-	2/21/49/49	0/2/3/3
7	GOL	A	404	-	-	2/4/4/4	-
9	MES	N	401	-	-	0/6/14/14	0/1/1/1

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	403	LAB	C16-N1	19.08	1.70	1.46
6	A	403	LAB	C2-C3	11.52	1.54	1.33
8	N	400	SOP	C3'-C4'	-8.35	1.31	1.52
8	N	400	SOP	O4'-C4'	7.74	1.62	1.45
9	N	401	MES	C8-S	-7.48	1.67	1.77
6	A	403	LAB	C7-C6	6.54	1.58	1.31
6	A	403	LAB	C18-S1	-6.04	1.65	1.77
8	N	400	SOP	CP6-NP2	5.32	1.46	1.33
8	N	400	SOP	P1-O6	5.31	1.65	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	N	400	SOP	CP3-NP1	5.23	1.45	1.33
8	N	400	SOP	P2-O6	5.19	1.65	1.59
6	A	403	LAB	O2-C1	5.01	1.45	1.34
6	A	403	LAB	C17-S1	-4.96	1.69	1.81
6	A	403	LAB	C12-C11	4.73	1.63	1.52
6	A	403	LAB	C15-C16	-4.66	1.45	1.54
8	N	400	SOP	O4'-C1'	-4.58	1.31	1.42
8	N	400	SOP	C6-N6	4.45	1.45	1.34
6	A	403	LAB	C10-C11	-4.42	1.37	1.51
5	A	402	ATP	C5-C4	4.37	1.46	1.39
6	A	403	LAB	C2-C1	4.04	1.56	1.46
6	A	403	LAB	O3-C11	3.76	1.52	1.44
8	N	400	SOP	P3-O3'	3.30	1.65	1.59
5	A	402	ATP	PB-O3B	3.23	1.63	1.59
6	A	403	LAB	C9-C10	3.10	1.61	1.53
6	A	403	LAB	C17-C16	-2.88	1.46	1.52
6	A	403	LAB	O3-C15	2.85	1.48	1.43
8	N	400	SOP	O2'-C2'	-2.64	1.36	1.43
5	A	402	ATP	C5-C6	2.60	1.48	1.41
6	A	403	LAB	C4-C3	2.55	1.56	1.51
8	N	400	SOP	C5-C4	-2.45	1.34	1.39
5	A	402	ATP	C8-N7	2.39	1.36	1.31
8	N	400	SOP	C8-N9	-2.34	1.33	1.37
5	A	402	ATP	C5-N7	-2.33	1.34	1.39
8	N	400	SOP	C5-N7	-2.24	1.35	1.39
5	A	402	ATP	PB-O3A	2.22	1.61	1.59
8	N	400	SOP	OP2-CP6	-2.21	1.19	1.23
5	A	402	ATP	PA-O3A	2.19	1.61	1.59
8	N	400	SOP	OP1-CP3	-2.17	1.18	1.23
5	A	402	ATP	C4-N9	-2.12	1.33	1.37

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	403	LAB	C17-S1-C18	21.76	104.64	92.04
8	N	400	SOP	C4-N9-C1'	-6.81	110.70	126.63
8	N	400	SOP	N3-C2-N1	-5.59	120.12	128.58
5	A	402	ATP	C5-C4-N3	-5.46	119.20	126.72
8	N	400	SOP	N9-C8-N7	-5.22	106.53	113.94
8	N	400	SOP	C1'-N9-C8	5.18	138.59	127.09
8	N	400	SOP	C5-C4-N3	-5.09	119.71	126.72
8	N	400	SOP	C4-N9-C8	4.71	110.68	105.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	N	400	SOP	N3-C4-N9	4.43	134.70	127.17
8	N	400	SOP	N6-C6-N1	-4.42	108.54	118.38
5	A	402	ATP	N3-C4-N9	4.30	134.47	127.17
5	A	402	ATP	C4-C5-N7	-3.76	106.28	110.58
5	A	402	ATP	C2-N3-C4	3.60	120.62	111.83
5	A	402	ATP	N3-C2-N1	-3.51	123.27	128.58
8	N	400	SOP	C2-N3-C4	3.39	120.11	111.83
5	A	402	ATP	C4-N9-C8	3.33	109.24	105.74
8	N	400	SOP	CP1-CP2-NP1	-3.30	105.53	112.41
8	N	400	SOP	C5-C6-N6	3.26	131.36	123.29
8	N	400	SOP	C5-N7-C8	3.09	108.31	103.45
6	A	403	LAB	C4-C5-C6	-2.94	106.17	112.57
5	A	402	ATP	C5-N7-C8	2.92	108.05	103.45
5	A	402	ATP	N9-C8-N7	-2.69	110.11	113.94
8	N	400	SOP	CA3-CA2-CA1	2.64	120.58	116.17
8	N	400	SOP	CPA-CP8-CP7	2.55	113.11	108.77
5	A	402	ATP	C6-C5-N7	2.34	136.60	132.09
5	A	402	ATP	C2-N1-C6	2.29	122.49	118.73
6	A	403	LAB	C19-C3-C4	2.27	119.17	115.23
8	N	400	SOP	CP2-NP1-CP3	-2.19	118.75	122.82
8	N	400	SOP	P3-O3'-C3'	-2.18	117.62	123.43
6	A	403	LAB	C9-C10-C11	-2.15	110.14	114.06
9	N	401	MES	O3S-S-C8	2.14	110.19	106.00
8	N	400	SOP	C3'-C2'-C1'	2.10	104.52	99.89
8	N	400	SOP	CP1-S-CA1	2.08	105.15	101.72
6	A	403	LAB	O5-C18-S1	2.08	126.77	123.06
7	A	404	GOL	C3-C2-C1	-2.01	104.42	111.80

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	402	ATP	PB-O3B-PG-O3G
7	A	404	GOL	C1-C2-C3-O3
7	A	405	GOL	C1-C2-C3-O3
8	N	400	SOP	C5'-O5'-P1-O11
8	N	400	SOP	C5'-O5'-P1-O12
8	N	400	SOP	C5'-O5'-P1-O6
8	N	400	SOP	CP9-O7-P2-O6
8	N	400	SOP	CP9-O7-P2-O21
8	N	400	SOP	CP9-O7-P2-O22
7	A	404	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	A	405	GOL	O2-C2-C3-O3
8	N	400	SOP	OP3-CP7-CP8-CP9
6	A	403	LAB	O2-C1-C2-C3
6	A	403	LAB	O1-C1-C2-C3
8	N	400	SOP	C3'-O3'-P3-O31
8	N	400	SOP	CP2-CP1-S-CA1
5	A	402	ATP	PB-O3B-PG-O1G
8	N	400	SOP	OP3-CP7-CP8-CPB
8	N	400	SOP	CP8-CP9-O7-P2
8	N	400	SOP	CP6-CP7-CP8-CP9


There are no ring outliers.

1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	403	LAB	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand SOP N 400



Ligand ATP A 402





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	366/374 (97%)	-0.38	8 (2%) 62 58	26, 38, 56, 81	1 (0%)
2	N	167/235 (71%)	-0.21	9 (5%) 31 27	32, 39, 90, 115	0
3	P	138/140 (98%)	-0.25	0 100 100	36, 45, 72, 81	0
All	All	671/749 (89%)	-0.31	17 (2%) 58 54	26, 40, 68, 115	1 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	N	133	HIS	4.0
1	A	3	ASP	3.7
1	A	2	ASP	3.7
1	A	4	ASP	3.4
2	N	228	ALA	3.3
2	N	229	THR	3.1
1	A	40	HIS	2.6
1	A	48	GLY	2.6
2	N	76	HIS	2.6
1	A	356	TRP	2.6
2	N	230	LEU	2.5
2	N	214	GLU	2.2
1	A	5	ILE	2.1
2	N	263	LEU	2.1
2	N	268	PRO	2.1
1	A	270	GLU	2.0
2	N	270	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	HIC	A	73	11/12	0.95	0.07	28,31,39,39	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	A	404	6/6	0.74	0.18	34,40,45,45	0
9	MES	N	401	12/12	0.78	0.17	41,50,55,69	0
7	GOL	A	405	6/6	0.92	0.08	44,52,55,59	0
6	LAB	A	403	27/27	0.96	0.07	26,31,34,36	0
8	SOP	N	400	52/52	0.97	0.07	29,35,40,43	0
5	ATP	A	402	31/31	0.98	0.05	24,29,35,38	0
4	CA	A	401	1/1	0.98	0.02	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LAB A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SOP N 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ATP A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.