



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:12 PM UTC

PDB ID : 7NBS / pdb_00007nbs
Title : Crystal structure of SARS-CoV-2 main protease (Mpro) in complex with the HCV NS3/4A inhibitor telaprevir
Authors : Zhang, L.; Hilgenfeld, R.
Deposited on : 2021-01-27
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	306	15% 89% 10%

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	305	Total	C	N	O	S	0	0	0
			2351	1488	400	441	22			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total Cl 1 1	0	0

- Molecule 3 is (1S,3aR,6aS)-2-[(2S)-2-({(2S)-2-cyclohexyl-2-[(pyrazin-2-ylcarbonyl)amino]acetyl}amino)-3,3-dimethylbutanoyl]-N-[(2R,3S)-1-(cyclopropylamino)-2-hydroxy-1-oxohexan-3-yl]octahydrocyclopenta[c]pyrrole-1-carboxamide (CCD ID: SV6) (formula: C₃₆H₅₅N₇O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	N	O	0	0
			49	36	7	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	155	Total 155	O 155	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.69Å 54.62Å 48.28Å 90.00° 101.40° 90.00°	Depositor
Resolution (Å)	48.70 – 1.70 48.70 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.70-1.70) 100.0 (48.70-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.206 , 0.256 0.213 , 0.266	Depositor DCC
R_{free} test set	1536 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2556	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SV6, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.02	0/2403	1.32	4/3266 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	127	GLN	CB-CA-C	-6.80	97.89	109.51
1	AAA	2	GLY	CA-C-O	-5.92	118.15	122.23
1	AAA	230	PHE	CA-CB-CG	-5.79	108.01	113.80
1	AAA	196	THR	CA-CB-OG1	-5.26	101.71	109.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2351	0	2298	27	0
2	AAA	1	0	0	0	0
3	AAA	49	0	54	5	0
4	AAA	155	0	0	6	0
All	All	2556	0	2352	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:19:GLN:NE2	1:AAA:119:ASN:HD22	1.57	1.02
1:AAA:19:GLN:HE21	1:AAA:119:ASN:HD22	1.19	0.81
1:AAA:197:ASP:OD2	4:AAA:501:HOH:O	2.02	0.76
1:AAA:194:ALA:O	4:AAA:502:HOH:O	2.06	0.73
1:AAA:243:THR:H	1:AAA:246:HIS:CD2	2.19	0.60
1:AAA:167:LEU:HD22	3:AAA:402:SV6:HBL	1.85	0.58
1:AAA:188:ARG:HD3	4:AAA:541:HOH:O	2.05	0.57
1:AAA:24:THR:HG23	4:AAA:576:HOH:O	2.05	0.57
1:AAA:95:ASN:HB3	1:AAA:98:THR:OG1	2.06	0.55
3:AAA:402:SV6:HALB	4:AAA:602:HOH:O	2.07	0.54
1:AAA:19:GLN:NE2	1:AAA:119:ASN:ND2	2.41	0.53
1:AAA:167:LEU:HD23	1:AAA:167:LEU:N	2.23	0.53
1:AAA:109:GLY:HA2	1:AAA:200:ILE:HD13	1.91	0.52
1:AAA:19:GLN:HE21	1:AAA:26:THR:HG21	1.77	0.49
1:AAA:233:VAL:HG11	1:AAA:269:LYS:HG3	1.94	0.48
1:AAA:10:SER:O	1:AAA:14:GLU:HG3	2.12	0.48
1:AAA:167:LEU:HD22	3:AAA:402:SV6:CBL	2.44	0.47
1:AAA:5:LYS:HD3	4:AAA:528:HOH:O	2.13	0.47
1:AAA:113:SER:O	1:AAA:149:GLY:HA2	2.15	0.47
1:AAA:243:THR:H	1:AAA:246:HIS:HD2	1.63	0.45
1:AAA:70:ALA:HB3	1:AAA:73:VAL:HG13	2.00	0.44
1:AAA:5:LYS:HG2	1:AAA:127:GLN:HB2	1.99	0.44
1:AAA:153:ASP:OD1	1:AAA:153:ASP:N	2.51	0.44
1:AAA:49:MET:HE2	3:AAA:402:SV6:HD2	2.01	0.43
1:AAA:96:PRO:C	1:AAA:98:THR:H	2.27	0.42
1:AAA:232:LEU:HA	1:AAA:235:MET:HE2	2.02	0.42
1:AAA:19:GLN:HE22	1:AAA:119:ASN:HD22	1.57	0.41
1:AAA:49:MET:HE2	3:AAA:402:SV6:HG	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	303/306 (99%)	294 (97%)	6 (2%)	3 (1%)	12 3

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	71	GLY
1	AAA	97	LYS
1	AAA	73	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	261/263 (99%)	258 (99%)	3 (1%)	65 54

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	73	VAL
1	AAA	153	ASP
1	AAA	235	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SV6	AAA	402	1	53,53,53	1.05	4 (7%)	69,76,76	1.61	11 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SV6	AAA	402	1	-	1/57/89/89	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	402	SV6	CA-C	-3.35	1.48	1.53
3	AAA	402	SV6	CAZ-CAW	-2.66	1.52	1.55
3	AAA	402	SV6	CAO-CAN	2.38	1.54	1.48
3	AAA	402	SV6	CAI-CAH	-2.19	1.51	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	402	SV6	CAP-CAN-NAA	-6.77	109.64	118.52
3	AAA	402	SV6	CAV-CAW-NAC	4.25	112.55	107.40
3	AAA	402	SV6	O-C-CA	-3.52	115.17	120.95
3	AAA	402	SV6	OBR-CAI-CAM	-3.42	103.49	110.63
3	AAA	402	SV6	CA-C-NAE	2.67	120.40	115.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	402	SV6	CD1-N-CA	-2.67	108.93	112.41
3	AAA	402	SV6	CBN-CBQ-NAG	-2.54	118.64	121.97
3	AAA	402	SV6	CAJ-CAH-NAE	2.53	113.59	110.30
3	AAA	402	SV6	CBP-CBO-NAF	-2.47	118.71	122.19
3	AAA	402	SV6	CAH-NAE-C	-2.29	119.26	123.25
3	AAA	402	SV6	CAI-CAM-NAA	-2.28	113.13	116.22

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	AAA	402	SV6	OBR-CAI-CAM-OBS

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	402	SV6	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	305/306 (99%)	0.77	45 (14%) 5 5	14, 27, 51, 64	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	94	ALA	4.8
1	AAA	305	PHE	4.7
1	AAA	50	LEU	4.5
1	AAA	73	VAL	4.2
1	AAA	154	TYR	3.7
1	AAA	70	ALA	3.6
1	AAA	155	ASP	3.4
1	AAA	223	PHE	3.4
1	AAA	24	THR	3.1
1	AAA	74	GLN	3.0
1	AAA	277	ASN	3.0
1	AAA	35	VAL	2.9
1	AAA	93	THR	2.8
1	AAA	153	ASP	2.8
1	AAA	78	ILE	2.8
1	AAA	294	PHE	2.7
1	AAA	31	TRP	2.7
1	AAA	68	VAL	2.7
1	AAA	249	ILE	2.6
1	AAA	98	THR	2.5
1	AAA	16	CYS	2.5
1	AAA	156	CYS	2.5
1	AAA	75	LEU	2.5
1	AAA	96	PRO	2.5
1	AAA	278	GLY	2.5
1	AAA	304	THR	2.4
1	AAA	224	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	222	ARG	2.4
1	AAA	76	ARG	2.4
1	AAA	91	VAL	2.3
1	AAA	99	PRO	2.3
1	AAA	72	ASN	2.2
1	AAA	101	TYR	2.2
1	AAA	79	GLY	2.2
1	AAA	97	LYS	2.2
1	AAA	77	VAL	2.1
1	AAA	15	GLY	2.0
1	AAA	303	VAL	2.0
1	AAA	67	LEU	2.0
1	AAA	64	HIS	2.0
1	AAA	47	GLU	2.0
1	AAA	235	MET	2.0
1	AAA	152	ILE	2.0
1	AAA	18	VAL	2.0
1	AAA	233	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SV6	AAA	402	49/49	0.94	0.08	19,24,28,37	0
2	CL	AAA	401	1/1	0.97	0.07	22,22,22,22	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.