



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 11:08 PM UTC

PDB ID : 2NC3 / pdb_00002nc3
BMRB ID : 26005
Title : Solution Structure of N-Allosylated Pin1 WW Domain
Authors : Hsu, C.; Park, S.; Mortenson, D.E.; Foley, B.; Wang, X.; Woods, R.J.; Case, D.A.; Powers, E.T.; Wong, C.; Dyson, H.; Kelly, J.W.
Deposited on : 2016-03-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 49%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	33	73% 27%

2 Ensemble composition and analysis

This entry contains 20 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:5-A:28 (24)	0.16	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 6, 7, 9, 10, 11, 12, 13, 14, 17, 19, 20
2	8, 15
3	5, 18
Single-model clusters	3; 16


3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 552 atoms, of which 268 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Pin1 WW Domain.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	33	530	176	257	48	48	1	0

- Molecule 2 is beta-D-allopyranose (CCD ID: ALL) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			
			Total	C	H	O
2	A	1	22	6	11	5

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Pin1 WW Domain




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Pin1 WW Domain



4.2.2 Score per residue for model 2


- Molecule 1: Pin1 WW Domain



4.2.3 Score per residue for model 3

- Molecule 1: Pin1 WW Domain


Chain A:  70% 27%



4.2.4 Score per residue for model 4

- Molecule 1: Pin1 WW Domain


Chain A:  70% 27%



4.2.5 Score per residue for model 5

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.6 Score per residue for model 6

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.7 Score per residue for model 7

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.8 Score per residue for model 8

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.9 Score per residue for model 9

- Molecule 1: Pin1 WW Domain


Chain A:  70% 27%



4.2.10 Score per residue for model 10

- Molecule 1: Pin1 WW Domain

Chain A:  73% 27%



4.2.11 Score per residue for model 11

- Molecule 1: Pin1 WW Domain

Chain A:  70% 27%



4.2.12 Score per residue for model 12

- Molecule 1: Pin1 WW Domain


Chain A:  70% 27%



4.2.13 Score per residue for model 13

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.14 Score per residue for model 14

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.15 Score per residue for model 15

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.16 Score per residue for model 16

- Molecule 1: Pin1 WW Domain


Chain A:  73% 27%



4.2.17 Score per residue for model 17

- Molecule 1: Pin1 WW Domain

Chain A:  70% 27%



4.2.18 Score per residue for model 18

- Molecule 1: Pin1 WW Domain

Chain A:  73% 27%



4.2.19 Score per residue for model 19

- Molecule 1: Pin1 WW Domain

Chain A:  73% 27%



4.2.20 Score per residue for model 20 (medoid)

- Molecule 1: Pin1 WW Domain

Chain A:  73% 27%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	structure solution	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	235
Number of shifts mapped to atoms	235
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	49%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ALL

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.2
All	All	0	1

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	9	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	4300	3880	3660	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	24/33 (73%)	24±0 (99±1%)	0±0 (1±1%)	0±0 (0±0%)	100	100
All	All	480/660 (73%)	477 (99%)	3 (1%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	20/28 (71%)	20±0 (98±2%)	0±0 (2±2%)	55	93
All	All	400/560 (71%)	394 (98%)	6 (2%)	55	93

All 2 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	9	ARG	4
1	A	24	ASN	2

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	ALL	A	101	-	11,11,12	1.40±0.02	1±0 (9±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	ALL	A	101	-	15,15,17	0.75±0.04	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALL	A	101	-	-	0±0,2,19,22	0±0,1,1,1

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	101	ALL	O5-C5	3.21	1.49	1.43	1	20

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 49% for the well-defined parts and 50% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	235
Number of shifts mapped to atoms	235
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	6

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 49%, i.e. 164 atoms were assigned a chemical shift out of a possible 335. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	50/122 (41%)	50/50 (100%)	0/48 (0%)	0/24 (0%)
Sidechain	85/146 (58%)	85/94 (90%)	0/44 (0%)	0/8 (0%)
Aromatic	29/67 (43%)	29/33 (88%)	0/32 (0%)	0/2 (0%)
Overall	164/335 (49%)	164/177 (93%)	0/124 (0%)	0/34 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 50%, i.e. 227 atoms were assigned a chemical shift out of a possible 456. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	65/162 (40%)	65/66 (98%)	0/66 (0%)	0/30 (0%)
Sidechain	133/227 (59%)	133/146 (91%)	0/69 (0%)	0/12 (0%)
Aromatic	29/67 (43%)	29/33 (88%)	0/32 (0%)	0/2 (0%)
Overall	227/456 (50%)	227/245 (93%)	0/167 (0%)	0/44 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	23	THR	HG1	5.08	0.08 – 2.19	18.7
1	A	20	ASN	HB2	-0.71	1.27 – 4.34	-11.5
1	A	9	ARG	HB2	-0.13	0.52 – 3.08	-7.5
1	A	9	ARG	HB3	-0.13	0.43 – 3.11	-7.1
1	A	31	PRO	HG2	-0.07	0.41 – 3.45	-6.6
1	A	20	ASN	HD22	4.13	4.69 – 9.61	-6.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis [i](#)

8.1 Conformationally restricting restraints [i](#)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	482
Intra-residue ($ i-j =0$)	250
Sequential ($ i-j =1$)	95
Medium range ($ i-j >1$ and $ i-j <5$)	39
Long range ($ i-j \geq 5$)	89
Inter-chain	9
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	169
Number of unmapped restraints	0
Number of restraints per residue	19.1
Number of long range restraints per residue ¹	2.6

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations [i](#)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model [i](#)

Distance violations less than 0.1 Å are not included in the calculation. There are no distance violations

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	250	51.9	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	38	7.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	111	23.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	101	21.0	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	95	19.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	37	7.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	47	9.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	11	2.3	0	0.0	0.0	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	39	8.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	2	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	12	2.5	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	25	5.2	0	0.0	0.0	0	0.0	0.0
Long range ($i-j \geq 5$)	89	18.5	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	14	2.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	28	5.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	47	9.8	0	0.0	0.0	0	0.0	0.0
Inter-chain	9	1.9	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	1	0.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	8	1.7	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	482	100.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	92	19.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	206	42.7	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	184	38.2	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfide bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

No violations found

9.3 Distance violation statistics for the ensemble [i](#)

No violations found

9.4 Most violated distance restraints in the ensemble [i](#)

No violations found

9.5 All violated distance restraints [i](#)

No violations found

10 Dihedral-angle violation analysis ⓘ


10.1 Summary of dihedral-angle violations ⓘ

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PSI	29	17.2	0	0.0	0.0	0	0.0	0.0
UNNAMED	81	47.9	0	0.0	0.0	0	0.0	0.0
OMEGA	32	18.9	0	0.0	0.0	0	0.0	0.0
PHI	27	16.0	0	0.0	0.0	0	0.0	0.0
Total	169	100.0	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations ⓘ



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model [i](#)

No violations found

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

No violations found

10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

No violations found

10.5 All violated dihedral-angle restraints [i](#)

No violations found