



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:17 AM UTC

PDB ID : 3NCG / pdb_00003ncg
Title : Activated Calcium-Dependent Protein Kinase 1 from *Cryptosporidium parvum* (CpCDPK1) in complex with bumped kinase inhibitor NM-PP1
Authors : Larson, E.T.; Merritt, E.A.; Medical Structural Genomics of Pathogenic Protozoa (MSGPP)
Deposited on : 2010-06-04
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	486	18% 73% 13% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3530 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calmodulin-domain protein kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	426	Total	C	N	O	S	0	5	0
			3455	2197	576	662	20			


There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	MET	-	expression tag	UNP A3FQ16
A	-15	HIS	-	expression tag	UNP A3FQ16
A	-14	HIS	-	expression tag	UNP A3FQ16
A	-13	HIS	-	expression tag	UNP A3FQ16
A	-12	HIS	-	expression tag	UNP A3FQ16
A	-11	HIS	-	expression tag	UNP A3FQ16
A	-10	HIS	-	expression tag	UNP A3FQ16
A	-9	SER	-	expression tag	UNP A3FQ16
A	-8	SER	-	expression tag	UNP A3FQ16
A	-7	GLY	-	expression tag	UNP A3FQ16
A	-6	ARG	-	expression tag	UNP A3FQ16
A	-5	GLU	-	expression tag	UNP A3FQ16
A	-4	ASN	-	expression tag	UNP A3FQ16
A	-3	LEU	-	expression tag	UNP A3FQ16
A	-2	TYR	-	expression tag	UNP A3FQ16
A	-1	PHE	-	expression tag	UNP A3FQ16
A	0	GLN	-	expression tag	UNP A3FQ16

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		

- Molecule 3 is 1-(1-methylethyl)-3-(naphthalen-1-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine (CCD ID: BK1) (formula: C₁₉H₁₉N₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			24	19	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	48	Total	O	0	0
			48	48		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Calmodulin-domain protein kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.99Å 55.97Å 81.97Å 90.00° 104.78° 90.00°	Depositor
Resolution (Å)	33.58 – 2.49 33.58 – 2.49	Depositor EDS
% Data completeness (in resolution range)	94.8 (33.58-2.49) 95.0 (33.58-2.49)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.48Å)	Xtriage
Refinement program	REFMAC refmac_5.5.0110	Depositor
R, R_{free}	0.250 , 0.305 0.251 , 0.297	Depositor DCC
R_{free} test set	957 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3530	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BK1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/3527	0.79	0/4745

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3455	0	3430	43	0
2	A	3	0	0	0	0
3	A	24	0	19	0	0
4	A	48	0	0	1	0
All	All	3530	0	3449	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:THR:HG21	1:A:456:ILE:HD13	1.67	0.76
1:A:347:SER:OG	1:A:388:ILE:HD11	1.89	0.72
1:A:373:ALA:O	1:A:377:THR:OG1	2.13	0.67
1:A:483:SER:OG	1:A:485:LYS:O	2.19	0.61
1:A:377:THR:HG21	1:A:382:THR:CG2	2.32	0.60
1:A:229:ASN:OD1	1:A:230:THR:HG23	2.02	0.59
1:A:377:THR:HG21	1:A:382:THR:HG21	1.84	0.58
1:A:119:THR:HG23	1:A:122[A]:ARG:NH2	2.21	0.56
1:A:357:GLN:O	1:A:361:GLU:N	2.42	0.53
1:A:471:MET:HE1	1:A:532:GLN:OE1	2.09	0.53
1:A:86:SER:OG	1:A:219:ASP:OD2	2.25	0.52
1:A:269:SER:HB3	1:A:298:TRP:CZ2	2.45	0.52
1:A:215:ILE:C	1:A:215:ILE:HD12	2.35	0.52
1:A:462:ARG:HA	1:A:465:LEU:HB3	1.90	0.52
1:A:140:GLU:OE2	1:A:142:LEU:HD11	2.10	0.52
1:A:524:PHE:O	1:A:528:VAL:HG23	2.09	0.51
1:A:352:MET:HA	1:A:352:MET:HE2	1.91	0.51
1:A:452:TYR:CE2	1:A:456:ILE:HD11	2.46	0.51
1:A:79:VAL:HG22	1:A:94:LYS:HE2	1.92	0.50
1:A:180:LYS:HD3	1:A:336:SER:CB	2.43	0.48
1:A:527:PHE:CE2	1:A:531:LEU:HD11	2.48	0.48
1:A:378:THR:CG2	1:A:379:LEU:N	2.77	0.48
1:A:409:TYR:HE1	1:A:413:MET:HE2	1.78	0.47
1:A:207:GLU:HG2	1:A:379:LEU:HD11	1.97	0.47
1:A:242:ILE:HG23	1:A:246:VAL:HB	1.96	0.47
1:A:373:ALA:HB1	1:A:456:ILE:HG22	1.97	0.46
1:A:442:LEU:HD23	1:A:450:ILE:HG13	1.97	0.46
1:A:411:GLU:OE2	1:A:414:ARG:NH1	2.43	0.45
1:A:70:GLY:O	1:A:74:GLU:HB2	2.17	0.45
1:A:196:HIS:O	1:A:197:ARG:HB2	2.17	0.45
1:A:345:LEU:HD11	1:A:416:LYS:HB2	1.99	0.45
1:A:246:VAL:HG23	1:A:255[B]:CYS:SG	2.57	0.44
1:A:409:TYR:CE1	1:A:413:MET:HE2	2.52	0.44
1:A:348:LEU:HD23	1:A:412:PHE:HD1	1.82	0.44
1:A:242:ILE:HG22	1:A:243:ALA:O	2.19	0.43
1:A:402:ARG:C	1:A:402:ARG:HD3	2.43	0.42
1:A:222:LEU:HD12	4:A:609:HOH:O	2.20	0.42
1:A:344:ASP:O	1:A:348:LEU:HB2	2.20	0.42
1:A:441:LEU:O	1:A:444:MET:HE2	2.19	0.42
1:A:355:ILE:HD11	1:A:459:ALA:HB2	2.03	0.41
1:A:337:SER:O	1:A:339:THR:HG22	2.20	0.41
1:A:378:THR:HG23	1:A:379:LEU:N	2.36	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:TYR:CE2	1:A:456:ILE:CD1	3.04	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/486 (87%)	404 (96%)	18 (4%)	1 (0%)	43 63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	211	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	380/439 (87%)	369 (97%)	11 (3%)	37 65

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	GLU
1	A	123	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	304	ASP
1	A	339	THR
1	A	343	SER
1	A	345	LEU
1	A	347	SER
1	A	352	MET
1	A	377	THR
1	A	378	THR
1	A	463	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	366	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BK1	A	561	-	27,27,27	2.01	5 (18%)	37,39,39	2.10	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BK1	A	561	-	-	0/8/8/8	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	561	BK1	NAX-NAO	-7.62	1.24	1.38
3	A	561	BK1	C4-NAX	-3.73	1.32	1.36
3	A	561	BK1	CAR-NAO	2.81	1.33	1.29
3	A	561	BK1	CAT-CAS	-2.39	1.38	1.43
3	A	561	BK1	C5-CAR	-2.01	1.40	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	561	BK1	C5-C4-N3	-6.66	119.90	126.97
3	A	561	BK1	N3-C4-NAX	5.29	132.94	126.05
3	A	561	BK1	N3-C2-N1	-4.74	121.41	128.58
3	A	561	BK1	C5-CAR-NAO	-4.07	106.11	110.21
3	A	561	BK1	C2-N3-C4	3.82	121.15	111.83

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/486 (87%)	1.24	86 (20%) 3 2	25, 53, 103, 141	5 (1%)

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	THR	6.5
1	A	364	LEU	6.1
1	A	236	ILE	5.8
1	A	371	TYR	5.7
1	A	478	PHE	5.0
1	A	275[A]	TYR	4.7
1	A	465	LEU	4.6
1	A	488	THR	4.5
1	A	484	GLY	4.2
1	A	491	LEU	4.2
1	A	452	TYR	4.0
1	A	486	ILE	4.0
1	A	251	TYR	4.0
1	A	363	LYS	3.9
1	A	420	SER	3.9
1	A	527	PHE	3.9
1	A	-3	LEU	3.8
1	A	365	ALA	3.7
1	A	277	LYS	3.7
1	A	367	ALA	3.7
1	A	379	LEU	3.7
1	A	368	ALA	3.6
1	A	464	ILE	3.5
1	A	475	PHE	3.5
1	A	-2	TYR	3.5
1	A	230	THR	3.5
1	A	463	THR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	360	ALA	3.4
1	A	372	MET	3.3
1	A	417	GLY	3.3
1	A	359	GLN	3.2
1	A	477	MET	3.2
1	A	369	LEU	3.2
1	A	237	GLY	3.2
1	A	355	ILE	3.2
1	A	530	MET	3.1
1	A	345	LEU	3.1
1	A	421	ASN	3.1
1	A	342	ILE	3.0
1	A	376	LEU	3.0
1	A	525	ASN	3.0
1	A	239	ALA	3.0
1	A	-1	PHE	3.0
1	A	462	ARG	2.9
1	A	418	VAL	2.8
1	A	466	LEU	2.8
1	A	250	THR	2.8
1	A	361	GLU	2.8
1	A	446	GLY	2.7
1	A	487	SER	2.7
1	A	377	THR	2.7
1	A	419	ASP	2.6
1	A	339	THR	2.6
1	A	482	GLY	2.6
1	A	71	THR	2.6
1	A	524	PHE	2.6
1	A	375	LYS	2.5
1	A	526	GLU	2.5
1	A	531	LEU	2.5
1	A	77	ASN	2.5
1	A	193	ASN	2.5
1	A	352	MET	2.5
1	A	479	ASP	2.4
1	A	358	PHE	2.4
1	A	416	LYS	2.4
1	A	276	GLY	2.4
1	A	441	LEU	2.4
1	A	533	ASN	2.4
1	A	300	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	398	GLY	2.4
1	A	405	LEU	2.4
1	A	431	ILE	2.3
1	A	284	LYS	2.3
1	A	281	ASP	2.3
1	A	378	THR	2.3
1	A	337	SER	2.2
1	A	70	GLY	2.2
1	A	409	TYR	2.2
1	A	299	ARG	2.2
1	A	383	LYS	2.2
1	A	407	ARG	2.1
1	A	483	SER	2.1
1	A	362	LYS	2.0
1	A	229	ASN	2.0
1	A	489	LYS	2.0
1	A	523	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	A	553	1/1	0.90	0.07	54,54,54,54	1
3	BK1	A	561	24/24	0.92	0.10	26,32,36,38	0
2	CA	A	552	1/1	0.97	0.04	48,48,48,48	0
2	CA	A	551	1/1	0.99	0.04	45,45,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.