



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 10:53 PM UTC

PDB ID : 5NC9 / pdb\_00005nc9  
Title : Crystal structure of the polysaccharide deacetylase Bc1974 from *Bacillus cereus* in complex with (2S)-2,6-diamino-N-hydroxyhexanamide  
Authors : Giastas, P.; Andreou, A.; Eliopoulos, E.E.  
Deposited on : 2017-03-03  
Resolution : 2.44 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
Sidechain outliers	187428	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	 2% 64% 19% 17%
1	B	247	 62% 21% 17%
1	C	247	 2% 69% 14% 17%
1	D	247	 66% 18% 17%

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 6768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidoglycan N-acetylglucosamine deacetylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	Total 1637	C 1049	N 280	O 301	S 7	0	0	0
1	B	206	Total 1645	C 1054	N 281	O 303	S 7	0	0	0
1	C	206	Total 1645	C 1054	N 281	O 303	S 7	0	0	0
1	D	206	Total 1664	C 1066	N 286	O 305	S 7	0	2	0

- Molecule 2 is (2 {S})-2,6-bis(azanyl)- {N}-oxidanyl-hexanamide (CCD ID: 8SZ) (formula: C<sub>6</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 11	C 6	N 3	O 2	0	0
2	B	1	Total 11	C 6	N 3	O 2	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			11	6	3	2		
2	D	1	Total	C	N	O	0	0
			11	6	3	2		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CITRIC ACID (CCD ID: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	B	1	4	2	2	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	D	1	10	6	4	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	25	Total 25	O 25	0	0
7	B	26	Total 26	O 26	0	0
7	C	29	Total 29	O 29	0	0
7	D	22	Total 22	O 22	0	0



- Molecule 1: Peptidoglycan N-acetylglucosamine deacetylase

Chain D:  66% 18% 17%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.77Å 117.83Å 99.06Å 90.00° 102.38° 90.00°	Depositor
Resolution (Å)	48.61 – 2.44 48.61 – 2.44	Depositor EDS
% Data completeness (in resolution range)	96.6 (48.61-2.44) 96.6 (48.61-2.44)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.21 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.220 , 0.279 0.223 , 0.280	Depositor DCC
$R_{free}$ test set	2052 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.4	Xtrriage
Anisotropy	0.739	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 69.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6768	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8SZ, CIT, ZN, PGE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/1682	0.88	0/2279
1	B	0.56	0/1690	0.85	0/2288
1	C	0.53	0/1690	0.92	2/2288 (0.1%)
1	D	0.58	0/1716	0.91	3/2322 (0.1%)
All	All	0.55	0/6778	0.89	5/9177 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	244	LEU	N-CA-C	6.87	118.77	111.28
1	D	239	ALA	CA-C-N	5.94	123.98	120.24
1	D	239	ALA	C-N-CA	5.94	123.98	120.24
1	C	205	VAL	N-CA-C	5.72	116.46	110.62
1	D	172	GLY	N-CA-C	5.64	119.49	112.73

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1637	0	1585	29	1
1	B	1645	0	1600	43	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1645	0	1600	20	0
1	D	1664	0	1624	30	1
2	A	11	0	0	1	0
2	B	11	0	0	0	0
2	C	11	0	0	0	0
2	D	11	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	B	13	0	5	1	0
5	B	4	0	6	1	0
6	D	10	0	14	4	0
7	A	25	0	0	3	0
7	B	26	0	0	5	1
7	C	29	0	0	3	1
7	D	22	0	0	3	0
All	All	6768	0	6434	118	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:161:LYS:NZ	7:D:401:HOH:O	2.04	0.90
5:B:304:EDO:O2	7:B:401:HOH:O	1.99	0.77
1:C:201:ASN:OD1	1:C:235:GLN:NE2	2.19	0.76
1:A:201:ASN:OD1	1:A:235:GLN:NE2	2.19	0.75
1:A:139:GLY:O	7:A:401:HOH:O	2.07	0.72

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:425:HOH:O	7:C:422:HOH:O[2_748]	1.91	0.29
1:A:114:LYS:NZ	1:D:147:GLU:OE2[2_647]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/247 (83%)	198 (97%)	6 (3%)	0	100	100
1	B	204/247 (83%)	195 (96%)	8 (4%)	1 (0%)	24	29
1	C	204/247 (83%)	198 (97%)	6 (3%)	0	100	100
1	D	206/247 (83%)	199 (97%)	7 (3%)	0	100	100
All	All	818/988 (83%)	790 (97%)	27 (3%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	127	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/208 (82%)	169 (99%)	1 (1%)	78	85
1	B	172/208 (83%)	169 (98%)	3 (2%)	53	66
1	C	172/208 (83%)	171 (99%)	1 (1%)	78	85
1	D	175/208 (84%)	173 (99%)	2 (1%)	65	75
All	All	689/832 (83%)	682 (99%)	7 (1%)	68	77

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	187	LYS
1	C	260	GLU
1	D	248	LYS
1	D	183	GLU
1	B	173	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	258	HIS
1	C	117	ASN
1	C	233	HIS
1	D	105	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	EDO	B	304	-	3,3,3	0.55	0	2,2,2	0.20	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	PGE	D	303	-	9,9,9	0.40	0	8,8,8	0.49	0
2	8SZ	D	301	3	9,10,10	0.71	0	9,11,11	1.13	1 (11%)
2	8SZ	C	301	3	9,10,10	0.71	0	9,11,11	1.27	1 (11%)
2	8SZ	A	301	3	9,10,10	0.77	0	9,11,11	1.27	1 (11%)
4	CIT	B	303	-	12,12,12	1.06	0	17,17,17	1.64	5 (29%)
2	8SZ	B	301	3	9,10,10	0.72	0	9,11,11	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	304	-	-	0/1/1/1	-
6	PGE	D	303	-	-	5/7/7/7	-
2	8SZ	D	301	3	-	5/11/11/11	-
2	8SZ	C	301	3	-	3/11/11/11	-
2	8SZ	A	301	3	-	4/11/11/11	-
4	CIT	B	303	-	-	9/16/16/16	-
2	8SZ	B	301	3	-	7/11/11/11	-

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	8SZ	C06-C08-C11	-3.54	103.53	110.76
2	A	301	8SZ	C06-C08-C11	-3.42	103.77	110.76
4	B	303	CIT	O6-C6-C3	3.20	119.27	113.14
2	D	301	8SZ	C08-C11-N05	2.67	120.16	115.93
4	B	303	CIT	O4-C5-C4	2.35	121.78	114.35

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	8SZ	C07-C06-C08-N03
2	B	301	8SZ	C07-C06-C08-C11
4	B	303	CIT	C1-C2-C3-C4
4	B	303	CIT	O7-C3-C6-O5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	B	303	CIT	O7-C3-C6-O6

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	304	EDO	1	0
6	D	303	PGE	4	0
2	A	301	8SZ	1	0
4	B	303	CIT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/247 (83%)	0.51	6 (2%) 53 53	39, 64, 102, 116	0
1	B	206/247 (83%)	0.33	1 (0%) 87 89	38, 57, 82, 100	0
1	C	206/247 (83%)	0.57	6 (2%) 53 53	40, 65, 99, 127	0
1	D	206/247 (83%)	0.36	1 (0%) 87 89	38, 58, 79, 102	2 (0%)
All	All	824/988 (83%)	0.44	14 (1%) 69 70	38, 61, 97, 127	2 (0%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	ASN	3.2
1	C	203	MET	2.8
1	C	68	ARG	2.8
1	B	157	GLY	2.5
1	A	218	THR	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	8SZ	A	301	11/11	0.73	0.23	55,73,82,83	0
2	8SZ	C	301	11/11	0.77	0.21	54,72,81,81	0
2	8SZ	D	301	11/11	0.85	0.18	60,69,75,76	0
4	CIT	B	303	13/13	0.85	0.14	61,79,88,92	0
5	EDO	B	304	4/4	0.85	0.16	53,62,69,71	0
2	8SZ	B	301	11/11	0.86	0.16	57,66,70,73	0
6	PGE	D	303	10/10	0.86	0.17	40,56,62,63	0
3	ZN	A	302	1/1	0.93	0.12	88,88,88,88	0
3	ZN	C	302	1/1	0.97	0.05	70,70,70,70	0
3	ZN	D	302	1/1	0.99	0.05	50,50,50,50	0
3	ZN	B	302	1/1	0.99	0.07	71,71,71,71	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.