



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:56 AM UTC

PDB ID : 1ND6 / pdb_00001nd6
Title : Crystal Structures of Human Prostatic Acid Phosphatase in Complex with a Phosphate Ion and alpha-Benzylaminobenzylphosphonic Acid Update the Mechanistic Picture and Offer New Insights into Inhibitor Design
Authors : Ortlund, E.; LaCount, M.W.; Lebioda, L.
Deposited on : 2002-12-07
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	354	57% 36% . .
1	B	354	61% 31% 5% .
1	C	354	60% 34% . .
1	D	354	59% 34% . .
2	E	4	75% 25%
3	F	2	50% 50%
4	G	5	60% 20% 20%
5	H	3	33% 67%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	I	5	 60% 40%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	E	2	X	-	-	-
2	MAN	E	4	X	-	-	-
3	NAG	F	2	X	-	-	-
4	NAG	G	1	X	-	-	-
4	MAN	G	4	X	-	-	-
4	MAN	G	5	X	-	-	-
5	NAG	H	1	X	-	-	-
5	NAG	H	2	X	-	-	-
5	MAN	H	3	X	-	-	-
6	NAG	I	1	X	-	-	-
9	NAG	B	1401	X	-	-	-

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 12048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called prostatic acid phosphatase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2800	1807	461	516	16			
1	B	343	Total	C	N	O	S	0	0	0
			2807	1811	462	518	16			
1	C	342	Total	C	N	O	S	0	0	0
			2800	1807	461	516	16			
1	D	342	Total	C	N	O	S	0	0	0
			2800	1807	461	516	16			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	4	Total	C	N	O	0	0	0
			54	28	2	24			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			30	16	2	12			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	5	Total	C	N	O	0	0	0
			66	34	2	30			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	3	Total	C	N	O	0	0	0
			42	22	2	18			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	I	5	Total	C	N	O	0	0	0
			66	34	2	30			

- Molecule 7 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		
7	B	1	Total	O	P	0	0
			5	4	1		
7	C	1	Total	O	P	0	0
			5	4	1		
7	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 8 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C₁₀H₂₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			16	10	6		
8	B	1	Total	C	O	0	0
			16	10	6		
8	C	1	Total	C	O	0	0
			16	10	6		
8	D	1	Total	C	O	0	0
			16	10	6		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	N	O	0	0
			15	8	1	6		
9	D	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 10 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	B	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 11 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	143	Total	O	0	0
			143	143		
11	B	131	Total	O	0	0
			131	131		
11	C	93	Total	O	0	0
			93	93		
11	D	97	Total	O	0	0
			97	97		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: prostatic acid phosphatase




- Molecule 1: prostatic acid phosphatase




- Molecule 1: prostatic acid phosphatase





- Molecule 1: prostatic acid phosphatase

Chain D: 59% 34%



- Molecule 2: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 75% 25%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 50% 50%



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  60% 20% 20%



- Molecule 5: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  33% 67%



- Molecule 6: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  60% 40%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.89Å 203.32Å 70.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.27 – 2.40	Depositor
% Data completeness (in resolution range)	90.6 (35.27-2.40)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.198 , 0.256	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	12048	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, MAN, NAG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2882	0.92	11/3914 (0.3%)
1	B	0.40	0/2889	0.92	11/3924 (0.3%)
1	C	0.40	0/2882	0.91	13/3914 (0.3%)
1	D	0.41	0/2882	0.91	6/3914 (0.2%)
All	All	0.40	0/11535	0.91	41/15666 (0.3%)

There are no bond length outliers.

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2040	LEU	N-CA-C	-6.69	104.07	111.36
1	B	1128	CYS	CA-C-N	6.58	126.08	119.24
1	B	1128	CYS	C-N-CA	6.58	126.08	119.24
1	D	3046	TYR	N-CA-C	-6.39	104.40	111.36
1	A	75	THR	N-CA-C	-6.35	102.15	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2800	0	2749	115	0
1	B	2807	0	2753	126	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2800	0	2747	122	0
1	D	2800	0	2746	123	0
2	E	54	0	47	4	0
3	F	30	0	27	1	0
4	G	66	0	57	2	0
5	H	42	0	37	3	0
6	I	66	0	59	1	0
7	A	5	0	0	1	0
7	B	5	0	0	1	0
7	C	5	0	0	1	0
7	D	5	0	0	1	0
8	A	16	0	22	1	0
8	B	16	0	22	3	0
8	C	16	0	22	2	0
8	D	16	0	22	2	0
9	B	15	0	14	1	0
9	D	15	0	14	0	0
10	B	5	0	2	3	0
11	A	143	0	0	7	0
11	B	131	0	0	7	0
11	C	93	0	0	3	0
11	D	97	0	0	4	0
All	All	12048	0	11340	466	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 466 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1304:HIS:HB3	10:B:1403:GLY:OXT	1.47	1.12
1:C:2310:MET:HE3	1:C:2316:PRO:HD3	1.25	1.06
1:B:1146:LYS:HE3	1:B:1146:LYS:HA	1.38	1.03
1:B:1127:ASN:HD22	1:B:1127:ASN:H	1.02	0.97
1:C:2127:ASN:HD22	1:C:2127:ASN:H	1.07	0.97

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/354 (96%)	319 (94%)	19 (6%)	2 (1%)	21	32
1	B	341/354 (96%)	311 (91%)	28 (8%)	2 (1%)	21	32
1	C	340/354 (96%)	310 (91%)	28 (8%)	2 (1%)	21	32
1	D	340/354 (96%)	318 (94%)	19 (6%)	3 (1%)	14	22
All	All	1361/1416 (96%)	1258 (92%)	94 (7%)	9 (1%)	18	28

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	SER
1	B	1064	TYR
1	C	2289	LYS
1	D	3064	TYR
1	A	65	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/325 (97%)	290 (92%)	24 (8%)	12	21
1	B	315/325 (97%)	299 (95%)	16 (5%)	21	37
1	C	314/325 (97%)	299 (95%)	15 (5%)	23	40
1	D	314/325 (97%)	301 (96%)	13 (4%)	27	46
All	All	1257/1300 (97%)	1189 (95%)	68 (5%)	20	35

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	3006	THR
1	D	3067	GLU
1	D	3211	LEU
1	B	1005	VAL
1	A	328	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	2145	GLN
1	D	3086	ASN
1	C	2186	HIS
1	C	2303	GLN
1	D	3127	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

19 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	1,2	15,15,15	0.48	0	21,21,21	0.98	1 (4%)
2	NAG	E	2	2	15,15,15	0.51	0	21,21,21	0.58	0
2	MAN	E	3	2	12,12,12	0.50	0	17,17,17	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MAN	E	4	2	12,12,12	0.40	0	17,17,17	0.62	0
3	NAG	F	1	1,3	15,15,15	0.41	0	21,21,21	0.92	2 (9%)
3	NAG	F	2	3	15,15,15	0.40	0	21,21,21	0.62	0
4	NAG	G	1	1,4	15,15,15	0.48	0	21,21,21	0.72	0
4	NAG	G	2	4	15,15,15	0.58	0	21,21,21	0.98	1 (4%)
4	MAN	G	3	4	12,12,12	0.51	0	17,17,17	0.60	0
4	MAN	G	4	4	12,12,12	0.39	0	17,17,17	0.43	0
4	MAN	G	5	4	12,12,12	0.40	0	17,17,17	0.39	0
5	NAG	H	1	1,5	15,15,15	0.53	0	21,21,21	1.17	1 (4%)
5	NAG	H	2	5	15,15,15	0.80	0	21,21,21	1.31	2 (9%)
5	MAN	H	3	5	12,12,12	0.50	0	17,17,17	0.39	0
6	NAG	I	1	6,1	15,15,15	0.44	0	21,21,21	0.66	0
6	NAG	I	2	6	15,15,15	0.45	0	21,21,21	0.99	1 (4%)
6	MAN	I	3	6	12,12,12	0.46	0	17,17,17	0.46	0
6	MAN	I	4	6	12,12,12	0.49	0	17,17,17	0.40	0
6	MAN	I	5	6	12,12,12	0.33	0	17,17,17	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	0/6/26/26	0/1/1/1
2	NAG	E	2	2	1/1/6/7	4/6/26/26	0/1/1/1
2	MAN	E	3	2	-	2/2/22/22	0/1/1/1
2	MAN	E	4	2	1/1/5/5	2/2/22/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/26/26	0/1/1/1
3	NAG	F	2	3	1/1/6/7	2/6/26/26	0/1/1/1
4	NAG	G	1	1,4	1/1/6/7	0/6/26/26	0/1/1/1
4	NAG	G	2	4	-	2/6/26/26	0/1/1/1
4	MAN	G	3	4	-	2/2/22/22	0/1/1/1
4	MAN	G	4	4	1/1/5/5	0/2/22/22	0/1/1/1
4	MAN	G	5	4	1/1/5/5	2/2/22/22	0/1/1/1
5	NAG	H	1	1,5	1/1/6/7	0/6/26/26	0/1/1/1
5	NAG	H	2	5	1/1/6/7	6/6/26/26	0/1/1/1
5	MAN	H	3	5	1/1/5/5	2/2/22/22	0/1/1/1
6	NAG	I	1	6,1	1/1/6/7	0/6/26/26	0/1/1/1
6	NAG	I	2	6	-	0/6/26/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MAN	I	3	6	-	0/2/22/22	0/1/1/1
6	MAN	I	4	6	-	0/2/22/22	0/1/1/1
6	MAN	I	5	6	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	2	NAG	C1-C2-C3	-4.02	105.06	110.54
6	I	2	NAG	O1-C1-C2	-3.46	102.03	109.22
4	G	2	NAG	O1-C1-C2	-3.31	102.33	109.22
5	H	1	NAG	C4-C3-C2	3.31	115.22	110.40
2	E	1	NAG	O1-C1-C2	-2.76	103.48	109.22

5 of 10 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	E	2	NAG	C1
2	E	4	MAN	C1
3	F	2	NAG	C1
4	G	1	NAG	C1
4	G	4	MAN	C1

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	H	2	NAG	C1-C2-N2-C7
2	E	2	NAG	C8-C7-N2-C2
2	E	2	NAG	O7-C7-N2-C2
4	G	3	MAN	O5-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6

There are no ring outliers.

11 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	2	NAG	2	0
2	E	4	MAN	1	0
5	H	3	MAN	1	0
4	G	3	MAN	2	0
6	I	1	NAG	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	2	NAG	1	0
5	H	1	NAG	2	0
3	F	1	NAG	1	0
2	E	1	NAG	3	0
2	E	2	NAG	2	0
2	E	3	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	1PE	A	402	-	15,15,15	0.79	0	14,14,14	1.50	4 (28%)
8	1PE	B	1404	-	15,15,15	0.75	0	14,14,14	1.50	4 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	GLY	B	1403	-	4,4,4	1.32	1 (25%)	3,4,4	0.34	0
9	NAG	B	1401	1	15,15,15	0.40	0	21,21,21	0.81	0
8	1PE	C	2402	-	15,15,15	0.79	0	14,14,14	1.48	4 (28%)
8	1PE	D	3403	-	15,15,15	0.80	0	14,14,14	1.48	4 (28%)
7	PO4	A	401	-	4,4,4	1.66	0	6,6,6	0.48	0
9	NAG	D	3401	1	15,15,15	0.46	0	21,21,21	1.00	1 (4%)
7	PO4	D	3402	-	4,4,4	1.56	0	6,6,6	0.46	0
7	PO4	B	1402	-	4,4,4	1.71	0	6,6,6	0.46	0
7	PO4	C	2401	-	4,4,4	1.65	1 (25%)	6,6,6	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	1PE	A	402	-	-	6/13/13/13	-
8	1PE	B	1404	-	-	6/13/13/13	-
10	GLY	B	1403	-	-	2/2/2/2	-
9	NAG	B	1401	1	1/1/6/7	4/6/26/26	0/1/1/1
8	1PE	C	2402	-	-	1/13/13/13	-
8	1PE	D	3403	-	-	6/13/13/13	-
9	NAG	D	3401	1	-	4/6/26/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	1403	GLY	OXT-C	-2.52	1.22	1.30
7	C	2401	PO4	P-O4	-2.01	1.48	1.54

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	402	1PE	C25-OH5-C14	2.68	125.01	113.26
8	C	2402	1PE	C25-OH5-C14	2.67	124.95	113.26
8	B	1404	1PE	C25-OH5-C14	2.65	124.85	113.26
8	D	3403	1PE	C25-OH5-C14	2.62	124.72	113.26
8	B	1404	1PE	OH6-C26-C16	2.54	121.31	110.11

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
9	B	1401	NAG	C1

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	D	3401	NAG	C8-C7-N2-C2
9	D	3401	NAG	O7-C7-N2-C2
10	B	1403	GLY	O-C-CA-N
10	B	1403	GLY	OXT-C-CA-N
9	D	3401	NAG	O5-C5-C6-O6

There are no ring outliers.

10 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	402	1PE	1	0
8	B	1404	1PE	3	0
10	B	1403	GLY	3	0
9	B	1401	NAG	1	0
8	C	2402	1PE	2	0
8	D	3403	1PE	2	0
7	A	401	PO4	1	0
7	D	3402	PO4	1	0
7	B	1402	PO4	1	0
7	C	2401	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.