



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:54 PM UTC

PDB ID : 1NFT / pdb_00001nft
Title : OVOTRANSFERRIN, N-TERMINAL LOBE, IRON LOADED OPEN FORM
Authors : Mizutani, K.; Yamashita, H.; Kurokawa, H.; Mikami, B.; Hirose, M.
Deposited on : 1999-01-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	329	 75% 20% 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2725 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (OVOTRANSFERRIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2543	1599	437	492	15			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	VAL	ILE	conflict	UNP P02789
A	64	VAL	ALA	conflict	UNP P02789
A	133	ILE	LEU	conflict	UNP P02789

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is NITRILOTRIACETIC ACID (CCD ID: NTA) (formula: C₆H₉NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			13	6	1	6		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	153	Total	O	0	0
			153	153		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROTEIN (OVOTRANSFERRIN)



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	125.29Å 125.29Å 87.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	7.00 – 2.10	Depositor
% Data completeness (in resolution range)	88.3 (7.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.189 , 0.256	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2725	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE, SO4, NTA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	9/2596 (0.3%)	1.75	52/3510 (1.5%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	283	HIS	CD2-NE2	-7.11	1.30	1.37
1	A	134	HIS	CD2-NE2	-6.94	1.30	1.37
1	A	210	HIS	CD2-NE2	-6.58	1.30	1.37
1	A	84	HIS	CD2-NE2	-6.54	1.30	1.37
1	A	250	HIS	CD2-NE2	-6.49	1.30	1.37
1	A	116	HIS	CD2-NE2	-6.32	1.30	1.37
1	A	196	HIS	CD2-NE2	-5.26	1.32	1.37
1	A	283	HIS	CG-ND1	-5.10	1.32	1.38
1	A	46	ILE	CA-CB	5.00	1.60	1.54

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	257	ASP	CA-CB-CG	10.30	122.91	112.60
1	A	145	SER	N-CA-C	-8.76	92.14	110.80
1	A	86	GLU	N-CA-C	7.91	120.90	111.02
1	A	28	GLN	OE1-CD-NE2	-7.70	114.90	122.60
1	A	83	GLU	N-CA-C	7.62	119.67	111.36
1	A	60	ASP	CA-CB-CG	7.13	119.73	112.60
1	A	84	HIS	CB-CG-CD2	-7.04	122.04	131.20
1	A	176	GLY	O-C-N	-6.84	117.62	123.43
1	A	279	LYS	N-CA-C	-6.79	104.62	113.17
1	A	329	GLN	OE1-CD-NE2	-6.78	115.82	122.60
1	A	22	ASN	OD1-CG-ND2	-6.78	115.82	122.60
1	A	146	GLY	N-CA-C	-6.73	101.34	110.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	ASN	CA-CB-CG	-6.68	105.92	112.60
1	A	102	THR	O-C-N	-6.67	115.18	122.79
1	A	315	ASP	CA-CB-CG	6.27	118.87	112.60
1	A	21	ASN	OD1-CG-ND2	-6.16	116.44	122.60
1	A	205	VAL	CB-CA-C	-6.01	101.25	110.50
1	A	221	LYS	CB-CG-CD	-5.90	97.73	111.30
1	A	290	LYS	N-CA-C	-5.87	100.27	109.25
1	A	264	TRP	CG-CD2-CE3	5.78	139.68	133.90
1	A	30	GLU	N-CA-C	-5.77	101.47	110.42
1	A	297	ASP	CA-CB-CG	5.75	118.35	112.60
1	A	144	GLU	CA-C-N	5.72	132.46	121.54
1	A	144	GLU	C-N-CA	5.72	132.46	121.54
1	A	257	ASP	O-C-N	-5.71	117.58	123.29
1	A	283	HIS	CB-CG-CD2	-5.70	123.78	131.20
1	A	291	LYS	CB-CG-CD	-5.69	98.22	111.30
1	A	210	HIS	CB-CG-CD2	-5.69	123.81	131.20
1	A	144	GLU	N-CA-C	-5.65	97.35	107.98
1	A	84	HIS	CB-CG-ND1	5.61	131.12	122.70
1	A	238	ASN	OD1-CG-ND2	-5.59	117.01	122.60
1	A	185	ASN	OD1-CG-ND2	-5.59	117.01	122.60
1	A	169	LYS	CA-CB-CG	5.46	125.02	114.10
1	A	309	ARG	N-CA-CB	-5.45	101.72	109.85
1	A	214	ASN	CA-CB-CG	-5.42	107.18	112.60
1	A	264	TRP	CB-CG-CD1	-5.41	118.79	126.90
1	A	145	SER	CA-CB-OG	5.37	121.84	111.10
1	A	144	GLU	O-C-N	5.37	129.41	122.43
1	A	329	GLN	CG-CD-NE2	5.32	124.38	116.40
1	A	250	HIS	CB-CG-ND1	5.26	130.59	122.70
1	A	22	ASN	CB-CG-ND2	5.24	124.25	116.40
1	A	127	ILE	CA-C-N	5.16	124.34	118.97
1	A	127	ILE	C-N-CA	5.16	124.34	118.97
1	A	322	PHE	N-CA-C	5.14	116.57	111.07
1	A	234	GLN	CA-C-N	5.09	125.02	119.78
1	A	234	GLN	C-N-CA	5.09	125.02	119.78
1	A	102	THR	N-CA-C	-5.08	103.83	110.53
1	A	221	LYS	CB-CA-C	-5.07	102.38	110.79
1	A	116	HIS	CB-CG-CD2	-5.06	124.62	131.20
1	A	172	ARG	N-CA-C	5.05	117.51	111.71
1	A	261	GLU	CB-CG-CD	5.03	121.15	112.60
1	A	121	ARG	CA-CB-CG	-5.02	104.06	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2543	0	2485	26	0
2	A	15	0	0	0	0
3	A	1	0	0	0	0
4	A	13	0	0	0	0
5	A	153	0	0	3	0
All	All	2725	0	2485	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:LEU:HG	1:A:63:GLN:HB3	1.70	0.74
1:A:271:GLN:HE22	1:A:307:LEU:H	1.41	0.69
1:A:271:GLN:HE22	1:A:307:LEU:N	1.94	0.64
1:A:210:HIS:HE1	5:A:482:HOH:O	1.85	0.59
1:A:214:ASN:HD21	1:A:221:LYS:HE2	1.69	0.57
1:A:271:GLN:NE2	1:A:307:LEU:H	2.06	0.53
1:A:294:VAL:HG13	1:A:295:LEU:HD13	1.90	0.52
1:A:6:VAL:HG13	1:A:35:THR:HG21	1.92	0.51
1:A:93:TYR:H	1:A:210:HIS:CD2	2.34	0.46
1:A:100:LYS:HE2	1:A:225:GLU:CG	2.45	0.46
1:A:6:VAL:HG13	1:A:35:THR:CG2	2.46	0.46
1:A:9:TRP:HB2	1:A:34:LEU:HD11	2.00	0.44
1:A:100:LYS:HE2	1:A:225:GLU:HG2	1.99	0.44
1:A:196:HIS:HD2	5:A:443:HOH:O	2.01	0.44
1:A:167:GLU:HB2	1:A:170:LEU:HG	2.00	0.44
1:A:24:ARG:HH21	1:A:25:ASP:CG	2.26	0.43
1:A:184:ARG:NE	1:A:184:ARG:H	2.16	0.42
1:A:93:TYR:H	1:A:210:HIS:HD2	1.65	0.42
1:A:94:ALA:HB1	5:A:420:HOH:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:VAL:HG21	1:A:308:LYS:HG3	2.02	0.41
1:A:8:ARG:HB2	1:A:54:ALA:HA	2.02	0.41
1:A:22:ASN:O	1:A:26:LEU:HD13	2.20	0.41
1:A:299:LEU:HD12	1:A:299:LEU:HA	1.91	0.41
1:A:150:GLN:HG2	1:A:167:GLU:OE2	2.21	0.41
1:A:214:ASN:ND2	1:A:221:LYS:HE2	2.35	0.41
1:A:24:ARG:HG3	1:A:34:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	327/329 (99%)	310 (95%)	15 (5%)	2 (1%)	21 18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	SER
1	A	122	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	275/275 (100%)	251 (91%)	24 (9%)	9 7

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	SER
1	A	6	VAL
1	A	28	GLN
1	A	30	GLU
1	A	31	ARG
1	A	35	THR
1	A	59	LEU
1	A	71	PRO
1	A	85	THR
1	A	91	SER
1	A	119	LEU
1	A	139	GLU
1	A	179	LYS
1	A	184	ARG
1	A	205	VAL
1	A	214	ASN
1	A	236	VAL
1	A	267	LEU
1	A	295	LEU
1	A	298	LEU
1	A	299	LEU
1	A	309	ARG
1	A	311	PRO
1	A	318	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	ASN
1	A	126	ASN
1	A	134	HIS
1	A	210	HIS
1	A	214	ASN
1	A	238	ASN
1	A	243	ASN
1	A	271	GLN
1	A	283	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	329	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	354	-	4,4,4	0.18	0	6,6,6	0.13	0
2	SO4	A	353	-	4,4,4	0.40	0	6,6,6	0.36	0
4	NTA	A	334	3	12,12,12	3.30	7 (58%)	15,15,15	2.10	5 (33%)
2	SO4	A	351	-	4,4,4	0.33	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NTA	A	334	3	-	3/12/12/12	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	334	NTA	CA-C	-5.54	1.40	1.51
4	A	334	NTA	C6-C7	-5.38	1.40	1.51
4	A	334	NTA	CA-N	4.14	1.55	1.47
4	A	334	NTA	C10-N	3.88	1.55	1.47
4	A	334	NTA	OXT-C	-3.78	1.18	1.30
4	A	334	NTA	O8-C7	-2.97	1.21	1.30
4	A	334	NTA	C6-N	2.89	1.53	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	334	NTA	CA-N-C10	5.02	123.43	111.66
4	A	334	NTA	C10-N-C6	3.13	119.00	111.66
4	A	334	NTA	O9-C7-C6	-2.41	112.32	122.38
4	A	334	NTA	O13-C11-C10	-2.32	112.68	122.38
4	A	334	NTA	O12-C11-O13	2.22	129.03	123.33

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	334	NTA	C7-C6-N-C10
4	A	334	NTA	C-CA-N-C6
4	A	334	NTA	C11-C10-N-C6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.