



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 11:22 AM UTC

PDB ID : 6NFG / pdb_00006nfg
Title : CYCLIC GMP-AMP SYNTHASE in complex with compound 16 inhibitor: 7-hydroxy-N-methyl-5-phenylpyrazolo[1,5-a]pyrimidine-3-carboxamide
Authors : Hall, J.
Deposited on : 2018-12-20
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	3% 81% 14% • •
1	B	362	4% 82% 12% • •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5622 atoms, of which 24 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYCLIC GMP-AMP SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	348	Total	C	N	O	S	0	0	0
			2730	1754	469	492	15			
1	B	346	Total	C	N	O	S	0	0	0
			2649	1701	451	482	15			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 7-hydroxy-N-methyl-5-phenylpyrazolo[1,5-a]pyrimidine-3-carboxamide (CCD ID: KKP) (formula: C₁₄H₁₂N₄O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			32	14	12	4	2		
3	B	1	Total	C	H	N	O	0	0
			32	14	12	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	94	Total	O	0	0
			94	94		
4	B	83	Total	O	0	0
			83	83		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYCLIC GMP-AMP SYNTHASE



• Molecule 1: CYCLIC GMP-AMP SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	217.15Å 45.60Å 86.82Å 90.00° 105.69° 90.00°	Depositor
Resolution (Å)	104.53 – 2.76 104.53 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.5 (104.53-2.76) 99.9 (104.53-2.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.77Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.182 , 0.244 0.194 , 0.261	Depositor DCC
R_{free} test set	1076 reflections (4.23%)	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.693	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 75.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5622	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KKP, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2783 (0.0%)	1.40	20/3752 (0.5%)
1	B	0.85	0/2702	1.38	15/3654 (0.4%)
All	All	0.85	1/5485 (0.0%)	1.39	35/7406 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	234	VAL	CA-C	5.27	1.56	1.52

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	313	SER	N-CA-C	-10.13	92.91	108.31
1	B	313	SER	N-CA-C	-7.73	94.33	110.80
1	A	253	PHE	CA-C-N	7.71	135.58	121.70
1	A	253	PHE	C-N-CA	7.71	135.58	121.70
1	A	234	VAL	CB-CA-C	7.24	117.16	110.13
1	B	235	PRO	N-CA-C	7.11	127.12	112.47
1	A	235	PRO	N-CA-C	6.95	126.78	112.47
1	A	346	ALA	CA-C-N	5.95	128.58	120.54
1	A	346	ALA	C-N-CA	5.95	128.58	120.54
1	B	468	VAL	CA-C-N	5.89	128.09	120.44
1	B	468	VAL	C-N-CA	5.89	128.09	120.44
1	A	468	VAL	CA-C-N	5.77	127.94	120.44
1	A	468	VAL	C-N-CA	5.77	127.94	120.44
1	A	216	GLU	CA-C-N	5.49	129.93	122.19
1	A	216	GLU	C-N-CA	5.49	129.93	122.19
1	A	390	HIS	N-CA-C	5.43	120.01	113.17
1	A	167	ALA	CA-C-N	5.22	127.62	120.46
1	A	167	ALA	C-N-CA	5.22	127.62	120.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	456	ASP	CA-C-N	5.17	127.46	120.38
1	A	456	ASP	C-N-CA	5.17	127.46	120.38
1	B	456	ASP	CA-C-N	5.16	127.45	120.38
1	B	456	ASP	C-N-CA	5.16	127.45	120.38
1	B	167	ALA	CA-C-N	5.13	127.49	120.46
1	B	167	ALA	C-N-CA	5.13	127.49	120.46
1	A	314	GLU	N-CA-CB	-5.12	104.49	111.65
1	B	346	ALA	CA-C-N	5.08	127.05	120.44
1	B	346	ALA	C-N-CA	5.08	127.05	120.44
1	A	461	GLY	CA-C-N	5.07	127.49	120.29
1	A	461	GLY	C-N-CA	5.07	127.49	120.29
1	B	390	HIS	N-CA-C	5.05	119.53	113.17
1	B	429	HIS	CA-C-N	5.04	128.66	120.60
1	B	429	HIS	C-N-CA	5.04	128.66	120.60
1	A	363	HIS	CA-CB-CG	5.01	118.81	113.80
1	B	506	LYS	CA-C-N	5.01	126.95	120.44
1	B	506	LYS	C-N-CA	5.01	126.95	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2730	0	2655	14	0
1	B	2649	0	2485	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	20	12	0	2	0
3	B	20	12	0	1	0
4	A	94	0	0	0	0
4	B	83	0	0	1	0
All	All	5598	24	5140	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:ILE:HD11	1:B:411:LYS:HA	1.64	0.79
1:B:336:GLU:HG2	1:B:486:PRO:HG2	1.75	0.68
1:A:234:VAL:HG23	1:A:237:ILE:HG21	1.77	0.66
1:A:234:VAL:CG2	1:A:237:ILE:HG21	2.32	0.59
1:A:234:VAL:HG23	1:A:237:ILE:CG2	2.35	0.56
1:B:482:ASN:ND2	3:B:602:KKP:C07	2.74	0.50
1:A:284:ILE:HG21	1:A:308:VAL:HG11	1.95	0.49
1:A:488:PHE:CE1	3:A:602:KKP:C08	2.95	0.49
1:B:220:ILE:HD11	1:B:411:LYS:CA	2.40	0.49
1:B:284:ILE:HG21	1:B:308:VAL:HG11	1.95	0.48
1:B:218:VAL:O	1:B:220:ILE:HD12	2.14	0.47
1:B:234:VAL:HG12	1:B:262:LEU:HD11	1.96	0.47
1:B:481:GLU:HG2	4:B:752:HOH:O	2.15	0.47
1:A:430:LEU:HD22	1:A:504:LEU:HD13	1.96	0.47
1:B:511:GLU:OE1	1:B:517:PRO:HD2	2.15	0.47
1:A:488:PHE:CD1	3:A:602:KKP:C08	2.99	0.46
1:B:263:SER:HA	1:B:266:LEU:HD12	1.98	0.46
1:B:311:LEU:HD23	1:B:314:GLU:HA	1.98	0.45
1:B:496:ILE:HB	1:B:501:LYS:HE3	1.98	0.45
1:A:263:SER:HA	1:A:266:LEU:HD12	1.99	0.45
1:A:511:GLU:OE1	1:A:517:PRO:HD2	2.16	0.45
1:B:233:GLU:HA	1:B:325:GLU:HB3	1.98	0.45
1:A:203:PHE:HE2	1:A:279:LYS:HG2	1.83	0.44
1:A:209:LEU:HD23	1:A:209:LEU:HA	1.92	0.43
1:A:216:GLU:O	1:A:218:VAL:HG23	2.19	0.43
1:A:496:ILE:HB	1:A:501:LYS:HE3	2.00	0.42
1:A:218:VAL:HG12	1:A:218:VAL:O	2.20	0.41
1:B:216:GLU:HG2	1:B:384:LYS:HA	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/362 (94%)	320 (94%)	18 (5%)	2 (1%)	21	38
1	B	336/362 (93%)	312 (93%)	22 (6%)	2 (1%)	21	38
All	All	676/724 (93%)	632 (94%)	40 (6%)	4 (1%)	21	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	345	SER
1	B	345	SER
1	A	235	PRO
1	B	235	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/334 (86%)	260 (91%)	27 (9%)	8	16
1	B	268/334 (80%)	246 (92%)	22 (8%)	10	20
All	All	555/668 (83%)	506 (91%)	49 (9%)	9	18

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	THR
1	A	196	ARG
1	A	208	LEU
1	A	209	LEU
1	A	213	SER
1	A	229	MET
1	A	233	GLU
1	A	246	ARG
1	A	274	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	276	MET
1	A	313	SER
1	A	315	LYS
1	A	316	ILE
1	A	341	GLN
1	A	347	LYS
1	A	359	LEU
1	A	371	GLN
1	A	385	GLU
1	A	386	ILE
1	A	432	LYS
1	A	438	VAL
1	A	457	ARG
1	A	462	LEU
1	A	499	ARG
1	A	506	LYS
1	A	507	GLN
1	A	518	VAL
1	B	181	THR
1	B	209	LEU
1	B	220	ILE
1	B	227	ASP
1	B	229	MET
1	B	246	ARG
1	B	274	SER
1	B	276	MET
1	B	311	LEU
1	B	316	ILE
1	B	336	GLU
1	B	347	LYS
1	B	359	LEU
1	B	385	GLU
1	B	386	ILE
1	B	425	LYS
1	B	429	HIS
1	B	438	VAL
1	B	457	ARG
1	B	462	LEU
1	B	506	LYS
1	B	518	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	ASN
1	A	224	ASN
1	A	289	ASN
1	A	341	GLN
1	A	451	GLN
1	A	466	ASN
1	B	210	ASN
1	B	342	ASN
1	B	371	GLN
1	B	451	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	KKP	B	602	-	21,22,22	2.71	9 (42%)	25,31,31	2.27	9 (36%)
3	KKP	A	602	-	21,22,22	2.73	9 (42%)	25,31,31	2.64	13 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KKP	B	602	-	-	4/10/10/10	0/3/3/3
3	KKP	A	602	-	-	0/10/10/10	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	KKP	C14-N15	9.22	1.46	1.33
3	B	602	KKP	C14-N15	9.08	1.46	1.33
3	B	602	KKP	C18-N19	4.29	1.39	1.32
3	A	602	KKP	C18-N19	4.21	1.39	1.32
3	A	602	KKP	C12-N20	-3.37	1.33	1.38
3	B	602	KKP	C05-C04	3.32	1.52	1.47
3	A	602	KKP	C18-C13	-2.92	1.37	1.41
3	B	602	KKP	C18-C13	-2.76	1.37	1.41
3	A	602	KKP	C05-C04	2.67	1.51	1.47
3	A	602	KKP	O01-C02	2.56	1.40	1.32
3	B	602	KKP	C12-N20	-2.49	1.34	1.38
3	B	602	KKP	O17-C14	-2.47	1.18	1.23
3	B	602	KKP	O01-C02	2.47	1.40	1.32
3	B	602	KKP	C03-C04	-2.43	1.38	1.43
3	A	602	KKP	C03-C04	-2.36	1.38	1.43
3	A	602	KKP	O17-C14	-2.28	1.19	1.23
3	B	602	KKP	C12-N11	2.14	1.39	1.34
3	A	602	KKP	C13-C12	-2.03	1.40	1.45

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	KKP	C13-C12-N20	6.27	108.87	105.58
3	A	602	KKP	C18-N19-N20	6.04	107.68	103.43
3	B	602	KKP	C18-N19-N20	5.72	107.45	103.43
3	B	602	KKP	C18-C13-C12	4.35	107.81	104.57
3	A	602	KKP	C13-C18-N19	-3.94	107.56	112.45
3	B	602	KKP	C13-C18-N19	-3.79	107.75	112.45
3	B	602	KKP	C13-C12-N20	3.60	107.47	105.58
3	A	602	KKP	C18-C13-C12	3.57	107.23	104.57
3	A	602	KKP	C12-N20-N19	-3.40	108.65	112.24
3	B	602	KKP	C13-C12-N11	3.06	131.86	123.54
3	B	602	KKP	C05-C04-N11	2.94	120.07	116.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	KKP	C10-C05-C04	-2.87	117.66	120.87
3	B	602	KKP	C12-N20-N19	-2.60	109.50	112.24
3	A	602	KKP	C13-C12-N11	2.58	130.55	123.54
3	B	602	KKP	O01-C02-C03	-2.52	120.29	125.58
3	A	602	KKP	C04-N11-C12	2.51	120.35	117.78
3	A	602	KKP	O01-C02-C03	-2.39	120.56	125.58
3	A	602	KKP	C03-C04-N11	-2.29	119.65	121.83
3	A	602	KKP	C16-N15-C14	-2.14	119.71	121.90
3	B	602	KKP	C03-C04-N11	-2.14	119.79	121.83
3	A	602	KKP	C06-C05-C10	2.11	121.25	118.57
3	A	602	KKP	C09-C08-C07	2.09	122.73	119.87

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	602	KKP	C12-C13-C14-N15
3	B	602	KKP	C12-C13-C14-O17
3	B	602	KKP	C18-C13-C14-O17
3	B	602	KKP	C18-C13-C14-N15


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	KKP	1	0
3	A	602	KKP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	348/362 (96%)	0.09	10 (2%) 53 52	24, 45, 77, 108	0
1	B	346/362 (95%)	0.27	13 (3%) 44 41	22, 53, 87, 109	0
All	All	694/724 (95%)	0.18	23 (3%) 49 47	22, 49, 84, 109	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	LYS	4.4
1	A	366	GLU	3.3
1	B	220	ILE	3.0
1	B	429	HIS	2.8
1	B	295	ASP	2.8
1	B	370	PHE	2.8
1	A	235	PRO	2.7
1	B	308	VAL	2.7
1	B	235	PRO	2.7
1	B	294	THR	2.6
1	A	302	ARG	2.5
1	B	297	ILE	2.5
1	A	466	ASN	2.4
1	A	293	ASP	2.4
1	B	365	LYS	2.3
1	A	307	ALA	2.3
1	B	259	GLU	2.3
1	B	301	LYS	2.2
1	B	194	LEU	2.2
1	B	266	LEU	2.2
1	A	370	PHE	2.1
1	A	297	ILE	2.1
1	A	220	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	KKP	B	602	20/20	0.77	0.20	102,106,109,109	0
3	KKP	A	602	20/20	0.82	0.18	88,99,102,103	0
2	ZN	A	601	1/1	0.99	0.02	41,41,41,41	0
2	ZN	B	601	1/1	0.99	0.03	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KKP B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KKP A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.