



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:14 AM UTC

PDB ID : 3NG0 / pdb_00003ng0
Title : Crystal Structure of Glutamine Synthetase from Synechocystis sp. PCC 6803
Authors : Saelices, L.; Cascio, D.; Florencio, F.J.; Muro-Pastor, M.I.
Deposited on : 2010-06-10
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	473	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3732 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutamine synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	465	3674	2353	606	697	18	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	115	ALA	VAL	variant	UNP P77961
A	204	ALA	GLY	variant	UNP P77961
A	205	PHE	LEU	variant	UNP P77961
A	206	GLY	CYS	variant	UNP P77961

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	10	6	12	3	0	0

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	Mn 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	24	Total 24	O 24	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamine synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.02Å 132.02Å 202.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.92 – 2.80 32.92 – 2.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (32.92-2.80) 97.8 (32.92-2.80)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.81Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.8.0	Depositor
R, R_{free}	0.285 , 0.313 0.288 , 0.321	Depositor DCC
R_{free} test set	1298 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.498	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	3732	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/3778	1.35	27/5128 (0.5%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	SER	N-CA-C	9.71	121.46	111.07
1	A	388	ASN	N-CA-C	8.32	125.68	114.12
1	A	248	VAL	N-CA-CB	7.86	122.38	110.58
1	A	388	ASN	CA-C-N	7.33	134.89	121.70
1	A	388	ASN	C-N-CA	7.33	134.89	121.70
1	A	26	ASP	CA-CB-CG	6.49	119.09	112.60
1	A	461	ARG	N-CA-C	6.34	118.06	109.24
1	A	183	PHE	CA-CB-CG	6.22	120.02	113.80
1	A	472	ASP	CA-CB-CG	6.22	118.82	112.60
1	A	133	ALA	N-CA-C	6.09	118.70	107.99
1	A	169	GLY	CA-C-N	6.00	132.50	121.70
1	A	169	GLY	C-N-CA	6.00	132.50	121.70
1	A	470	TYR	N-CA-C	5.74	122.09	114.12
1	A	60	TRP	N-CA-C	5.56	114.09	108.75
1	A	375	PHE	CA-CB-CG	5.49	119.29	113.80
1	A	428	ASP	CA-CB-CG	5.48	118.08	112.60
1	A	191	ALA	N-CA-C	5.45	118.94	112.72
1	A	109	THR	CB-CA-C	5.41	119.89	110.79
1	A	432	LEU	CA-C-N	5.31	127.85	120.63
1	A	432	LEU	C-N-CA	5.31	127.85	120.63
1	A	215	GLU	CA-C-N	5.26	127.38	120.60
1	A	215	GLU	C-N-CA	5.26	127.38	120.60
1	A	42	ASP	CA-CB-CG	5.22	117.82	112.60
1	A	436	GLY	CA-C-N	5.09	127.07	120.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	436	GLY	C-N-CA	5.09	127.07	120.70
1	A	39	ASP	CA-CB-CG	5.07	117.67	112.60
1	A	236	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3674	0	3533	29	0
2	A	31	0	12	0	0
3	A	3	0	0	0	0
4	A	24	0	0	0	0
All	All	3732	0	3545	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ARG:HE	1:A:212:HIS:CD2	2.10	0.70
1:A:332:PRO:HB2	1:A:345:SER:HA	1.73	0.69
1:A:195:ARG:HE	1:A:212:HIS:HD2	1.42	0.66
1:A:312:LEU:HD12	1:A:415:PRO:HD2	1.79	0.65
1:A:331:ALA:HB1	1:A:332:PRO:HA	1.79	0.65
1:A:195:ARG:HH11	1:A:222:ASN:ND2	1.99	0.61
1:A:195:ARG:HD3	1:A:222:ASN:HD22	1.72	0.54
1:A:435:THR:HG22	1:A:437:VAL:HG23	1.88	0.54
1:A:332:PRO:HG3	1:A:363:ARG:HB2	1.91	0.52
1:A:329:PHE:HD2	1:A:330:GLU:HG3	1.76	0.50
1:A:422:LEU:HD22	1:A:447:ILE:HA	1.94	0.50
1:A:260:PRO:HD3	1:A:368:THR:CG2	2.42	0.49
1:A:52:PHE:CD1	1:A:70:MET:HG3	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ARG:HD3	1:A:222:ASN:ND2	2.29	0.47
1:A:118:LEU:HD11	1:A:124:ALA:HB3	1.97	0.46
1:A:22:LEU:HD22	1:A:72:PRO:HB3	1.97	0.46
1:A:194:ILE:HG12	1:A:198:MET:HE3	1.98	0.46
1:A:332:PRO:HG3	1:A:363:ARG:CB	2.46	0.45
1:A:351:SER:OG	1:A:358:LYS:HE3	2.18	0.44
1:A:337:TYR:HA	1:A:346:VAL:O	2.18	0.44
1:A:19:ILE:HG22	1:A:87:THR:HG23	2.00	0.44
1:A:197:GLU:O	1:A:201:THR:HG23	2.19	0.42
1:A:274:HIS:HB3	1:A:359:ARG:HG3	2.01	0.42
1:A:327:PRO:HA	1:A:333:VAL:HG21	2.02	0.42
1:A:160:ARG:HD3	1:A:177:GLY:HA2	2.02	0.42
1:A:174:TYR:CG	1:A:187:PRO:HG2	2.55	0.41
1:A:433:THR:HA	1:A:438:PHE:O	2.20	0.41
1:A:71:VAL:HG21	1:A:95:LYS:HD2	2.02	0.41
1:A:162:ASN:O	1:A:175:LYS:HE3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/473 (98%)	431 (94%)	26 (6%)	4 (1%)	14	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	ALA
1	A	390	ILE
1	A	231	LEU
1	A	332	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	392/399 (98%)	362 (92%)	30 (8%)	12	36

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LYS
1	A	19	ILE
1	A	20	ILE
1	A	27	THR
1	A	31	TRP
1	A	44	ASN
1	A	109	THR
1	A	147	GLU
1	A	150	SER
1	A	157	VAL
1	A	183	PHE
1	A	205	PHE
1	A	223	GLU
1	A	234	SER
1	A	244	VAL
1	A	248	VAL
1	A	267	ASN
1	A	273	VAL
1	A	280	ASP
1	A	293	PHE
1	A	298	LEU
1	A	311	LEU
1	A	330	GLU
1	A	360	LEU
1	A	408	LEU
1	A	410	LYS
1	A	426	GLU
1	A	448	GLU
1	A	460	LEU
1	A	468	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	ASN
1	A	192	GLN
1	A	212	HIS
1	A	222	ASN
1	A	316	ASN
1	A	334	ASN
1	A	354	ASN
1	A	453	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ANP	A	474	3	33,33,33	2.37	10 (30%)	45,52,52	2.21	13 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	474	3	-	9/18/38/38	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	474	ANP	PA-O3A	6.58	1.66	1.59
2	A	474	ANP	PG-O1G	5.03	1.53	1.46
2	A	474	ANP	PB-O1B	4.93	1.53	1.46
2	A	474	ANP	C6-N6	4.45	1.45	1.34
2	A	474	ANP	PG-O3G	-4.09	1.45	1.56
2	A	474	ANP	O2'-C2'	-3.10	1.35	1.43
2	A	474	ANP	PB-O3A	-2.88	1.55	1.59
2	A	474	ANP	C5'-C4'	-2.74	1.43	1.51
2	A	474	ANP	C2'-C3'	-2.20	1.47	1.53
2	A	474	ANP	C3'-C4'	-2.06	1.47	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	474	ANP	O4'-C1'-N9	5.47	118.60	108.09
2	A	474	ANP	N3-C2-N1	-5.28	120.58	128.58
2	A	474	ANP	C5-C4-N3	-4.99	119.84	126.72
2	A	474	ANP	N9-C8-N7	-4.14	108.06	113.94
2	A	474	ANP	N3-C4-N9	4.09	134.13	127.17
2	A	474	ANP	O5'-C5'-C4'	3.92	122.35	108.99
2	A	474	ANP	C4-N9-C8	3.83	109.75	105.74
2	A	474	ANP	C2-N3-C4	3.67	120.80	111.83
2	A	474	ANP	C5-N7-C8	3.36	108.72	103.45
2	A	474	ANP	C4-C5-N7	-2.76	107.42	110.58
2	A	474	ANP	C4-N9-C1'	-2.23	121.42	126.63
2	A	474	ANP	O2G-PG-O1G	-2.12	108.12	113.45
2	A	474	ANP	O3'-C3'-C4'	-2.00	105.33	111.08

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	474	ANP	PB-N3B-PG-O1G
2	A	474	ANP	PG-N3B-PB-O1B

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	474	ANP	PG-N3B-PB-O3A
2	A	474	ANP	C5'-O5'-PA-O1A
2	A	474	ANP	O4'-C4'-C5'-O5'
2	A	474	ANP	PB-O3A-PA-O1A
2	A	474	ANP	C2'-C1'-N9-C8
2	A	474	ANP	C5'-O5'-PA-O3A
2	A	474	ANP	O4'-C1'-N9-C8

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	465/473 (98%)	0.37	32 (6%) 23 16	8, 26, 53, 85	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	400	ILE	11.3
1	A	329	PHE	7.7
1	A	330	GLU	7.3
1	A	331	ALA	5.8
1	A	410	LYS	5.5
1	A	327	PRO	5.1
1	A	352	GLY	5.1
1	A	407	GLU	4.3
1	A	473	CYS	4.3
1	A	288	ASP	4.1
1	A	394	GLU	3.6
1	A	388	ASN	3.4
1	A	387	LYS	3.2
1	A	98	ARG	3.1
1	A	427	LYS	3.1
1	A	408	LEU	3.1
1	A	44	ASN	2.9
1	A	353	GLY	2.8
1	A	389	GLN	2.8
1	A	351	SER	2.8
1	A	399	ASP	2.7
1	A	326	VAL	2.7
1	A	328	GLY	2.6
1	A	101	GLU	2.4
1	A	444	GLU	2.4
1	A	356	LYS	2.4
1	A	341	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	332	PRO	2.1
1	A	409	ALA	2.1
1	A	344	ALA	2.1
1	A	339	GLN	2.1
1	A	398	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	A	476	1/1	0.84	0.21	119,119,119,119	0
3	MN	A	477	1/1	0.94	0.09	78,78,78,78	0
2	ANP	A	474	31/31	0.95	0.10	3,36,123,188	0
3	MN	A	475	1/1	0.95	0.06	60,60,60,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP A 474:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.