



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:16 AM UTC

PDB ID : 7NHV / pdb_00007nhv
Title : Crystal structure of the human METTL3-METTL14 complex with compound UOZ016
Authors : Bedi, R.K.; Huang, D.; Caflisch, A.
Deposited on : 2021-02-11
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	5% 73% 9% 18%
2	B	290	5% 75% 9% 16%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3756 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	0	0
			1563	1001	272	281	9			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	335	MET	-	initiating methionine	UNP Q86U44
A	336	GLY	-	expression tag	UNP Q86U44
A	337	HIS	-	expression tag	UNP Q86U44
A	338	HIS	-	expression tag	UNP Q86U44
A	339	HIS	-	expression tag	UNP Q86U44
A	340	HIS	-	expression tag	UNP Q86U44
A	341	HIS	-	expression tag	UNP Q86U44
A	342	HIS	-	expression tag	UNP Q86U44
A	343	SER	-	expression tag	UNP Q86U44
A	344	SER	-	expression tag	UNP Q86U44
A	345	GLY	-	expression tag	UNP Q86U44
A	346	ARG	-	expression tag	UNP Q86U44
A	347	GLU	-	expression tag	UNP Q86U44
A	348	ASN	-	expression tag	UNP Q86U44
A	349	LEU	-	expression tag	UNP Q86U44
A	350	TYR	-	expression tag	UNP Q86U44
A	351	PHE	-	expression tag	UNP Q86U44
A	352	GLN	-	expression tag	UNP Q86U44
A	353	GLY	-	expression tag	UNP Q86U44


- Molecule 2 is a protein called N6-adenosine-methyltransferase non-catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	1	0
			1880	1206	318	343	13			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	106	MET	-	initiating methionine	UNP Q9HCE5

- Molecule 3 is (R)-N-((1-(6-(benzylamino)pyrimidin-4-yl)-3-hydroxypiperidin-3-yl)methyl)-4-((4,4-dimethylpiperidin-1-yl)methyl)benzamide (CCD ID: UE5) (formula: C₃₂H₄₂N₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			40	32	6	2		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	114	Total	O	0	0
			114	114		
5	B	155	Total	O	0	0
			155	155		

- Molecule 1: N6-adenosine-methyltransferase catalytic subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	63.65Å 63.65Å 225.42Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.45 – 1.91 44.45 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.45-1.91) 99.9 (44.45-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 1.91Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.196 , 0.212 0.200 , 0.218	Depositor DCC
R_{free} test set	2122 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3756	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UE5, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1603	0.60	0/2185
2	B	0.38	0/1931	0.60	0/2629
All	All	0.38	0/3534	0.60	0/4814

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1563	0	1472	14	0
2	B	1880	0	1737	20	0
3	A	40	0	0	1	0
4	B	4	0	3	0	0
5	A	114	0	0	1	0
5	B	155	0	0	4	0
All	All	3756	0	3212	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:456:ILE:HD11	2:B:262:ILE:HD11	1.34	1.03
2:B:250:LYS:NZ	5:B:501:HOH:O	2.19	0.73
1:A:459:LYS:HE2	1:A:481:GLU:HG2	1.71	0.72
2:B:179:ILE:HG12	2:B:224:ILE:HD13	1.80	0.63
1:A:391:VAL:HB	1:A:530:LYS:HG2	1.81	0.63
2:B:168:MET:HG2	2:B:358:LEU:HD13	1.85	0.58
2:B:128:HIS:CE1	2:B:269:PRO:HG3	2.39	0.57
1:A:374:ILE:HG23	1:A:554:ILE:HD12	1.88	0.55
2:B:185:LYS:NZ	5:B:507:HOH:O	2.38	0.54
1:A:473:GLY:HA3	1:A:478:HIS:CE1	2.42	0.54
2:B:239:GLU:HG3	2:B:243:LEU:HD12	1.91	0.53
1:A:539:ASN:O	1:A:541:GLN:HG2	2.10	0.50
2:B:160:ILE:HD13	2:B:335:GLU:HB3	1.91	0.50
1:A:459:LYS:HB3	1:A:507:VAL:HG23	1.94	0.49
2:B:219:LEU:O	5:B:501:HOH:O	2.20	0.48
2:B:163:SER:OG	2:B:387:GLY:HA2	2.15	0.47
1:A:477:ASN:C	2:B:311:ILE:HD12	2.40	0.46
2:B:260[A]:CYS:SG	2:B:287:HIS:CE1	3.09	0.46
1:A:560:ASP:O	1:A:564:ARG:HG3	2.16	0.45
1:A:549:ASN:OD1	3:A:601:UE5:O37	2.34	0.45
2:B:168:MET:SD	2:B:374:TYR:HB2	2.57	0.44
2:B:254:ARG:HD3	5:B:556:HOH:O	2.17	0.44
2:B:230:PHE:CE1	2:B:339:LEU:HD22	2.53	0.43
2:B:254:ARG:H	2:B:254:ARG:HG2	1.67	0.43
2:B:168:MET:HE3	2:B:168:MET:HB2	1.93	0.42
1:A:456:ILE:HD11	2:B:262:ILE:CD1	2.26	0.42
2:B:178:ASP:O	2:B:181:GLU:HG3	2.20	0.41
2:B:217:MET:HE3	2:B:217:MET:HB2	1.91	0.41
1:A:533:LEU:CD2	1:A:547:LEU:HD12	2.51	0.41
1:A:480:LYS:HD2	1:A:482:HIS:NE2	2.37	0.40
1:A:400:ILE:HA	5:A:752:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/246 (79%)	192 (98%)	3 (2%)	0	100	100
2	B	237/290 (82%)	230 (97%)	6 (2%)	1 (0%)	30	19
All	All	432/536 (81%)	422 (98%)	9 (2%)	1 (0%)	43	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	271	LYS

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	B	401	-	3,3,3	2.11	1 (33%)	3,3,3	1.09	0
3	UE5	A	601	-	44,44,44	2.61	12 (27%)	54,62,62	1.83	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UE5	A	601	-	-	2/23/47/47	0/5/5/5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	UE5	C07-N06	-7.10	1.34	1.47
3	A	601	UE5	C27-N28	6.64	1.46	1.36
3	A	601	UE5	C05-N06	-6.54	1.29	1.46
3	A	601	UE5	C14-N15	6.37	1.47	1.33
3	A	601	UE5	C23-N22	4.33	1.47	1.37
3	A	601	UE5	C39-N06	-4.08	1.35	1.46
3	A	601	UE5	C21-N22	3.93	1.49	1.46
3	A	601	UE5	C07-C08	3.43	1.57	1.51
4	B	401	ACT	CH3-C	3.17	1.61	1.49
3	A	601	UE5	C04-C02	-2.36	1.50	1.53
3	A	601	UE5	C16-C17	2.33	1.56	1.53
3	A	601	UE5	C25-N24	2.23	1.37	1.33
3	A	601	UE5	C40-C39	2.01	1.58	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	UE5	N26-C25-N24	-5.74	119.90	128.58
3	A	601	UE5	C25-N24-C23	5.40	119.48	114.95
3	A	601	UE5	C16-N15-C14	-4.72	115.41	122.88
3	A	601	UE5	C20-N22-C21	3.61	121.37	113.86
3	A	601	UE5	C03-C02-C04	-3.60	104.86	110.04
3	A	601	UE5	C36-C23-N22	-3.09	118.62	122.35
3	A	601	UE5	C40-C02-C04	2.68	112.17	108.67
3	A	601	UE5	C04-C05-N06	2.35	113.55	111.16

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	UE5	C08-C07-N06-C05
3	A	601	UE5	N15-C16-C17-O37

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	UE5	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/246 (81%)	0.18	12 (5%) 27 32	26, 36, 59, 67	0
2	B	244/290 (84%)	0.28	15 (6%) 27 31	20, 35, 68, 82	1 (0%)
All	All	445/536 (83%)	0.24	27 (6%) 27 31	20, 36, 62, 82	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	153	ILE	5.3
1	A	368	LEU	4.8
2	B	296	VAL	4.4
2	B	392	ILE	3.8
2	B	272	THR	3.8
2	B	152	LEU	3.7
1	A	405	PRO	3.7
1	A	406	TYR	3.5
1	A	510	THR	3.5
1	A	369	PHE	3.4
2	B	390	GLU	3.1
1	A	401	HIS	2.9
2	B	274	THR	2.9
2	B	309	VAL	2.8
2	B	201	GLU	2.8
2	B	151	GLU	2.8
2	B	155	LEU	2.7
2	B	200	ARG	2.7
1	A	467	ILE	2.7
2	B	270	GLY	2.5
1	A	573	ILE	2.4
1	A	407	GLY	2.4
2	B	321	ILE	2.3
1	A	490	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	512	HIS	2.2
2	B	393	GLU	2.2
1	A	446	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UE5	A	601	40/40	0.81	0.14	34,46,55,56	0
4	ACT	B	401	4/4	0.97	0.07	31,32,34,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UE5 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.