



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:32 AM UTC

PDB ID : 1NIF / pdb_00001nif
Title : THE STRUCTURE OF CU-NITRITE REDUCTASE FROM ACHROMOBACTER CYCLOCLASTES AT FIVE PH VALUES, WITH NITRITE BOUND AND WITH TYPE II CU DEPLETED
Authors : Adman, E.T.; Godden, J.W.; Turley, S.
Deposited on : 1995-07-03
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	65% 29% ...

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2829 atoms, of which 28 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRITE REDUCTASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	333	Total	C	H	N	O	S	27	11	0
			2653	1676	28	448	490	11			

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	174	Total	O	0	0
			174	174		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRITE REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	98.40 Å 98.40 Å 98.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.70 10.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	86.0 (10.00-1.70) 90.3 (10.00-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.61 Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.175 , (Not available) 0.173 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 77.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.041 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2829	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.48	21/2739 (0.8%)	1.34	27/3731 (0.7%)

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	322	VAL	CA-CB	10.13	1.67	1.54
1	A	210	MET	SD-CE	9.89	2.04	1.79
1	A	73	MET	SD-CE	-8.81	1.57	1.79
1	A	236	ALA	CA-C	7.17	1.61	1.52
1	A	175	ILE	CA-CB	6.71	1.64	1.54
1	A	43	MET	SD-CE	-6.62	1.62	1.79
1	A	275	ASP	CA-CB	6.59	1.63	1.53
1	A	287	THR	CA-CB	5.98	1.64	1.53
1	A	244	VAL	CA-CB	5.81	1.61	1.54
1	A	264	VAL	CA-CB	5.74	1.62	1.54
1	A	197	GLU	CA-C	5.58	1.59	1.52
1	A	316	ALA	CA-CB	5.40	1.59	1.52
1	A	74	VAL	CA-CB	5.38	1.61	1.54
1	A	92	THR	CA-CB	5.31	1.61	1.54
1	A	81	VAL	CA-CB	5.26	1.60	1.54
1	A	28	HIS	CA-C	5.23	1.58	1.52
1	A	75	VAL	CA-C	5.18	1.57	1.52
1	A	264	VAL	C-O	-5.13	1.18	1.24
1	A	31	VAL	CA-CB	5.04	1.60	1.54
1	A	188	ASP	CA-C	5.02	1.59	1.53
1	A	185	VAL	CA-C	5.01	1.58	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	305	ASN	N-CA-C	-9.60	94.57	109.25
1	A	104	GLY	N-CA-C	9.35	120.83	111.95
1	A	307	ASN	N-CA-C	-8.77	95.08	108.67
1	A	268	GLY	N-CA-C	8.24	126.54	114.10
1	A	222	GLY	N-CA-C	7.58	125.44	115.36
1	A	312	PHE	N-CA-C	6.70	121.54	113.23
1	A	270	PHE	N-CA-C	6.22	120.34	112.87
1	A	203	TYR	N-CA-C	6.13	121.42	111.37
1	A	63	THR	N-CA-C	6.08	118.94	109.52
1	A	275	ASP	CA-CB-CG	-6.02	106.58	112.60
1	A	79	ASP	N-CA-C	-5.92	103.13	110.41
1	A	272	ASN	CA-C-N	-5.78	114.42	120.38
1	A	272	ASN	C-N-CA	-5.78	114.42	120.38
1	A	314	LEU	N-CA-C	5.62	120.20	113.23
1	A	169	PRO	N-CA-C	5.57	119.82	111.14
1	A	48	LYS	N-CA-C	5.54	117.68	108.99
1	A	218	ILE	N-CA-C	-5.53	99.54	107.78
1	A	127	THR	N-CA-C	5.38	119.70	113.18
1	A	54	ARG	N-CA-C	5.29	117.95	111.82
1	A	22	PRO	N-CA-C	-5.25	105.43	110.47
1	A	303	TYR	N-CA-C	-5.23	100.20	108.73
1	A	261	GLY	N-CA-C	-5.19	100.89	113.18
1	A	234	THR	N-CA-C	5.17	117.99	109.72
1	A	64	PHE	N-CA-C	-5.16	100.11	108.52
1	A	142	VAL	CA-C-N	-5.10	113.67	119.28
1	A	142	VAL	C-N-CA	-5.10	113.67	119.28
1	A	279	GLU	N-CA-C	-5.08	105.39	112.45

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2625	28	2561	60	0
2	A	2	0	0	0	0
3	A	174	0	0	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2801	28	2561	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:MET:CE	1:A:210:MET:SD	2.04	1.44
1:A:250[B]:ARG:NH1	3:A:1180:HOH:O	1.73	1.12
1:A:335:VAL:CG1	3:A:1336:HOH:O	2.35	0.74
1:A:22:PRO:HB2	1:A:221:ASN:HD21	1.53	0.74
1:A:229:GLY:O	1:A:321:LYS:NZ	2.23	0.72
1:A:335:VAL:HG12	3:A:1336:HOH:O	1.89	0.71
1:A:32:ALA:HA	3:A:1076:HOH:O	1.91	0.70
1:A:36:PRO:HG3	1:A:159[B]:ARG:NH2	2.06	0.70
1:A:36:PRO:HG3	1:A:159[B]:ARG:HH21	1.61	0.66
1:A:76:HIS:NE2	3:A:1076:HOH:O	2.30	0.64
1:A:26:HIS:HD2	1:A:27:ALA:O	1.81	0.63
1:A:26:HIS:HE1	1:A:74:VAL:H	1.45	0.63
1:A:237[B]:VAL:HG23	3:A:1237:HOH:O	2.00	0.61
1:A:60:HIS:HD2	3:A:3060:HOH:O	1.82	0.61
1:A:232:ALA:CB	1:A:321:LYS:HZ3	2.15	0.59
1:A:232:ALA:HB2	1:A:321:LYS:HZ3	1.67	0.59
1:A:307:ASN:HD22	1:A:307:ASN:C	2.13	0.57
1:A:60:HIS:HE1	1:A:196:TYR:O	1.88	0.57
1:A:232:ALA:CB	1:A:321:LYS:NZ	2.68	0.57
1:A:240:ARG:HB3	1:A:292:PHE:CZ	2.40	0.56
1:A:143:PRO:HA	1:A:146[A]:VAL:HG22	1.87	0.55
1:A:250[B]:ARG:NH2	1:A:310:GLU:CD	2.64	0.55
1:A:28:HIS:HE1	1:A:172:TYR:OH	1.90	0.54
1:A:237[B]:VAL:HG21	3:A:1325:HOH:O	2.07	0.54
1:A:319:HIS:HD2	3:A:1229:HOH:O	1.90	0.53
1:A:26:HIS:CE1	1:A:74:VAL:H	2.25	0.52
1:A:232:ALA:HB2	1:A:321:LYS:NZ	2.24	0.52
1:A:20[B]:VAL:HG22	3:A:1072:HOH:O	2.10	0.52
1:A:305:ASN:O	1:A:311:ALA:HB2	2.11	0.50
1:A:48:LYS:HE2	1:A:66:GLY:O	2.12	0.49
1:A:237[B]:VAL:HG22	1:A:324:GLY:HA3	1.94	0.49
1:A:257:ILE:HD12	1:A:302:ALA:HB3	1.95	0.48
1:A:17[A]:VAL:HG13	1:A:41:PHE:CD1	2.48	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:LYS:HD2	1:A:40:GLU:O	2.14	0.47
1:A:197:GLU:HB2	1:A:201:GLU:OE1	2.13	0.47
1:A:187:LYS:HD2	1:A:191:GLY:O	2.14	0.47
1:A:250[B]:ARG:NH2	1:A:310:GLU:OE1	2.46	0.47
1:A:11:THR:O	1:A:11:THR:CG2	2.62	0.47
1:A:232:ALA:HB3	1:A:321:LYS:NZ	2.29	0.47
1:A:250[B]:ARG:HH2	1:A:310:GLU:CD	2.20	0.47
1:A:43:MET:HG3	1:A:83:LEU:HD11	1.97	0.46
1:A:335:VAL:HG11	3:A:1336:HOH:O	2.06	0.45
1:A:11:THR:O	1:A:11:THR:HG22	2.15	0.45
1:A:22:PRO:HB2	1:A:221:ASN:ND2	2.27	0.45
1:A:26:HIS:O	1:A:28:HIS:HD2	2.00	0.45
1:A:207:VAL:O	1:A:211:ARG:HG3	2.18	0.43
1:A:83:LEU:O	1:A:121:THR:HA	2.18	0.43
1:A:237[B]:VAL:CG2	1:A:324:GLY:HA3	2.49	0.43
1:A:168:GLN:HA	1:A:169:PRO:HD3	1.87	0.42
1:A:141[A]:MET:SD	1:A:145:HIS:NE2	2.91	0.42
1:A:237[B]:VAL:CG2	3:A:1325:HOH:O	2.64	0.42
1:A:173:ASP:OD2	1:A:240:ARG:HD3	2.21	0.41
1:A:227:LEU:HA	1:A:227:LEU:HD23	1.78	0.41
1:A:17[A]:VAL:CG1	1:A:41:PHE:CE1	3.03	0.41
1:A:164:ASP:OD2	1:A:168:GLN:HB2	2.21	0.41
1:A:94:LEU:HG	1:A:95:HIS:N	2.36	0.40
1:A:49:LYS:HD2	1:A:58:GLU:CD	2.45	0.40
1:A:162:LEU:O	1:A:163:LYS:HD3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/340 (101%)	337 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	282/273 (103%)	259 (92%)	23 (8%)	10 2

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	SER
1	A	17[A]	VAL
1	A	17[B]	VAL
1	A	33	LYS
1	A	50	LEU
1	A	94	LEU
1	A	98	ASP
1	A	106	LEU
1	A	111	LEU
1	A	131	VAL
1	A	166	LYS
1	A	194	LYS
1	A	213	LEU
1	A	221	ASN
1	A	230	ASP
1	A	240	ARG
1	A	247	GLN
1	A	282	LEU
1	A	305	ASN
1	A	307	ASN
1	A	330	LEU
1	A	339	SER
1	A	340	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	HIS
1	A	28	HIS
1	A	60	HIS
1	A	76	HIS
1	A	221	ASN
1	A	307	ASN
1	A	319	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/340 (97%)	-0.44	2 (0%) 85 88	9, 17, 31, 59	22 (6%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	165	GLU	2.7
1	A	340	MET	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU	A	502	1/1	0.97	0.06	4,4,4,4	1
2	CU	A	501	1/1	0.99	0.02	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.