



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:03 AM UTC

PDB ID : 5NIW / pdb_00005niw
Title : Glucose oxydase mutant A2
Authors : Hoffmann, K.; Frank, D.
Deposited on : 2017-03-27
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.



■ Percentile relative to all X-ray structures

Percentile relative to X-ray structures of similar resolution

Metric

Whole archive
(#Entries)

Similar resolution
(#Entries, resolution range(Å))

 R_{free}

Clashscore

Ramachandran outliers

Sidechain outliers

RSRZ outliers

180053

190562

187476

187428

180081

7662 (1.80-1.80)

8479 (1.80-1.80)

8391 (1.80-1.80)

8390 (1.80-1.80)

7663 (1.80-1.80)

fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled.

The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol

Chain

Length

Quality of chain

1

A

581

0.

91°

91%

100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	607	X	-	-	-
6	DIO	A	611	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glucose oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	578	Total	C	N	O	S	2	4	0
			4427	2792	756	867	12			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	30	VAL	THR	conflict	UNP P13006
A	37	LYS	ARG	conflict	UNP P13006
A	94	VAL	ILE	conflict	UNP P13006
A	106	ILE	VAL	conflict	UNP P13006
A	162	THR	ALA	conflict	UNP P13006
A	556	VAL	MET	conflict	UNP P13006

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is OXYGEN MOLECULE (CCD ID: OXY) (formula: O₂).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			2	2		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 1,4-DIETHYLENE DIOXIDE (CCD ID: DIO) (formula: $C_4H_8O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0
6	A	1	Total C O 6 4 2	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is O-ACETALDEHYDYL-HEXAETHYLENE GLYCOL (CCD ID: P4C) (formula: C₁₄H₂₈O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			22	14	8		
8	A	1	Total	C	O	0	0
			22	14	8		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	364	Total	O	0	0
			364	364		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucose oxidase



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.10Å 128.10Å 77.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	110.94 – 1.80 110.94 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (110.94-1.80) 99.8 (110.94-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.154 , 0.191 (Not available) , 0.175	Depositor DCC
R_{free} test set	3389 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	16.3	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5085	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, PEG, OXY, DIO, P4C, NAG, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.41	14/4543 (0.3%)	1.15	8/6202 (0.1%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	207	GLY	C-O	6.46	1.31	1.24
1	A	310	GLU	N-CA	6.29	1.52	1.46
1	A	373	LEU	C-O	6.15	1.35	1.23
1	A	451	ASP	N-CA	-6.08	1.39	1.46
1	A	346	GLY	C-O	5.84	1.32	1.23
1	A	263	ARG	CZ-NH2	5.65	1.40	1.33
1	A	321	PRO	CA-C	5.35	1.58	1.53
1	A	282	LYS	C-O	5.26	1.30	1.24
1	A	464	GLN	C-O	5.19	1.30	1.24
1	A	379	GLU	C-O	5.18	1.30	1.24
1	A	164	CYS	N-CA	-5.09	1.40	1.46
1	A	527	GLU	CA-CB	5.08	1.61	1.53
1	A	448	PHE	N-CA	5.05	1.52	1.45
1	A	239	ARG	CA-C	5.04	1.59	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	507	ILE	N-CA-CB	-7.92	105.51	110.50
1	A	263	ARG	NE-CZ-NH1	-6.55	114.95	121.50
1	A	263	ARG	NE-CZ-NH2	6.15	124.73	119.20
1	A	458	GLU	N-CA-C	5.71	117.51	111.28
1	A	547	ILE	N-CA-C	5.55	117.53	112.43
1	A	551	ILE	CA-C-N	-5.31	114.91	120.38
1	A	551	ILE	C-N-CA	-5.31	114.91	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	51	SER	N-CA-C	5.16	116.90	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4427	0	4200	32	0
2	B	39	0	34	0	0
3	A	53	0	31	3	0
4	A	2	0	0	0	0
5	A	70	0	65	1	1
6	A	72	0	96	8	0
7	A	14	0	20	0	1
8	A	44	0	56	8	0
9	A	364	0	0	5	1
All	All	5085	0	4502	33	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ASN:O	9:A:701:HOH:O	1.86	0.91
1:A:57[A]:ASP:OD2	9:A:702:HOH:O	1.89	0.90
1:A:275:ASN:HB3	9:A:701:HOH:O	1.77	0.83
1:A:277:HIS:ND1	9:A:704:HOH:O	2.15	0.78
1:A:107:ASN:HB2	3:A:601:FAD:C5X	2.18	0.73
1:A:273:LYS:HZ2	8:A:625:P4C:H91	1.62	0.65
1:A:542:GLN:HE22	5:A:607:NAG:C7	2.12	0.63
1:A:273:LYS:HZ3	8:A:625:P4C:H151	1.68	0.58
1:A:196:ARG:NH1	9:A:703:HOH:O	2.02	0.57
1:A:447:HIS:NE2	8:A:626:P4C:C20	2.69	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:447:HIS:NE2	8:A:626:P4C:H202	2.20	0.55
1:A:107:ASN:HB2	3:A:601:FAD:N5	2.22	0.54
1:A:273:LYS:HZ2	8:A:625:P4C:C9	2.20	0.53
1:A:204:PHE:CZ	1:A:213:SER:HB2	2.47	0.49
1:A:30:VAL:HG11	1:A:547:ILE:O	2.13	0.49
1:A:164:CYS:HA	6:A:611:DIO:H11	1.95	0.48
1:A:563:VAL:HG22	6:A:619:DIO:H22	1.95	0.48
8:A:625:P4C:H91	8:A:625:P4C:H52	1.96	0.48
1:A:164:CYS:HA	6:A:611:DIO:C1	2.44	0.48
1:A:141:LEU:HD21	6:A:611:DIO:H12	1.97	0.46
1:A:243:GLN:HG3	6:A:622:DIO:O1'	2.16	0.46
1:A:109:GLY:O	1:A:217:ASN:HA	2.17	0.45
1:A:273:LYS:HE2	8:A:625:P4C:O19	2.18	0.44
1:A:543:GLY:HA2	6:A:620:DIO:H1'1	1.99	0.43
1:A:126:PHE:O	1:A:525:PRO:HD3	2.19	0.42
1:A:107:ASN:HB2	3:A:601:FAD:C4X	2.49	0.42
1:A:507:ILE:N	1:A:508:PRO:CD	2.83	0.42
1:A:15:VAL:HA	1:A:18:ARG:HD2	2.02	0.42
1:A:37[A]:LYS:HD2	1:A:37[A]:LYS:HA	1.85	0.42
1:A:397:GLU:HB3	6:A:612:DIO:H2'1	2.03	0.41
1:A:273:LYS:HE2	8:A:625:P4C:C21	2.50	0.41
1:A:133:TRP:HE1	6:A:611:DIO:H2'1	1.86	0.41
1:A:524:MET:O	1:A:530:GLY:HA3	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:607:NAG:O3	5:A:607:NAG:O3[6_767]	1.86	0.34
7:A:624:PEG:O2	9:A:974:HOH:O[2_774]	1.93	0.27

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	578/581 (100%)	559 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	461/467 (99%)	460 (100%)	1 (0%)	87	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	219	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	ASN
1	A	243	GLN
1	A	278	ASN
1	A	375	GLN
1	A	407	ASN
1	A	493	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	1.00	0	17,19,21	1.75	5 (29%)
2	NAG	B	2	2	14,14,15	1.06	1 (7%)	17,19,21	1.64	3 (17%)
2	BMA	B	3	2	11,11,12	1.51	2 (18%)	15,15,17	2.06	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	BMA	C2-C3	3.53	1.57	1.52
2	B	3	BMA	O5-C5	2.49	1.48	1.43
2	B	2	NAG	O5-C1	-2.12	1.40	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	BMA	O5-C5-C6	4.13	115.71	107.66
2	B	2	NAG	O7-C7-C8	-4.13	114.70	122.05
2	B	3	BMA	O2-C2-C3	3.65	117.70	110.15
2	B	2	NAG	O7-C7-N2	3.41	128.01	121.98
2	B	3	BMA	O3-C3-C2	3.38	116.94	110.05
2	B	1	NAG	C1-C2-N2	-3.11	105.53	110.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C4-C3-C2	-2.97	106.67	111.02
2	B	1	NAG	O5-C1-C2	-2.95	106.72	111.29
2	B	1	NAG	O3-C3-C2	2.88	115.38	109.40
2	B	1	NAG	O5-C5-C6	-2.48	102.84	107.66
2	B	3	BMA	C1-O5-C5	2.44	115.46	112.19
2	B	2	NAG	O4-C4-C5	-2.31	103.63	109.32

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C1-C2-N2-C7
2	B	2	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	DIO	A	618	-	6,6,6	0.81	0	6,6,6	1.12	0
5	NAG	A	606	1	14,14,15	1.22	1 (7%)	17,19,21	2.21	5 (29%)
5	NAG	A	608	1	14,14,15	0.95	0	17,19,21	2.07	9 (52%)
6	DIO	A	619	-	6,6,6	0.79	0	6,6,6	1.46	1 (16%)
6	DIO	A	612	-	6,6,6	0.47	0	6,6,6	2.23	2 (33%)
6	DIO	A	611	-	6,6,6	0.54	0	6,6,6	1.22	0
6	DIO	A	615	-	6,6,6	0.51	0	6,6,6	1.20	0
6	DIO	A	614	-	6,6,6	0.78	0	6,6,6	0.93	0
5	NAG	A	609	1	14,14,15	1.43	2 (14%)	17,19,21	1.94	3 (17%)
4	OXY	A	602	-	1,1,1	0.15	0	-	-	-
8	P4C	A	625	-	21,21,21	0.65	0	20,20,20	1.10	1 (5%)
8	P4C	A	626	-	21,21,21	0.89	0	20,20,20	2.03	7 (35%)
6	DIO	A	616	-	6,6,6	0.49	0	6,6,6	0.83	0
6	DIO	A	620	-	6,6,6	0.58	0	6,6,6	1.11	1 (16%)
3	FAD	A	601	-	58,58,58	1.78	13 (22%)	85,89,89	1.89	20 (23%)
6	DIO	A	613	-	6,6,6	0.71	0	6,6,6	0.59	0
5	NAG	A	610	1	14,14,15	0.67	0	17,19,21	1.37	1 (5%)
7	PEG	A	624	-	6,6,6	0.41	0	5,5,5	0.87	0
6	DIO	A	617	-	6,6,6	0.97	0	6,6,6	1.38	0
6	DIO	A	621	-	6,6,6	0.48	0	6,6,6	0.66	0
6	DIO	A	622	-	6,6,6	0.52	0	6,6,6	0.39	0
5	NAG	A	607	1	14,14,15	0.81	1 (7%)	17,19,21	2.41	8 (47%)
7	PEG	A	623	-	6,6,6	0.42	0	5,5,5	1.25	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	DIO	A	618	-	-	-	0/1/1/1
5	NAG	A	606	1	-	1/6/23/26	0/1/1/1
5	NAG	A	608	1	-	0/6/23/26	0/1/1/1
6	DIO	A	619	-	-	-	0/1/1/1
6	DIO	A	612	-	-	-	0/1/1/1
6	DIO	A	611	-	-	-	0/1/1/1
6	DIO	A	615	-	-	-	0/1/1/1
6	DIO	A	614	-	-	-	0/1/1/1
5	NAG	A	609	1	-	0/6/23/26	0/1/1/1
8	P4C	A	626	-	-	11/18/19/19	-
8	P4C	A	625	-	-	13/18/19/19	-
6	DIO	A	616	-	-	-	0/1/1/1
6	DIO	A	620	-	-	-	0/1/1/1
3	FAD	A	601	-	-	5/34/50/50	0/6/6/6
6	DIO	A	613	-	-	-	0/1/1/1
5	NAG	A	610	1	-	2/6/23/26	0/1/1/1
7	PEG	A	624	-	-	3/4/4/4	-
6	DIO	A	617	-	-	-	0/1/1/1
6	DIO	A	621	-	-	-	0/1/1/1
6	DIO	A	622	-	-	-	0/1/1/1
5	NAG	A	607	1	1/1/5/7	3/6/23/26	0/1/1/1
7	PEG	A	623	-	-	3/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	FAD	C9A-C5X	5.77	1.50	1.41
5	A	609	NAG	O5-C1	-4.16	1.36	1.43
3	A	601	FAD	C5A-C4A	3.96	1.46	1.39
3	A	601	FAD	C8-C7	3.61	1.49	1.40
3	A	601	FAD	C1'-C2'	-3.48	1.47	1.52
3	A	601	FAD	C2-N1	3.43	1.44	1.36
3	A	601	FAD	P-O3P	2.90	1.62	1.59
3	A	601	FAD	C4X-N5	2.76	1.36	1.30
3	A	601	FAD	O4B-C1B	2.68	1.48	1.42
3	A	601	FAD	C2A-N3A	2.61	1.38	1.33
3	A	601	FAD	C1'-N10	-2.58	1.41	1.47
5	A	609	NAG	C2-N2	-2.54	1.42	1.46
3	A	601	FAD	O2'-C2'	-2.52	1.38	1.43
5	A	606	NAG	O5-C5	-2.33	1.38	1.43
5	A	607	NAG	C1-C2	2.27	1.55	1.52
3	A	601	FAD	C4-N3	-2.18	1.34	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	FAD	C6-C5X	2.01	1.43	1.40

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	FAD	N3A-C4A-N9A	7.00	139.06	127.17
5	A	606	NAG	C4-C3-C2	-6.64	101.29	111.02
3	A	601	FAD	C5A-C4A-N3A	-6.57	117.67	126.72
5	A	607	NAG	C1-C2-N2	5.57	119.21	110.43
3	A	601	FAD	C4A-N9A-C8A	5.20	111.20	105.74
5	A	609	NAG	C4-C3-C2	-4.44	104.51	111.02
5	A	609	NAG	C1-C2-N2	-4.40	103.50	110.43
8	A	626	P4C	O16-C15-C14	4.38	130.30	110.35
8	A	626	P4C	O7-C6-C5	4.13	129.20	110.35
5	A	607	NAG	O5-C5-C4	-3.87	101.40	110.83
6	A	612	DIO	O1-C2-C2'	-3.78	92.99	110.93
5	A	607	NAG	C2-N2-C7	3.74	127.91	122.90
5	A	608	NAG	C1-O5-C5	3.69	117.13	112.19
3	A	601	FAD	C4'-C3'-C2'	-3.55	107.67	113.57
3	A	601	FAD	O2'-C2'-C3'	3.45	117.31	109.25
3	A	601	FAD	O4-C4-C4X	-3.33	117.74	126.53
3	A	601	FAD	C4A-N9A-C1B	-3.32	118.86	126.63
6	A	612	DIO	C2'-O1'-C1'	3.32	120.60	109.88
5	A	609	NAG	C2-N2-C7	-3.31	118.46	122.90
5	A	610	NAG	O3-C3-C4	3.30	118.16	110.38
5	A	608	NAG	C1-C2-N2	-3.30	105.24	110.43
3	A	601	FAD	C2A-N3A-C4A	3.20	119.64	111.83
8	A	626	P4C	O22-C21-C20	-3.06	117.04	126.10
5	A	608	NAG	C6-C5-C4	-2.84	106.06	113.02
5	A	608	NAG	C3-C4-C5	-2.83	105.11	110.23
3	A	601	FAD	N3A-C2A-N1A	-2.82	124.31	128.58
6	A	619	DIO	C2-O1-C1	2.81	118.97	109.88
5	A	608	NAG	O6-C6-C5	-2.72	102.06	111.33
5	A	606	NAG	O6-C6-C5	-2.68	102.19	111.33
3	A	601	FAD	C2'-C1'-N10	2.64	122.67	110.20
8	A	626	P4C	O10-C9-C8	2.60	122.20	110.35
5	A	606	NAG	O5-C5-C4	-2.60	104.51	110.83
3	A	601	FAD	C6-C7-C8	-2.58	115.91	119.69
3	A	601	FAD	O2P-P-O1P	2.53	124.20	112.44
5	A	606	NAG	C2-N2-C7	-2.47	119.58	122.90
3	A	601	FAD	O4-C4-N3	2.46	124.75	120.11
5	A	606	NAG	O5-C1-C2	-2.41	107.56	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	623	PEG	O2-C3-C4	2.40	120.69	110.11
5	A	607	NAG	C4-C3-C2	2.40	114.53	111.02
5	A	607	NAG	C1-O5-C5	2.38	115.38	112.19
3	A	601	FAD	C4-C4X-N5	2.37	121.49	118.21
5	A	607	NAG	O7-C7-N2	2.35	126.13	121.98
3	A	601	FAD	C5'-C4'-C3'	-2.34	107.80	112.22
3	A	601	FAD	C5A-C4A-N9A	-2.34	103.26	105.81
5	A	608	NAG	C4-C3-C2	2.33	114.43	111.02
8	A	626	P4C	O7-C8-C9	2.28	120.74	110.35
5	A	608	NAG	O5-C5-C4	2.26	116.32	110.83
6	A	620	DIO	C2-O1-C1	2.26	117.18	109.88
5	A	608	NAG	O3-C3-C4	2.25	115.69	110.38
5	A	607	NAG	C8-C7-N2	-2.17	112.51	116.12
8	A	626	P4C	O4-C3-C2	2.17	119.68	110.11
5	A	607	NAG	C3-C4-C5	-2.17	106.30	110.23
5	A	608	NAG	O4-C4-C5	-2.16	103.99	109.32
8	A	626	P4C	C11-O10-C9	2.16	122.71	113.26
3	A	601	FAD	N9A-C8A-N7A	-2.13	110.92	113.94
3	A	601	FAD	C9A-C5X-N5	-2.12	120.20	122.45
8	A	625	P4C	O10-C9-C8	2.09	119.86	110.35
3	A	601	FAD	C5A-C6A-N6A	-2.09	118.12	123.29
3	A	601	FAD	C4A-C5A-N7A	-2.05	108.24	110.58

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	607	NAG	C1

All (41) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	FAD	N10-C1'-C2'-O2'
3	A	601	FAD	PA-O3P-P-O5'
5	A	607	NAG	C1-C2-N2-C7
8	A	626	P4C	C14-C15-O16-C17
5	A	610	NAG	O5-C5-C6-O6
8	A	625	P4C	C15-C14-O13-C12
5	A	607	NAG	C4-C5-C6-O6
8	A	625	P4C	C12-C11-O10-C9
5	A	607	NAG	O5-C5-C6-O6
5	A	610	NAG	C4-C5-C6-O6
8	A	626	P4C	O16-C17-C18-O19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	A	625	P4C	O4-C5-C6-O7
8	A	625	P4C	C18-C17-O16-C15
8	A	626	P4C	C9-C8-O7-C6
8	A	625	P4C	O10-C11-C12-O13
7	A	624	PEG	O1-C1-C2-O2
8	A	626	P4C	O7-C8-C9-O10
8	A	625	P4C	O13-C14-C15-O16
7	A	623	PEG	O1-C1-C2-O2
8	A	626	P4C	O4-C5-C6-O7
8	A	626	P4C	C6-C5-O4-C3
8	A	625	P4C	O1-C2-C3-O4
8	A	626	P4C	O1-C2-C3-O4
3	A	601	FAD	P-O3P-PA-O1A
8	A	625	P4C	C5-C6-O7-C8
8	A	625	P4C	O7-C8-C9-O10
8	A	625	P4C	C14-C15-O16-C17
7	A	623	PEG	C4-C3-O2-C2
7	A	624	PEG	C1-C2-O2-C3
8	A	625	P4C	C2-C3-O4-C5
8	A	626	P4C	C11-C12-O13-C14
8	A	625	P4C	O16-C17-C18-O19
7	A	623	PEG	O2-C3-C4-O4
8	A	625	P4C	C9-C8-O7-C6
3	A	601	FAD	P-O3P-PA-O2A
8	A	626	P4C	C21-C20-O19-C18
7	A	624	PEG	O2-C3-C4-O4
5	A	606	NAG	C1-C2-N2-C7
3	A	601	FAD	C2B-C1B-N9A-C8A
8	A	626	P4C	C18-C17-O16-C15
8	A	626	P4C	O10-C11-C12-O13

There are no ring outliers.

10 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	619	DIO	1	0
6	A	612	DIO	1	0
6	A	611	DIO	4	0
8	A	625	P4C	6	0
8	A	626	P4C	2	0
6	A	620	DIO	1	0
3	A	601	FAD	3	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	624	PEG	0	1
6	A	622	DIO	1	0
5	A	607	NAG	1	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	578/581 (99%)	-0.39	5 (0%) 81 81	9, 22, 35, 66	5 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	581	SER	4.5
1	A	260	THR	4.1
1	A	257	GLN	2.6
1	A	256	SER	2.1
1	A	258	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	B	3	11/12	0.70	0.19	47,51,65,65	0
2	NAG	B	2	14/15	0.92	0.10	29,30,35,39	0
2	NAG	B	1	14/15	0.97	0.06	25,27,31,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	607	14/15	0.54	0.22	69,79,86,87	0
5	NAG	A	610	14/15	0.68	0.19	51,58,68,76	0
6	DIO	A	618	6/6	0.76	0.23	49,51,56,56	0
6	DIO	A	617	6/6	0.77	0.19	42,46,48,48	0
6	DIO	A	620	6/6	0.82	0.20	62,64,65,66	0
7	PEG	A	624	7/7	0.85	0.18	43,55,59,61	0
8	P4C	A	626	22/22	0.85	0.18	32,51,58,60	0
8	P4C	A	625	22/22	0.86	0.18	36,50,77,84	0
6	DIO	A	612	6/6	0.87	0.14	41,43,46,49	0
6	DIO	A	621	6/6	0.87	0.20	64,68,70,71	0
6	DIO	A	622	6/6	0.87	0.19	59,60,61,64	0
6	DIO	A	614	6/6	0.88	0.16	44,47,52,54	0
6	DIO	A	615	6/6	0.88	0.16	49,52,53,53	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	A	608	14/15	0.89	0.12	37,44,48,50	0
5	NAG	A	609	14/15	0.90	0.11	32,37,48,52	0
7	PEG	A	623	7/7	0.92	0.21	31,31,32,33	7
6	DIO	A	619	6/6	0.92	0.12	38,40,41,42	0
5	NAG	A	606	14/15	0.94	0.08	19,22,31,32	0
6	DIO	A	611	6/6	0.94	0.11	37,40,45,45	0
4	OXY	A	602	2/2	0.94	0.10	28,28,28,38	0
6	DIO	A	613	6/6	0.96	0.09	29,32,37,40	0
6	DIO	A	616	6/6	0.96	0.09	35,36,37,37	6
3	FAD	A	601	53/53	0.98	0.05	16,18,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P4C A 626:


2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around P4C A 625:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.