



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:31 AM UTC

PDB ID : 5NJF / pdb_00005njf
Title : E. coli Microcin-processing metalloprotease TldD/E (TldD H262A mutant) with pentapeptide bound
Authors : Ghilarov, D.; Serebryakova, M.; Stevenson, C.E.M.; Hearnshaw, S.J.; Volkov, D.; Maxwell, A.; Lawson, D.M.; Severinov, K.
Deposited on : 2017-03-28
Resolution : 1.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	 91% 5%
1	C	495	 94%
2	B	450	 96%
2	D	450	 96%
3	E	5	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	5	 100%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 17169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metalloprotease TldD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	38	0
			3867	2403	674	771	19			
1	C	480	Total	C	N	O	S	0	31	0
			3837	2392	667	760	18			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP P0AGG8
A	-12	GLY	-	expression tag	UNP P0AGG8
A	-11	SER	-	expression tag	UNP P0AGG8
A	-10	SER	-	expression tag	UNP P0AGG8
A	-9	HIS	-	expression tag	UNP P0AGG8
A	-8	HIS	-	expression tag	UNP P0AGG8
A	-7	HIS	-	expression tag	UNP P0AGG8
A	-6	HIS	-	expression tag	UNP P0AGG8
A	-5	HIS	-	expression tag	UNP P0AGG8
A	-4	HIS	-	expression tag	UNP P0AGG8
A	-3	SER	-	expression tag	UNP P0AGG8
A	-2	GLN	-	expression tag	UNP P0AGG8
A	-1	ASP	-	expression tag	UNP P0AGG8
A	0	PRO	-	expression tag	UNP P0AGG8
A	262	ALA	HIS	engineered mutation	UNP P0AGG8
A	401	ASP	GLY	engineered mutation	UNP P0AGG8
C	-13	MET	-	initiating methionine	UNP P0AGG8
C	-12	GLY	-	expression tag	UNP P0AGG8
C	-11	SER	-	expression tag	UNP P0AGG8
C	-10	SER	-	expression tag	UNP P0AGG8
C	-9	HIS	-	expression tag	UNP P0AGG8
C	-8	HIS	-	expression tag	UNP P0AGG8
C	-7	HIS	-	expression tag	UNP P0AGG8
C	-6	HIS	-	expression tag	UNP P0AGG8
C	-5	HIS	-	expression tag	UNP P0AGG8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	HIS	-	expression tag	UNP P0AGG8
C	-3	SER	-	expression tag	UNP P0AGG8
C	-2	GLN	-	expression tag	UNP P0AGG8
C	-1	ASP	-	expression tag	UNP P0AGG8
C	0	PRO	-	expression tag	UNP P0AGG8
C	262	ALA	HIS	engineered mutation	UNP P0AGG8
C	401	ASP	GLY	engineered mutation	UNP P0AGG8

- Molecule 2 is a protein called Metalloprotease PmbA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	444	Total	C	N	O	S	0	32	0
			3599	2236	634	713	16			
2	D	445	Total	C	N	O	S	0	34	0
			3627	2251	636	724	16			

- Molecule 3 is a protein called ALA-ALA-ALA-ALA-ALA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	5	Total	C	N	O	0	0	0
			26	15	5	6			
3	F	5	Total	C	N	O	0	0	0
			26	15	5	6			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	C	1	Total	Zn	0	0
			1	1		

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 8 4 4	0	1
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 8 4 4	0	1
6	D	1	Total C O 4 2 2	0	0

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Na 1 1	0	0
7	B	1	Total Na 1 1	0	0
7	C	1	Total Na 1 1	0	0
7	D	1	Total Na 1 1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	507	Total O 532 532	0	25
8	B	497	Total O 525 525	0	28
8	C	436	Total O 461 461	0	25
8	D	532	Total O 554 554	0	22

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	F	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Metalloprotease TldD




- Molecule 1: Metalloprotease TldD



- Molecule 2: Metalloprotease PmbA



- Molecule 2: Metalloprotease PmbA



- Molecule 3: ALA-ALA-ALA-ALA-ALA



There are no outlier residues recorded for this chain.

- Molecule 3: ALA-ALA-ALA-ALA-ALA

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.16Å 174.48Å 83.65Å 90.00° 90.04° 90.00°	Depositor
Resolution (Å)	65.16 – 1.42 65.16 – 1.42	Depositor EDS
% Data completeness (in resolution range)	98.8 (65.16-1.42) 98.8 (65.16-1.42)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.42Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.139 , 0.171 0.147 , 0.178	Depositor DCC
R_{free} test set	17317 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	15.7	Xtriage
Anisotropy	0.443	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.389 for h,-k,-l	Xtriage
Reported twinning fraction	0.631 for H, K, L 0.369 for -h,-k,l	Depositor
Outliers	0 of 346357 reflections	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	17169	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, NA, ZN, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/3921	0.80	0/5293
1	C	0.71	0/3891	0.79	0/5262
2	B	0.72	0/3659	0.78	0/4947
2	D	0.74	0/3688	0.77	1/4988 (0.0%)
3	E	0.82	0/25	1.06	0/32
3	F	0.66	0/25	0.94	0/32
All	All	0.72	0/15209	0.79	1/20554 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	244	ALA	N-CA-C	5.19	112.89	108.22

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3867	0	3810	26	0
1	C	3837	0	3780	17	0
2	B	3599	0	3524	10	0
2	D	3627	0	3534	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	26	0	24	0	0
3	F	26	0	24	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	24	0	25	0	0
5	B	12	0	13	0	0
5	C	12	0	13	0	0
5	D	12	0	13	0	0
6	A	8	0	12	0	0
6	B	16	0	24	0	0
6	C	4	0	6	0	0
6	D	20	0	30	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
8	A	532	0	0	0	0
8	B	525	0	0	1	0
8	C	461	0	0	4	0
8	D	554	0	0	1	0
8	F	1	0	0	0	0
All	All	17169	0	14832	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:202[B]:SER:OG	8:C:958[B]:HOH:O	1.58	1.21
1:C:340[B]:ARG:HH11	1:C:340[B]:ARG:HG2	1.09	1.14
1:A:101[A]:ARG:HH11	1:A:101[A]:ARG:HG2	1.06	1.07
2:B:115[A]:GLU:OE2	8:B:804[A]:HOH:O	1.86	0.92
1:A:44[A]:TYR:CD2	1:A:62[A]:ILE:CD1	2.56	0.88
1:A:101[A]:ARG:HG2	1:A:101[A]:ARG:NH1	1.83	0.88
1:A:101[A]:ARG:HH11	1:A:101[A]:ARG:CG	1.88	0.87
1:C:340[B]:ARG:HG2	1:C:340[B]:ARG:NH1	1.86	0.84
1:A:44[A]:TYR:CD2	1:A:62[A]:ILE:HD12	2.14	0.82
1:A:44[A]:TYR:HD2	1:A:62[A]:ILE:CD1	1.97	0.77
1:A:44[A]:TYR:HD2	1:A:62[A]:ILE:HD11	1.55	0.72
2:B:314[GLU:OE1	2:B:316[B]:ARG:NH2	2.29	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:340[B]:ARG:NH1	1:C:340[B]:ARG:CG	2.60	0.61
1:A:101[A]:ARG:NH1	1:A:101[A]:ARG:CG	2.54	0.60
1:C:101[A]:ARG:HG2	8:C:617[A]:HOH:O	2.03	0.58
1:A:250:VAL:HG13	1:A:386[B]:ILE:HG12	1.87	0.57
1:C:66[A]:VAL:HG21	1:C:89:LEU:HD21	1.89	0.55
1:A:66[B]:VAL:HG21	1:A:89:LEU:HD21	1.91	0.53
2:B:11[A]:VAL:HG22	2:B:167:TYR:CE2	2.45	0.51
1:C:357:HIS:HD2	8:C:978:HOH:O	1.93	0.51
2:B:8[B]:ILE:O	2:B:11[B]:VAL:HG12	2.11	0.51
2:B:7:VAL:O	2:B:11[A]:VAL:HG23	2.11	0.50
1:C:228:LYS:NZ	8:C:603:HOH:O	2.45	0.50
1:C:101[A]:ARG:HG2	1:C:101[A]:ARG:NH1	2.26	0.49
2:D:115[B]:GLU:H	2:D:115[B]:GLU:CD	2.19	0.49
1:C:261:LEU:O	1:C:265[B]:VAL:HG22	2.13	0.48
1:A:404:VAL:CG2	2:B:377[B]:MET:HE3	2.44	0.47
1:A:470[A]:LEU:C	1:A:470[A]:LEU:HD12	2.39	0.47
1:C:337[B]:LEU:HD22	1:C:337[B]:LEU:O	2.13	0.47
1:A:254:ALA:HB1	1:A:446:LYS:HG2	1.97	0.47
1:C:101[A]:ARG:HG2	1:C:101[A]:ARG:HH11	1.79	0.47
1:C:470[A]:LEU:HD12	1:C:470[A]:LEU:C	2.40	0.46
1:A:278:THR:HG22	2:B:194[A]:ILE:HD12	1.98	0.46
1:C:337[B]:LEU:HD22	1:C:337[B]:LEU:C	2.41	0.46
2:D:7:VAL:HG11	2:D:129:PRO:HB2	1.97	0.46
1:A:233[A]:MET:CE	1:A:390:ASN:HD22	2.30	0.45
1:C:278:THR:HG22	2:D:194[B]:ILE:HD12	2.00	0.44
1:A:402:LYS:NZ	2:B:408:GLU:OE1	2.50	0.44
1:C:254:ALA:HB1	1:C:446:LYS:HG2	1.98	0.44
2:D:279:SER:HA	2:D:282:LYS:HD3	2.00	0.43
1:A:220:GLU:HG3	1:A:225:ALA:HB2	2.01	0.43
2:D:213[B]:MET:HE2	2:D:213[B]:MET:HB3	1.71	0.43
1:A:44[A]:TYR:CD1	1:A:44[A]:TYR:N	2.88	0.42
1:A:404:VAL:HG23	2:B:377[B]:MET:HE3	2.02	0.42
2:D:335:ARG:NH1	8:D:812:HOH:O	2.52	0.42
1:A:44[A]:TYR:HA	1:A:62[A]:ILE:HD13	2.02	0.41
1:A:430[B]:SER:HB2	1:A:433[B]:GLU:HB3	2.01	0.41
1:C:78:PHE:HB2	1:C:358:LEU:HD21	2.01	0.41
2:B:186[B]:ARG:HH11	2:B:186[B]:ARG:HD3	1.64	0.41
1:A:78:PHE:HB2	1:A:358:LEU:HD21	2.02	0.41
2:D:417[A]:ASP:OD1	2:D:421:ASN:ND2	2.53	0.41
1:A:66[B]:VAL:HG21	1:A:89:LEU:CD2	2.51	0.41
1:A:90[B]:GLU:CG	1:A:94:GLN:NE2	2.84	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:LEU:C	1:A:260:LEU:HD23	2.46	0.40
1:A:250:VAL:HG13	1:A:386[B]:ILE:CG1	2.49	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/495 (103%)	493 (97%)	15 (3%)	0	100	100
1	C	510/495 (103%)	497 (98%)	13 (2%)	0	100	100
2	B	475/450 (106%)	470 (99%)	5 (1%)	0	100	100
2	D	478/450 (106%)	470 (98%)	8 (2%)	0	100	100
3	E	3/5 (60%)	3 (100%)	0	0	100	100
3	F	3/5 (60%)	3 (100%)	0	0	100	100
All	All	1977/1900 (104%)	1936 (98%)	41 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/395 (105%)	412 (100%)	2 (0%)	81	62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	407/395 (103%)	405 (100%)	2 (0%)	81	62
2	B	381/360 (106%)	375 (98%)	6 (2%)	55	22
2	D	385/360 (107%)	380 (99%)	5 (1%)	61	29
All	All	1587/1510 (105%)	1572 (99%)	15 (1%)	76	43

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101[A]	ARG
1	A	101[B]	ARG
2	B	115[A]	GLU
2	B	115[B]	GLU
2	B	200	ASP
2	B	270	TYR
2	B	398[A]	GLU
2	B	398[B]	GLU
1	C	423[A]	LYS
1	C	423[B]	LYS
2	D	200	ASP
2	D	270	TYR
2	D	282	LYS
2	D	398[A]	GLU
2	D	398[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	GLN
1	A	349	ASN
1	A	357	HIS
1	A	474	ASN
2	B	57	ASN
2	B	150	GLN
1	C	27	GLN
1	C	64	GLN
1	C	94	GLN
1	C	349	ASN
1	C	357	HIS
1	C	474	ASN
2	D	57	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	356	GLN
2	D	379	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 23 ligands modelled in this entry, 6 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MES	C	502	-	12,12,12	0.74	0	15,16,16	0.51	0
6	EDO	D	702	-	3,3,3	0.59	0	2,2,2	0.24	0
6	EDO	C	503	-	3,3,3	0.44	0	2,2,2	0.46	0
6	EDO	B	702[B]	-	3,3,3	0.46	0	2,2,2	0.37	0
6	EDO	B	704	-	3,3,3	0.48	0	2,2,2	0.39	0
5	MES	A	503	-	12,12,12	0.70	0	15,16,16	0.43	0
6	EDO	D	704[B]	-	3,3,3	0.27	0	2,2,2	0.46	0
5	MES	D	701	-	12,12,12	0.63	0	15,16,16	0.47	0
6	EDO	D	705	-	3,3,3	0.54	0	2,2,2	0.23	0
6	EDO	B	702[A]	-	3,3,3	0.49	0	2,2,2	0.30	0
6	EDO	A	506	-	3,3,3	0.58	0	2,2,2	0.28	0
5	MES	A	502	-	12,12,12	0.80	0	15,16,16	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	504	-	3,3,3	0.57	0	2,2,2	0.34	0
5	MES	B	701	-	12,12,12	0.66	0	15,16,16	0.30	0
6	EDO	D	703	-	3,3,3	0.49	0	2,2,2	0.35	0
6	EDO	B	703	-	3,3,3	0.33	0	2,2,2	0.31	0
6	EDO	D	704[A]	-	3,3,3	0.30	0	2,2,2	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MES	C	502	-	-	0/6/14/14	0/1/1/1
6	EDO	D	702	-	-	0/1/1/1	-
6	EDO	C	503	-	-	0/1/1/1	-
6	EDO	B	702[B]	-	-	1/1/1/1	-
6	EDO	B	704	-	-	0/1/1/1	-
5	MES	A	503	-	-	2/6/14/14	0/1/1/1
6	EDO	D	704[B]	-	-	1/1/1/1	-
5	MES	D	701	-	-	0/6/14/14	0/1/1/1
6	EDO	D	705	-	-	1/1/1/1	-
6	EDO	B	702[A]	-	-	0/1/1/1	-
6	EDO	A	506	-	-	0/1/1/1	-
5	MES	A	502	-	-	0/6/14/14	0/1/1/1
6	EDO	A	504	-	-	0/1/1/1	-
5	MES	B	701	-	-	0/6/14/14	0/1/1/1
6	EDO	D	703	-	-	0/1/1/1	-
6	EDO	B	703	-	-	0/1/1/1	-
6	EDO	D	704[A]	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	MES	C8-C7-N4-C3
5	A	503	MES	C8-C7-N4-C5
6	B	702[B]	EDO	O1-C1-C2-O2
6	D	704[B]	EDO	O1-C1-C2-O2
6	D	705	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/495 (96%)	-1.01	0 100 100	7, 18, 26, 40	38 (7%)
1	C	480/495 (96%)	-0.97	0 100 100	7, 19, 28, 36	31 (6%)
2	B	444/450 (98%)	-1.10	0 100 100	5, 17, 26, 41	32 (7%)
2	D	445/450 (98%)	-1.14	0 100 100	5, 16, 25, 40	34 (7%)
3	E	5/5 (100%)	0.49	0 100 100	13, 14, 14, 22	5 (100%)
3	F	5/5 (100%)	0.39	0 100 100	13, 14, 14, 22	5 (100%)
All	All	1855/1900 (97%)	-1.04	0 100 100	5, 17, 27, 41	145 (7%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MES	A	503	12/12	0.93	0.11	17,19,28,34	12

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	D	704[A]	4/4	0.95	0.10	23,24,28,29	4
6	EDO	D	704[B]	4/4	0.95	0.10	13,19,22,24	4
6	EDO	B	704	4/4	0.97	0.07	29,35,37,40	0
5	MES	C	502	12/12	0.97	0.07	27,30,32,35	0
6	EDO	B	703	4/4	0.97	0.07	23,27,31,33	0
6	EDO	D	705	4/4	0.97	0.06	33,34,35,38	0
6	EDO	D	702	4/4	0.98	0.05	25,27,28,29	0
6	EDO	A	504	4/4	0.98	0.04	16,21,22,25	0
5	MES	A	502	12/12	0.99	0.05	18,20,21,22	0
5	MES	D	701	12/12	0.99	0.04	10,13,14,14	12
6	EDO	C	503	4/4	0.99	0.04	24,26,28,29	0
5	MES	B	701	12/12	0.99	0.04	12,14,15,15	12
6	EDO	D	703	4/4	0.99	0.04	24,25,25,25	0
6	EDO	A	506	4/4	0.99	0.04	20,20,21,24	0
6	EDO	B	702[A]	4/4	0.99	0.07	13,21,21,21	4
6	EDO	B	702[B]	4/4	0.99	0.07	22,22,24,26	4
7	NA	A	505	1/1	0.99	0.15	25,25,25,25	0
7	NA	B	705	1/1	0.99	0.11	29,29,29,29	0
4	ZN	C	501	1/1	1.00	0.01	17,17,17,17	0
4	ZN	A	501	1/1	1.00	0.02	16,16,16,16	0
7	NA	C	504	1/1	1.00	0.05	22,22,22,22	0
7	NA	D	706	1/1	1.00	0.07	25,25,25,25	0

6.5 Other polymers ⓘ

There are no such residues in this entry.