



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 06:50 AM UTC

PDB ID : 6NKQ / pdb_00006nkq
Title : The structure of bovine beta-lactoglobulin in novel crystals grown at pH 3.8
Authors : McPherson, A.
Deposited on : 2019-01-07
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	62% 25% • 11%
1	B	178	% 60% 28% •• 10%
1	C	178	% 66% 22% • 11%
1	D	178	54% 33% • 11%
1	E	178	67% 20% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	178	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: green (60%), yellow (27%), red (2%), and grey (11%). The red segment is marked with two dots (..).

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactoglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	160	Total	C	N	O	S	0	0	0
			1260	802	204	245	9			
1	C	159	Total	C	N	O	S	0	2	0
			1262	806	201	246	9			
1	D	159	Total	C	N	O	S	0	2	0
			1267	807	203	248	9			
1	E	159	Total	C	N	O	S	0	0	0
			1250	796	201	244	9			
1	F	159	Total	C	N	O	S	0	1	0
			1261	803	203	246	9			
1	A	158	Total	C	N	O	S	0	0	0
			1248	794	202	243	9			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	16	Total	O	0	0
			16	16		
3	C	12	Total	O	0	0
			12	12		
3	D	8	Total	O	0	1
			9	9		
3	E	7	Total	O	0	0
			7	7		
3	F	9	Total	O	0	0
			9	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-lactoglobulin




• Molecule 1: Beta-lactoglobulin




• Molecule 1: Beta-lactoglobulin




• Molecule 1: Beta-lactoglobulin





• Molecule 1: Beta-lactoglobulin



• Molecule 1: Beta-lactoglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	65.89Å 114.12Å 140.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.06 – 2.30 57.06 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (57.06-2.30) 99.5 (57.06-2.30)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	0.19	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.16 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.207 , 0.261 0.207 , 0.261	Depositor DCC
R_{free} test set	2370 reflections (4.50%)	wwPDB-VP
Wilson B-factor (Å ²)	42.2	Xtriage
Anisotropy	0.956	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 132.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.18$	Xtriage
Estimated twinning fraction	0.448 for -1/2*h+1/2*k,3/2*h+1/2*k,-l 0.439 for -1/2*h-1/2*k,-3/2*h+1/2*k,-l 0.447 for 1/2*h+1/2*k,3/2*h-1/2*k,-l 0.437 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.467 for -h,-k,l	Xtriage
Reported twinning fraction	0.162 for H, K, L 0.159 for -h,-k,l 0.164 for -1/2H-1/2K, 3/2H-1/2K, L 0.175 for 1/2H-1/2K, 3/2H+1/2K, L 0.158 for 1/2H+1/2K, -3/2H+1/2K, L 0.183 for -1/2H+1/2K, -3/2H-1/2K, L	Depositor
Outliers	1 of 49499 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7617	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/1269	1.27	0/1717
1	B	0.99	0/1281	1.30	0/1734
1	C	0.98	0/1289	1.28	0/1745
1	D	1.02	1/1294 (0.1%)	1.30	0/1751
1	E	0.98	0/1270	1.28	0/1719
1	F	1.08	1/1285 (0.1%)	1.33	3/1739 (0.2%)
All	All	1.01	2/7688 (0.0%)	1.29	3/10405 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	161	HIS	CG-ND1	-7.20	1.30	1.38
1	F	161	HIS	CG-ND1	-6.21	1.31	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	4	THR	CA-C-O	-6.04	114.84	121.31
1	F	58	LEU	CA-C-N	5.82	131.03	123.00
1	F	58	LEU	C-N-CA	5.82	131.03	123.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	158	GLU	Peptide
1	B	4	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1248	0	1259	47	0
1	B	1260	0	1270	55	0
1	C	1262	0	1277	48	1
1	D	1267	0	1280	87	2
1	E	1250	0	1263	38	1
1	F	1261	0	1274	70	2
2	B	1	0	0	0	0
3	A	15	0	0	7	0
3	B	16	0	0	7	0
3	C	12	0	0	8	0
3	D	9	0	0	3	0
3	E	7	0	0	3	0
3	F	9	0	0	3	0
All	All	7617	0	7623	320	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 320 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:GLN:O	1:B:161:HIS:HB2	1.17	1.32
1:D:60:LYS:HE3	1:D:61:TRP:N	1.48	1.26
1:F:131:GLU:OE2	3:F:201:HOH:O	1.56	1.21
1:F:155:GLN:O	1:F:161:HIS:HB3	1.43	1.13
1:B:155:GLN:O	1:B:161:HIS:CB	1.96	1.12

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:51:GLU:OE1	1:E:53:ASP:OD2[2_655]	2.11	0.09
1:D:53:ASP:OD2	1:F:51:GLU:OE1[2_656]	2.16	0.04
1:D:49:THR:OG1	1:F:51:GLU:OE2[2_656]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/178 (88%)	139 (89%)	15 (10%)	2 (1%)	9	10
1	B	158/178 (89%)	140 (89%)	17 (11%)	1 (1%)	21	27
1	C	159/178 (89%)	142 (89%)	16 (10%)	1 (1%)	21	27
1	D	159/178 (89%)	141 (89%)	16 (10%)	2 (1%)	9	10
1	E	157/178 (88%)	139 (88%)	16 (10%)	2 (1%)	9	10
1	F	158/178 (89%)	141 (89%)	16 (10%)	1 (1%)	21	27
All	All	947/1068 (89%)	842 (89%)	96 (10%)	9 (1%)	12	15

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	39	LEU
1	A	160	CYS
1	D	128	VAL
1	B	128	VAL
1	C	128	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/154 (90%)	135 (97%)	4 (3%)	37	55
1	B	140/154 (91%)	135 (96%)	5 (4%)	31	47
1	C	141/154 (92%)	140 (99%)	1 (1%)	76	87
1	D	142/154 (92%)	136 (96%)	6 (4%)	26	40
1	E	139/154 (90%)	135 (97%)	4 (3%)	37	55
1	F	141/154 (92%)	137 (97%)	4 (3%)	38	56
All	All	842/924 (91%)	818 (97%)	24 (3%)	37	55

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	128	VAL
1	F	47	LYS
1	F	4	THR
1	F	58	LEU
1	D	22	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	115	GLN
1	E	35	GLN
1	A	109	ASN
1	E	109	ASN
1	D	35	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/178 (88%)	-0.57	0 100 100	50, 101, 139, 175	1 (0%)
1	B	160/178 (89%)	-0.56	1 (0%) 85 86	42, 107, 143, 160	0
1	C	159/178 (89%)	-0.66	2 (1%) 75 76	53, 103, 144, 192	2 (1%)
1	D	159/178 (89%)	-0.58	0 100 100	33, 96, 136, 150	2 (1%)
1	E	159/178 (89%)	-0.57	0 100 100	62, 103, 136, 155	0
1	F	159/178 (89%)	-0.60	0 100 100	28, 92, 125, 144	1 (0%)
All	All	954/1068 (89%)	-0.59	3 (0%) 90 90	28, 100, 138, 192	6 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	139	ALA	2.5
1	B	126	PRO	2.2
1	C	156	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	B	201	1/1	0.98	0.05	111,111,111,111	1

6.5 Other polymers [i](#)

There are no such residues in this entry.