



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 05:50 AM UTC

PDB ID : 2NLK / pdb_00002nlk
Title : Crystal structure of D1 and D2 catalytic domains of human Protein Tyrosine Phosphatase Gamma (D1+D2 PTPRG)
Authors : Filippakopoulos, P.; Gileadi, O.; Johansson, C.; Ugochukwu, E.; Edwards, A.; Arrowsmith, C.; Sundstrom, M.; von Delft, F.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2006-10-20
Resolution : 2.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	627	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4536 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein tyrosine phosphatase, receptor type, G variant (Fragment).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	553	Total	C	N	O	S	0	0	0
			4386	2790	753	819	24			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	826	MET	-	initiating methionine	UNP Q59EE0
A	1446	ALA	-	cloning artifact	UNP Q59EE0
A	1447	HIS	-	expression tag	UNP Q59EE0
A	1448	HIS	-	expression tag	UNP Q59EE0
A	1449	HIS	-	expression tag	UNP Q59EE0
A	1450	HIS	-	expression tag	UNP Q59EE0
A	1451	HIS	-	expression tag	UNP Q59EE0
A	1452	HIS	-	expression tag	UNP Q59EE0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	150	Total	O	0	0
			150	150		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein tyrosine phosphatase, receptor type, G variant (Fragment)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.51Å 134.00Å 59.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.94 – 2.40 39.94 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.2 (39.94-2.40) 99.2 (39.94-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.237 , 0.274 0.234 , 0.268	Depositor DCC
R_{free} test set	1865 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4536	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	0/4486	0.97	5/6097 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1356	VAL	N-CA-C	6.09	116.84	110.62
1	A	1295	ILE	CB-CA-C	-5.62	102.46	110.83
1	A	906	TYR	N-CA-C	5.59	118.75	109.46
1	A	1390	ARG	CA-C-N	5.46	125.77	119.93
1	A	1390	ARG	C-N-CA	5.46	125.77	119.93

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1101	TYR	Peptide
1	A	1102	LEU	Peptide
1	A	1103	VAL	Peptide
1	A	1171	GLN	Peptide
1	A	1172	LYS	Peptide
1	A	1173	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4386	0	4215	43	0
2	A	150	0	0	1	0
All	All	4536	0	4215	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1096:ARG:HD2	1:A:1102:LEU:O	1.84	0.77
1:A:1096:ARG:CD	1:A:1102:LEU:O	2.39	0.71
1:A:1096:ARG:CZ	1:A:1102:LEU:O	2.40	0.69
1:A:1272:CYS:O	1:A:1275:PHE:O	2.10	0.69
1:A:855:VAL:HG13	1:A:1096:ARG:HD3	1.78	0.66
1:A:1096:ARG:NE	1:A:1102:LEU:O	2.31	0.63
1:A:1096:ARG:NH1	1:A:1102:LEU:O	2.31	0.63
1:A:1133:HIS:CE1	1:A:1408:LEU:HD22	2.35	0.61
1:A:1370:GLN:HG2	1:A:1376:ALA:O	2.01	0.60
1:A:970:SER:HG	1:A:979:THR:HG1	1.50	0.59
1:A:970:SER:OG	1:A:979:THR:OG1	2.19	0.58
1:A:890:ARG:HD3	2:A:57:HOH:O	2.03	0.58
1:A:1171:GLN:HA	1:A:1205:TYR:CE1	2.39	0.57
1:A:1064:VAL:HB	1:A:1101:TYR:O	2.05	0.57
1:A:1099:ARG:HB3	1:A:1102:LEU:HD11	1.85	0.57
1:A:855:VAL:CG1	1:A:1096:ARG:HD3	2.36	0.56
1:A:1411:VAL:O	1:A:1412:SER:C	2.50	0.53
1:A:1230:THR:O	1:A:1230:THR:CG2	2.57	0.53
1:A:1276:THR:O	1:A:1300:LEU:HA	2.09	0.53
1:A:1050:MET:O	1:A:1053:THR:HG22	2.10	0.51
1:A:1217:SER:O	1:A:1218:ASN:CB	2.55	0.51
1:A:1170:ALA:C	1:A:1172:LYS:H	2.20	0.50
1:A:1337:LYS:NZ	1:A:1368:SER:OG	2.43	0.49
1:A:1133:HIS:HA	1:A:1408:LEU:HD21	1.96	0.48
1:A:1099:ARG:HB3	1:A:1102:LEU:CD1	2.44	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1230:THR:O	1:A:1230:THR:HG22	2.13	0.47
1:A:1092:LEU:HD11	1:A:1103:VAL:H	1.80	0.47
1:A:1349:VAL:HG21	1:A:1361:LEU:HD22	1.96	0.47
1:A:1385:MET:HB3	1:A:1389:MET:HE2	1.97	0.47
1:A:1068:GLY:C	1:A:1103:VAL:HG12	2.40	0.46
1:A:1132:LEU:O	1:A:1136:VAL:HG23	2.15	0.45
1:A:1277:VAL:HG22	1:A:1300:LEU:HG	1.98	0.45
1:A:1238:ILE:CD1	1:A:1348:ILE:HG13	2.46	0.45
1:A:1068:GLY:O	1:A:1072:VAL:HG23	2.17	0.45
1:A:1170:ALA:C	1:A:1172:LYS:N	2.75	0.44
1:A:1314:PHE:CZ	1:A:1339:GLU:HG3	2.52	0.44
1:A:1175:ASN:HB3	1:A:1205:TYR:CD2	2.53	0.43
1:A:1212:MET:H	1:A:1389:MET:HE1	1.83	0.43
1:A:1133:HIS:NE2	1:A:1408:LEU:HD22	2.34	0.42
1:A:1377:VAL:CG1	1:A:1379:VAL:HG23	2.49	0.42
1:A:1333:ILE:HD11	1:A:1364:LEU:HB2	2.02	0.42
1:A:893:LEU:HD11	1:A:934:ASP:HB3	2.00	0.42
1:A:1314:PHE:HZ	1:A:1339:GLU:HG3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	541/627 (86%)	520 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	471/559 (84%)	465 (99%)	6 (1%)	61 80

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1103	VAL
1	A	1124	GLU
1	A	1140	LEU
1	A	1230	THR
1	A	1238	ILE
1	A	1377	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	836	HIS
1	A	868	HIS
1	A	952	ASN
1	A	1100	ASN
1	A	1289	ASN
1	A	1315	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	553/627 (88%)	0.46	36 (6%) 25 21	16, 25, 44, 55	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1204	ASP	4.7
1	A	1168	PHE	4.7
1	A	827	ALA	3.5
1	A	1169	SER	3.4
1	A	1155	LYS	3.3
1	A	1183	SER	3.3
1	A	897	PRO	3.2
1	A	1122	GLY	3.1
1	A	1196	PRO	3.1
1	A	1250	PRO	3.1
1	A	1129	SER	3.0
1	A	1176	LYS	2.9
1	A	1412	SER	2.8
1	A	1174	CYS	2.8
1	A	1102	LEU	2.8
1	A	1159	GLN	2.7
1	A	1170	ALA	2.6
1	A	1101	TYR	2.6
1	A	1172	LYS	2.6
1	A	1142	PRO	2.6
1	A	852	PHE	2.6
1	A	1124	GLU	2.6
1	A	828	ILE	2.5
1	A	916	ASN	2.4
1	A	1007	GLY	2.4
1	A	1133	HIS	2.4
1	A	1123	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1028	ASP	2.3
1	A	1292	GLN	2.3
1	A	1288	SER	2.2
1	A	1160	CYS	2.2
1	A	1186	PRO	2.2
1	A	1178	LYS	2.1
1	A	830	VAL	2.1
1	A	1184	VAL	2.0
1	A	1175	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.