



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:49 PM UTC

PDB ID : 2NN3 / pdb_00002nn3
Title : structure of pro-sf-caspase-1
Authors : Fisher, A.J.; Ni, L.
Deposited on : 2006-10-23
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	310	2% 43% 25% 6% 24%
1	D	310	2% 46% 30% 6% 16%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	235	Total	C	N	O	S	0	0	0
			1852	1186	314	338	14			
1	D	260	Total	C	N	O	S	0	0	0
			2014	1281	345	373	15			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	136	ALA	HIS	engineered mutation	UNP P89116
C	300	GLY	-	expression tag	UNP P89116
C	301	GLY	-	expression tag	UNP P89116
C	302	GLY	-	expression tag	UNP P89116
C	303	LEU	-	expression tag	UNP P89116
C	304	GLU	-	expression tag	UNP P89116
C	305	HIS	-	expression tag	UNP P89116
C	306	HIS	-	expression tag	UNP P89116
C	307	HIS	-	expression tag	UNP P89116
C	308	HIS	-	expression tag	UNP P89116
C	309	HIS	-	expression tag	UNP P89116
C	310	HIS	-	expression tag	UNP P89116
D	136	ALA	HIS	engineered mutation	UNP P89116
D	300	GLY	-	expression tag	UNP P89116
D	301	GLY	-	expression tag	UNP P89116
D	302	GLY	-	expression tag	UNP P89116
D	303	LEU	-	expression tag	UNP P89116
D	304	GLU	-	expression tag	UNP P89116
D	305	HIS	-	expression tag	UNP P89116
D	306	HIS	-	expression tag	UNP P89116
D	307	HIS	-	expression tag	UNP P89116
D	308	HIS	-	expression tag	UNP P89116
D	309	HIS	-	expression tag	UNP P89116
D	310	HIS	-	expression tag	UNP P89116


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	3	Total 3	O 3	0	0
2	D	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Caspase-1



• Molecule 1: Caspase-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	106.22Å 106.22Å 113.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.34 – 3.00 48.34 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.5 (48.34-3.00) 97.5 (48.34-3.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.228 , 0.275 0.228 , 0.225	Depositor DCC
R_{free} test set	722 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	108.1	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 139.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.049 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3872	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	1.32	6/1897 (0.3%)	1.07	3/2572 (0.1%)
1	D	1.35	8/2061 (0.4%)	1.16	13/2798 (0.5%)
All	All	1.33	14/3958 (0.4%)	1.12	16/5370 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	74	HIS	CE1-NE2	34.97	1.67	1.32
1	D	74	HIS	CE1-NE2	27.09	1.59	1.32
1	D	43	ARG	CZ-NH1	20.67	1.61	1.32
1	C	74	HIS	CG-ND1	19.24	1.59	1.38
1	D	74	HIS	CG-ND1	18.58	1.58	1.38

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	224	ARG	NE-CZ-NH2	-11.76	108.61	119.20
1	D	43	ARG	NE-CZ-NH2	-7.76	112.21	119.20
1	C	266	SER	N-CA-C	-7.05	102.96	112.26
1	C	215	SER	N-CA-C	6.70	125.07	110.80
1	D	74	HIS	ND1-CG-CD2	6.52	112.62	106.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	224	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1852	0	1797	82	1
1	D	2014	0	1943	81	1
2	C	3	0	0	0	0
2	D	3	0	0	0	0
All	All	3872	0	3740	143	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 143 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:74:HIS:NE2	1:C:74:HIS:CE1	1.67	1.43
1:D:44:MET:HB3	1:D:45:PRO:HD3	1.16	1.11
1:C:167:ALA:HB1	1:D:276:GLN:CB	1.90	1.02
1:D:44:MET:CB	1:D:45:PRO:HD3	1.96	0.94
1:D:44:MET:HB3	1:D:45:PRO:CD	1.96	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:223:TRP:O	1:D:218:PRO:O[5_555]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	231/310 (74%)	194 (84%)	26 (11%)	11 (5%)	2	11
1	D	258/310 (83%)	203 (79%)	42 (16%)	13 (5%)	1	10
All	All	489/620 (79%)	397 (81%)	68 (14%)	24 (5%)	1	10

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	265	GLU
1	C	278	LYS
1	D	44	MET
1	D	46	VAL
1	D	243	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	199/267 (74%)	174 (87%)	25 (13%)	4	20
1	D	214/267 (80%)	184 (86%)	30 (14%)	3	16
All	All	413/534 (77%)	358 (87%)	55 (13%)	4	18

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	74	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	101	THR
1	D	295	LYS
1	D	246	GLU
1	D	75	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	88	ASN
1	D	176	GLN
1	D	279	GLN
1	D	153	ASN
1	C	207	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	235/310 (75%)	0.04	4 (1%) 69 45	84, 101, 123, 138	0
1	D	260/310 (83%)	0.11	6 (2%) 61 38	82, 101, 137, 141	0
All	All	495/620 (79%)	0.08	10 (2%) 65 41	82, 101, 134, 141	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	75	SER	4.9
1	D	178	CYS	3.0
1	D	263	ASP	2.3
1	C	72	ASP	2.3
1	C	277	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.