



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:29 AM UTC

PDB ID : 5NOP / pdb_00005nop
Title : Structure of Mojiang virus attachment glycoprotein
Authors : Rissanen, I.R.; Ahmed, A.A.; Beaty, S.; Azarm, K.; Hong, P.; Nambulli, S.; Duprex, P.W.; Lee, B.; Bowden, T.A.
Deposited on : 2017-04-12
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	
1	B	471	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Attachment glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	423	Total	C	N	O	S	0	1	0
			3305	2100	542	638	25			
1	B	434	Total	C	N	O	S	0	2	0
			3378	2142	556	655	25			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	164	THR	-	expression tag	UNP W8TIR7
A	165	GLY	-	expression tag	UNP W8TIR7
A	626	GLY	-	expression tag	UNP W8TIR7
A	627	THR	-	expression tag	UNP W8TIR7
A	628	LYS	-	expression tag	UNP W8TIR7
A	629	HIS	-	expression tag	UNP W8TIR7
A	630	HIS	-	expression tag	UNP W8TIR7
A	631	HIS	-	expression tag	UNP W8TIR7
A	632	HIS	-	expression tag	UNP W8TIR7
A	633	HIS	-	expression tag	UNP W8TIR7
A	634	HIS	-	expression tag	UNP W8TIR7
B	164	THR	-	expression tag	UNP W8TIR7
B	165	GLY	-	expression tag	UNP W8TIR7
B	626	GLY	-	expression tag	UNP W8TIR7
B	627	THR	-	expression tag	UNP W8TIR7
B	628	LYS	-	expression tag	UNP W8TIR7
B	629	HIS	-	expression tag	UNP W8TIR7
B	630	HIS	-	expression tag	UNP W8TIR7
B	631	HIS	-	expression tag	UNP W8TIR7
B	632	HIS	-	expression tag	UNP W8TIR7
B	633	HIS	-	expression tag	UNP W8TIR7
B	634	HIS	-	expression tag	UNP W8TIR7

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	Cl 3	0	0
2	B	3	Total 3	Cl 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	492	Total 492	O 492	0	0
3	B	448	Total 448	O 448	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Attachment glycoprotein



• Molecule 1: Attachment glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.04Å 102.31Å 162.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.33 – 1.94 49.33 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.33-1.94) 99.3 (49.33-1.94)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.94Å)	Xtriage
Refinement program	PHENIX dev_2283	Depositor
R, R_{free}	0.182 , 0.222 0.182 , 0.223	Depositor DCC
R_{free} test set	3627 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.729	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7629	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/3389	0.47	0/4597
1	B	0.27	0/3464	0.47	0/4700
All	All	0.27	0/6853	0.47	0/9297

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3305	0	3200	29	0
1	B	3378	0	3265	17	0
2	A	3	0	0	0	0
2	B	3	0	0	1	0
3	A	492	0	0	5	0
3	B	448	0	0	3	0
All	All	7629	0	6465	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:475:VAL:HB	1:A:476:ARG:NH1	1.76	0.99
1:B:260:ASP:OD2	1:B:572:LYS:NZ	2.07	0.88
1:A:339:LEU:HG	1:B:339:LEU:HG	1.64	0.80
1:A:475:VAL:HB	1:A:476:ARG:HH12	1.51	0.76
1:A:474:SER:HB2	1:A:476:ARG:NH2	2.02	0.75
1:A:430:GLN:OE1	1:A:476:ARG:NH2	2.23	0.72
1:B:509:ARG:NH2	3:B:803:HOH:O	2.22	0.69
1:B:500:ASP:OD2	3:B:802:HOH:O	2.14	0.66
1:A:352:GLY:HA3	1:A:444:ARG:HB2	1.79	0.64
1:A:281:ILE:HD12	1:A:303:THR:HG22	1.81	0.60
1:A:475:VAL:HB	1:A:476:ARG:HH11	1.66	0.57
1:B:216:SER:HA	1:B:590:PRO:HG3	1.87	0.56
1:B:422:SER:HB3	2:B:701:CL:CL	2.43	0.55
1:B:286:ALA:HB1	1:B:353:GLY:HA2	1.87	0.55
1:A:361:LEU:HG	1:A:420:LEU:HD11	1.90	0.53
1:B:352:GLY:HA3	1:B:444:ARG:HB2	1.91	0.53
1:A:398:ARG:HA	1:A:401:MET:HE2	1.91	0.53
1:B:189:ASN:ND2	3:B:812:HOH:O	2.42	0.53
1:A:318:TRP:CE2	1:A:332:ARG:HD3	2.47	0.50
1:A:181:GLU:OE1	3:A:801:HOH:O	2.19	0.50
1:B:489:LYS:HD2	1:B:530[A]:THR:HG22	1.93	0.49
1:A:509:ARG:NH1	3:A:814:HOH:O	2.42	0.48
1:A:474:SER:HB2	1:A:476:ARG:HH21	1.74	0.48
1:A:476:ARG:CZ	1:A:476:ARG:H	2.26	0.48
1:B:302:LEU:HD23	1:B:315:ASP:HB2	1.95	0.48
1:A:401:MET:HE3	1:A:404:GLY:HA2	1.94	0.48
1:A:398:ARG:HG3	1:A:401:MET:HE2	1.94	0.47
1:A:500:ASP:OD2	3:A:802:HOH:O	2.20	0.47
1:A:177:ILE:HD13	1:A:258:ILE:HB	1.97	0.46
1:B:459:SER:HA	1:B:508:THR:O	2.16	0.46
1:A:372:ARG:NH2	3:A:825:HOH:O	2.49	0.45
1:B:318:TRP:CE2	1:B:332:ARG:HD3	2.52	0.45
1:A:417:VAL:HG22	1:A:427:ILE:HG12	1.98	0.45
1:A:199:PRO:HD3	1:A:542:VAL:HG23	1.99	0.45
1:B:370:ARG:HA	1:B:407:GLU:O	2.18	0.44
1:B:371:ASN:ND2	1:B:373:GLN:O	2.42	0.42
1:A:357:LYS:HD3	3:A:895:HOH:O	2.20	0.42
1:A:318:TRP:CZ2	1:A:332:ARG:HD3	2.55	0.41
1:A:323:GLU:HB3	1:A:326:THR:HB	2.03	0.41
1:A:223:PRO:HA	1:A:235:SER:O	2.20	0.41
1:A:523:TYR:HA	1:A:543:ARG:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:LEU:HD23	1:A:315:ASP:HB2	2.03	0.40
1:B:528:GLY:HA3	1:B:539:PHE:CZ	2.56	0.40
1:A:459:SER:HA	1:A:508:THR:O	2.22	0.40
1:B:318:TRP:CZ2	1:B:332:ARG:HD3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/471 (89%)	405 (96%)	15 (4%)	0	100	100
1	B	428/471 (91%)	412 (96%)	16 (4%)	0	100	100
All	All	848/942 (90%)	817 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/405 (90%)	364 (100%)	1 (0%)	86	86
1	B	374/405 (92%)	373 (100%)	1 (0%)	86	86
All	All	739/810 (91%)	737 (100%)	2 (0%)	86	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	604	CYS
1	B	229	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	A	263	GLN
1	A	371	ASN
1	A	418	ASN
1	A	538	ASN
1	A	548	HIS
1	B	430	GLN
1	B	538	ASN
1	B	589	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	423/471 (89%)	-0.21	9 (2%) 63 70	8, 17, 33, 68	1 (0%)
1	B	434/471 (92%)	-0.06	20 (4%) 37 42	9, 18, 39, 65	2 (0%)
All	All	857/942 (90%)	-0.13	29 (3%) 48 54	8, 18, 34, 68	3 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	403	PHE	4.5
1	A	312	HIS	4.3
1	B	176	THR	3.6
1	A	605	TYR	3.4
1	B	404	GLY	3.3
1	B	605	TYR	3.1
1	A	175	THR	3.1
1	B	405	SER	3.0
1	A	194	HIS	2.9
1	B	407	GLU	2.7
1	B	388	GLY	2.7
1	B	615	GLY	2.6
1	A	245	ALA	2.5
1	A	476	ARG	2.5
1	B	187	GLY	2.5
1	B	533	ASN	2.4
1	B	185	PRO	2.4
1	B	611	THR	2.4
1	B	612	VAL	2.4
1	B	189	ASN	2.3
1	B	229	SER	2.3
1	B	373	GLN	2.2
1	A	229	SER	2.1
1	B	312	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	325	ASP	2.1
1	A	533	ASN	2.1
1	B	184	LYS	2.1
1	B	476	ARG	2.0
1	B	406	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	703	1/1	0.83	0.12	45,45,45,45	0
2	CL	B	701	1/1	0.93	0.09	41,41,41,41	0
2	CL	B	702	1/1	0.93	0.10	50,50,50,50	0
2	CL	A	701	1/1	0.97	0.05	29,29,29,29	0
2	CL	A	702	1/1	0.99	0.06	24,24,24,24	0
2	CL	B	703	1/1	0.99	0.03	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.