



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 12:48 PM UTC

PDB ID : 2NPO / pdb_00002np0
Title : Crystal structure of putative transferase from *Campylobacter jejuni* subsp. *jejuni* NCTC 11168
Authors : Jin, X.; Bera, A.; Wasserman, S.; Smith, D.; Sauder, J.M.; Burley, S.K.; Shapiro, L.; New York SGX Research Center for Structural Genomics (NYS-GXRC)
Deposited on : 2006-10-27
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	13% 66% 16% 8% • 7%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	193	Total	C	N	O	S	0	0	0
			1460	938	242	270	10			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	UNP Q0P9D1
A	0	SER	-	cloning artifact	UNP Q0P9D1
A	1	LEU	-	cloning artifact	UNP Q0P9D1
A	196	GLU	-	cloning artifact	UNP Q0P9D1
A	197	GLY	-	cloning artifact	UNP Q0P9D1
A	198	GLY	-	cloning artifact	UNP Q0P9D1
A	199	SER	-	cloning artifact	UNP Q0P9D1
A	200	HIS	-	expression tag	UNP Q0P9D1
A	201	HIS	-	expression tag	UNP Q0P9D1
A	202	HIS	-	expression tag	UNP Q0P9D1
A	203	HIS	-	expression tag	UNP Q0P9D1
A	204	HIS	-	expression tag	UNP Q0P9D1
A	205	HIS	-	expression tag	UNP Q0P9D1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	141	Total	O	0	0
			141	141		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	87.14Å 87.14Å 65.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.20 50.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.20) 99.8 (50.00-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.95 (at 2.21Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.251 0.199 , 0.243	Depositor DCC
R_{free} test set	721 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.300	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.056 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1601	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	6/1482 (0.4%)	1.84	28/1991 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	12

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	196	GLU	C-N	18.46	1.59	1.33
1	A	194	ARG	C-N	9.94	1.46	1.33
1	A	194	ARG	CZ-NH1	5.60	1.40	1.32
1	A	195	MET	CA-C	5.55	1.60	1.52
1	A	193	LYS	C-N	5.50	1.41	1.33
1	A	194	ARG	CZ-NH2	5.17	1.40	1.33

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	VAL	CA-C-N	26.01	152.36	119.84
1	A	190	VAL	C-N-CA	26.01	152.36	119.84
1	A	58	ASN	N-CA-C	-23.03	86.27	111.14
1	A	59	GLU	N-CA-CB	-22.54	81.03	111.00
1	A	81	LYS	N-CA-C	-19.41	89.62	113.50
1	A	58	ASN	CB-CA-C	18.93	140.81	110.90
1	A	128	VAL	N-CA-C	15.29	131.18	108.54
1	A	182	ASP	N-CA-C	14.46	130.92	113.28
1	A	113	LYS	N-CA-C	-13.49	87.64	109.76
1	A	59	GLU	N-CA-C	-9.66	102.29	114.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	MET	CA-C-N	8.83	130.88	119.84
1	A	100	MET	C-N-CA	8.83	130.88	119.84
1	A	194	ARG	NE-CZ-NH1	-7.58	113.92	121.50
1	A	187	PHE	N-CA-C	-6.98	95.07	107.28
1	A	194	ARG	NH1-CZ-NH2	6.90	128.28	119.30
1	A	186	VAL	CA-C-N	6.88	131.52	121.26
1	A	186	VAL	C-N-CA	6.88	131.52	121.26
1	A	55	ILE	N-CA-C	-6.79	98.39	108.85
1	A	190	VAL	C-N-CD	-6.70	97.52	125.00
1	A	26	MET	N-CA-C	-6.12	102.61	110.19
1	A	190	VAL	N-CA-C	-5.82	101.19	108.05
1	A	57	ASN	N-CA-C	5.78	123.11	110.80
1	A	128	VAL	N-CA-CB	-5.70	104.57	110.95
1	A	127	CYS	CA-C-N	5.23	129.12	121.80
1	A	127	CYS	C-N-CA	5.23	129.12	121.80
1	A	187	PHE	N-CA-CB	5.19	120.51	111.74
1	A	81	LYS	CA-C-N	5.13	128.10	120.31
1	A	81	LYS	C-N-CA	5.13	128.10	120.31

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	100	MET	Peptide
1	A	112	GLU	Peptide
1	A	127	CYS	Peptide
1	A	181	GLN	Peptide
1	A	186	VAL	Peptide
1	A	190	VAL	Peptide
1	A	55	ILE	Peptide
1	A	56	GLY	Peptide
1	A	57	ASN	Peptide
1	A	58	ASN	Peptide
1	A	59	GLU	Peptide
1	A	80	HIS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1460	0	1498	55	1
2	A	141	0	0	10	0
All	All	1601	0	1498	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:LYS:HD3	1:A:38:LYS:H	1.11	1.14
1:A:58:ASN:O	1:A:61:ARG:HB2	1.55	1.02
1:A:38:LYS:H	1:A:38:LYS:CD	1.69	0.99
1:A:55:ILE:HG22	1:A:56:GLY:HA3	1.45	0.98
1:A:38:LYS:CD	1:A:38:LYS:N	2.30	0.93
1:A:61:ARG:HH11	1:A:61:ARG:HG3	1.37	0.90
1:A:66:GLN:O	1:A:70:GLU:HG2	1.73	0.88
1:A:38:LYS:HD3	1:A:38:LYS:N	1.90	0.86
1:A:26:MET:HE2	1:A:78:LEU:HD22	1.60	0.83
1:A:26:MET:HE1	1:A:78:LEU:HD23	1.66	0.76
1:A:195:MET:SD	2:A:323:HOH:O	2.45	0.75
1:A:102:TYR:CE1	2:A:264:HOH:O	2.40	0.74
1:A:57:ASN:ND2	1:A:59:GLU:HB3	2.04	0.72
1:A:58:ASN:O	1:A:61:ARG:CB	2.36	0.72
1:A:61:ARG:HH11	1:A:61:ARG:CG	2.03	0.72
1:A:26:MET:CE	1:A:78:LEU:CD2	2.67	0.71
1:A:92:GLU:OE1	2:A:340:HOH:O	2.08	0.70
1:A:4:THR:CG2	1:A:6:LYS:O	2.41	0.69
1:A:41:LYS:C	2:A:313:HOH:O	2.36	0.68
1:A:26:MET:HE2	1:A:78:LEU:CD2	2.23	0.68
1:A:26:MET:CE	1:A:78:LEU:HD23	2.25	0.67
1:A:55:ILE:O	1:A:61:ARG:HG3	1.96	0.66
1:A:112:GLU:OE1	2:A:324:HOH:O	2.12	0.66
1:A:26:MET:CE	1:A:78:LEU:HD22	2.27	0.65
1:A:57:ASN:HD22	1:A:60:ILE:H	1.45	0.64
1:A:55:ILE:HG22	1:A:56:GLY:CA	2.26	0.63
1:A:4:THR:HG21	1:A:50:ASP:H	1.64	0.62
1:A:57:ASN:O	1:A:58:ASN:C	2.43	0.61
1:A:80:HIS:ND1	1:A:82:SER:HB2	2.14	0.61
1:A:5:GLU:OE1	2:A:245:HOH:O	2.17	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:ILE:CG2	1:A:56:GLY:HA3	2.29	0.59
1:A:63:LYS:HZ1	1:A:66:GLN:HE22	1.50	0.58
1:A:4:THR:HG22	1:A:6:LYS:O	2.04	0.57
1:A:38:LYS:N	1:A:38:LYS:HD2	2.17	0.56
1:A:158:CYS:SG	1:A:176:THR:HB	2.46	0.56
1:A:183:GLU:HG2	2:A:296:HOH:O	2.07	0.54
1:A:182:ASP:OD2	1:A:182:ASP:N	2.39	0.54
1:A:26:MET:HE3	1:A:28:TYR:HE1	1.73	0.53
1:A:57:ASN:ND2	1:A:60:ILE:H	2.05	0.53
1:A:51:PHE:CE1	1:A:68:ILE:HD13	2.43	0.53
1:A:118:ASN:HD22	1:A:134:HIS:HE1	1.57	0.52
1:A:53:ILE:CD1	1:A:65:TYR:HA	2.40	0.52
1:A:55:ILE:O	1:A:61:ARG:CG	2.58	0.51
1:A:63:LYS:NZ	1:A:66:GLN:HE22	2.08	0.51
1:A:79:ILE:HG12	1:A:99:ILE:HD12	1.92	0.50
1:A:160:LEU:HB3	1:A:161:PRO:HD2	1.94	0.49
1:A:193:LYS:NZ	2:A:315:HOH:O	2.48	0.47
1:A:61:ARG:CG	1:A:61:ARG:NH1	2.68	0.46
1:A:171:LEU:HB2	2:A:344:HOH:O	2.16	0.45
1:A:4:THR:CG2	1:A:50:ASP:H	2.30	0.42
1:A:57:ASN:ND2	1:A:60:ILE:HD12	2.36	0.41
1:A:63:LYS:NZ	1:A:66:GLN:NE2	2.69	0.40
1:A:57:ASN:HD21	1:A:59:GLU:HB3	1.81	0.40
1:A:18:VAL:HG13	1:A:100:MET:HE1	2.03	0.40
1:A:38:LYS:NZ	2:A:325:HOH:O	1.94	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:SER:O	1:A:102:TYR:OH[2_665]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	189/207 (91%)	178 (94%)	8 (4%)	3 (2%)	7 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	191	PRO
1	A	101	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	159/172 (92%)	145 (91%)	14 (9%)	9 10

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	18	VAL
1	A	38	LYS
1	A	40	MET
1	A	61	ARG
1	A	63	LYS
1	A	82	SER
1	A	119	THR
1	A	128	VAL
1	A	160	LEU
1	A	176	THR
1	A	182	ASP
1	A	190	VAL
1	A	193	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	66	GLN
1	A	71	ASN
1	A	134	HIS
1	A	162	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	193/207 (93%)	1.02	26 (13%) 7 5	21, 34, 46, 57	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	197	GLY	7.3
1	A	196	GLU	4.7
1	A	56	GLY	4.5
1	A	55	ILE	4.0
1	A	57	ASN	3.9
1	A	70	GLU	3.6
1	A	58	ASN	3.5
1	A	59	GLU	3.2
1	A	37	PHE	3.1
1	A	181	GLN	3.1
1	A	195	MET	3.1
1	A	102	TYR	3.0
1	A	194	ARG	2.8
1	A	82	SER	2.7
1	A	41	LYS	2.6
1	A	11	GLY	2.6
1	A	38	LYS	2.5
1	A	27	GLY	2.5
1	A	142	ALA	2.2
1	A	101	PRO	2.1
1	A	176	THR	2.1
1	A	182	ASP	2.1
1	A	4	THR	2.1
1	A	160	LEU	2.1
1	A	159	VAL	2.1
1	A	192	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.