



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 11:46 AM UTC

PDB ID : 4NPH / pdb_00004nph
Title : Crystal structure of SsaN from Salmonella enterica
Authors : Sugiman-Marangos, S.N.; Zhang, K.; Cooper, C.A.; Coombes, B.K.; Junop, M.S.
Deposited on : 2013-11-21
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable secretion system apparatus ATP synthase SsaN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	0	0
			2441	1539	443	450	9			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	89	MET	-	initiating methionine	UNP P74857
A	434	HIS	-	expression tag	UNP P74857
A	435	HIS	-	expression tag	UNP P74857
A	436	HIS	-	expression tag	UNP P74857
A	437	HIS	-	expression tag	UNP P74857
A	438	HIS	-	expression tag	UNP P74857
A	439	HIS	-	expression tag	UNP P74857


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	157	Total	O	0	0
			157	157		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable secretion system apparatus ATP synthase SsaN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	138.14Å 76.31Å 39.09Å 90.00° 103.71° 90.00°	Depositor
Resolution (Å)	37.03 – 2.09 37.03 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.4 (37.03-2.09) 97.3 (37.03-2.09)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.08Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.191 , 0.229 0.193 , 0.229	Depositor DCC
R_{free} test set	2017 reflections (8.59%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.036 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2598	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	2/2475 (0.1%)	0.87	3/3359 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	GLN	C-N	5.06	1.40	1.33
1	A	134	PRO	N-CD	5.02	1.54	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	MET	N-CA-C	6.94	118.50	111.07
1	A	133	GLN	CA-C-N	-5.93	113.86	119.85
1	A	133	GLN	C-N-CA	-5.93	113.86	119.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2441	0	2446	57	1
2	A	157	0	0	23	1
All	All	2598	0	2446	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 12.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:GLY:HA2	1:A:297:LYS:O	1.35	1.23
1:A:93:VAL:HG11	1:A:239:PHE:CG	1.78	1.17
1:A:309:GLU:CB	2:A:627:HOH:O	1.94	1.13
1:A:157:ARG:HD3	2:A:605:HOH:O	1.51	1.10
1:A:270:GLU:N	2:A:632:HOH:O	1.87	1.07
1:A:93:VAL:CG1	1:A:239:PHE:CG	2.40	1.04
1:A:90:GLN:N	2:A:641:HOH:O	1.91	1.03
1:A:93:VAL:CG1	1:A:239:PHE:CD2	2.45	0.99
1:A:298:GLY:N	2:A:655:HOH:O	1.90	0.99
1:A:309:GLU:N	2:A:627:HOH:O	1.87	0.97
1:A:93:VAL:HG11	1:A:239:PHE:CB	1.95	0.96
1:A:93:VAL:HG12	1:A:239:PHE:CD2	2.00	0.96
1:A:157:ARG:NH1	2:A:605:HOH:O	1.97	0.94
1:A:270:GLU:CB	2:A:632:HOH:O	2.15	0.94
1:A:332:SER:OG	1:A:334:ARG:HG2	1.70	0.91
1:A:163:ALA:O	2:A:544:HOH:O	1.88	0.89
1:A:244:GLY:CA	1:A:297:LYS:O	2.25	0.82
1:A:256:ARG:NE	2:A:645:HOH:O	2.14	0.81
1:A:168:LYS:NZ	2:A:647:HOH:O	1.94	0.79
1:A:157:ARG:CD	2:A:605:HOH:O	2.16	0.76
1:A:241:ARG:NH2	2:A:560:HOH:O	2.15	0.76
1:A:256:ARG:CZ	2:A:645:HOH:O	2.38	0.71
1:A:157:ARG:CZ	2:A:605:HOH:O	2.37	0.67
1:A:149:VAL:HG21	1:A:346:VAL:HG13	1.78	0.66
1:A:152:CYS:SG	2:A:582:HOH:O	2.52	0.66
1:A:93:VAL:CG1	1:A:239:PHE:CB	2.71	0.66
1:A:334:ARG:HE	1:A:334:ARG:H	1.43	0.65
1:A:390:ARG:C	1:A:392:VAL:H	2.09	0.60
1:A:333:ARG:H	1:A:334:ARG:HH21	1.52	0.57
1:A:93:VAL:HG11	1:A:239:PHE:HB2	1.83	0.56
1:A:228:ARG:NH2	2:A:641:HOH:O	2.38	0.54
1:A:289:LEU:HD13	1:A:325:LEU:HD21	1.87	0.54
1:A:334:ARG:H	1:A:334:ARG:NE	2.06	0.54
1:A:308:VAL:HB	1:A:309:GLU:HA	1.90	0.54
1:A:100:ARG:NH1	1:A:115:PRO:HD3	2.24	0.53
1:A:361:GLU:OE2	1:A:428:HIS:NE2	2.26	0.52
1:A:377:GLN:NE2	2:A:653:HOH:O	2.24	0.52
1:A:93:VAL:HG12	1:A:239:PHE:CE2	2.45	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:GLU:HB2	2:A:580:HOH:O	2.10	0.50
1:A:338:ARG:NH1	2:A:640:HOH:O	2.44	0.49
1:A:213:VAL:HG21	1:A:236:ILE:HG21	1.93	0.49
1:A:155:GLY:O	1:A:293:GLY:HA2	2.13	0.49
1:A:103:ASP:HB3	1:A:109:LEU:HD11	1.95	0.48
1:A:194:ARG:O	1:A:198:GLU:HG2	2.14	0.47
1:A:390:ARG:C	1:A:392:VAL:N	2.73	0.47
1:A:332:SER:HG	1:A:334:ARG:HG2	1.75	0.45
1:A:185:VAL:HG22	1:A:213:VAL:CG2	2.47	0.45
1:A:255:THR:HA	1:A:317:LEU:CD2	2.48	0.43
1:A:93:VAL:CG1	1:A:239:PHE:HB3	2.49	0.43
1:A:256:ARG:NH1	2:A:645:HOH:O	2.46	0.43
1:A:207:GLU:H	1:A:207:GLU:CD	2.29	0.41
1:A:422:LEU:HG	1:A:426:LYS:HD2	2.01	0.41
1:A:269:GLY:C	2:A:632:HOH:O	2.47	0.41
1:A:403:TYR:HB3	1:A:404:PRO:HD3	2.03	0.41
1:A:97:LEU:C	1:A:213:VAL:HG12	2.45	0.41
1:A:254:LEU:HG	1:A:304:TYR:HB3	2.02	0.41
1:A:157:ARG:NE	2:A:605:HOH:O	2.41	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:VAL:CG2	2:A:612:HOH:O[1_554]	1.90	0.30

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/351 (90%)	312 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	250/293 (85%)	246 (98%)	4 (2%)	55 64

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ASP
1	A	131	VAL
1	A	307	LEU
1	A	334	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	328	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/351 (92%)	0.51	29 (8%) 15 16	20, 39, 80, 115	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	93	VAL	7.0
1	A	129	ALA	6.9
1	A	272	ALA	5.9
1	A	131	VAL	4.4
1	A	271	THR	4.3
1	A	120	LYS	3.9
1	A	118	CYS	3.6
1	A	315	GLU	3.5
1	A	432	THR	3.5
1	A	130	MET	3.4
1	A	314	ASN	3.4
1	A	111	GLY	3.3
1	A	116	ASP	3.3
1	A	114	LEU	3.3
1	A	394	THR	3.1
1	A	388	TYR	3.1
1	A	112	ARG	3.0
1	A	393	ASP	2.9
1	A	295	GLY	2.7
1	A	105	PHE	2.5
1	A	220	ASP	2.5
1	A	90	GLN	2.5
1	A	297	LYS	2.3
1	A	296	GLU	2.2
1	A	113	GLU	2.2
1	A	243	ASN	2.2
1	A	392	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	107	ARG	2.1
1	A	280	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.