



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:39 PM UTC

PDB ID : 4NQC / pdb_00004nqc
Title : Crystal structure of TCR-MR1 ternary complex and covalently bound 5-(2-oxopropylideneamino)-6-D-ribitylaminouracil
Authors : Birkinshaw, R.W.; Rossjohn, J.
Deposited on : 2013-11-25
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	85% 9% 6%
1	C	271	88% 8% ..
2	B	99	3% 88% 8% ..
2	F	99	89% 8% ..
3	D	203	7% 83% 7% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	203	90% 7%
4	E	245	2% 90% 8%
4	H	245	% 91% 9%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 13506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	1	0
			2098	1342	367	378	11			
1	C	263	Total	C	N	O	S	0	0	0
			2149	1377	370	391	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP Q95460
A	261	SER	CYS	engineered mutation	UNP Q95460
C	0	MET	-	expression tag	UNP Q95460
C	261	SER	CYS	engineered mutation	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	96	Total	C	N	O	S	0	0	0
			757	487	129	139	2			
2	F	97	Total	C	N	O	S	0	0	0
			774	497	132	143	2			


- Molecule 3 is a protein called TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	185	Total	C	N	O	S	0	0	0
			1376	888	220	260	8			
3	G	198	Total	C	N	O	S	0	0	0
			1516	960	242	305	9			

- Molecule 4 is a protein called TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	240	Total	C	N	O	S	0	1	0
			1851	1169	317	356	9			
4	H	243	Total	C	N	O	S	0	2	0
			1900	1195	331	365	9			

- Molecule 5 is 1-deoxy-1-({2,6-dioxo-5-[(E)-propylideneamino]-1,2,3,6-tetrahydropyrimidin-4-yl}amino)-D-ribitol (CCD ID: 2LJ) (formula: C₁₂H₂₀N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			22	12	4	6		
5	C	1	Total	C	N	O	0	0
			22	12	4	6		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	1	Total	Na	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	174	Total	O	0	0
			174	174		
7	B	56	Total	O	0	0
			56	56		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	194	Total 194	O 194	0	0
7	D	87	Total 87	O 87	0	0
7	E	139	Total 139	O 139	0	0
7	F	67	Total 67	O 67	0	0
7	G	143	Total 143	O 143	0	0
7	H	180	Total 180	O 180	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 3: TCR alpha chain





- Molecule 3: TCR alpha chain

Chain G: 90% 7% .



- Molecule 4: TCR beta chain

Chain E: 90% 2% 8% .



- Molecule 4: TCR beta chain

Chain H: 91% 9% .



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	218.76Å 71.11Å 144.28Å 90.00° 104.87° 90.00°	Depositor
Resolution (Å)	33.42 – 2.50 33.42 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.42-2.50) 99.9 (33.42-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.165 , 0.216 0.172 , 0.226	Depositor DCC
R_{free} test set	3757 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.663	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13506	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 2LJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.84	1/2160 (0.0%)	1.26	10/2933 (0.3%)
1	C	0.86	1/2213 (0.0%)	1.26	10/3008 (0.3%)
2	B	0.76	0/780	1.20	4/1066 (0.4%)
2	F	0.78	0/797	1.21	4/1087 (0.4%)
3	D	0.85	1/1407 (0.1%)	1.15	3/1915 (0.2%)
3	G	0.87	1/1550 (0.1%)	1.16	6/2106 (0.3%)
4	E	0.78	0/1901	1.18	7/2595 (0.3%)
4	H	0.84	0/1951	1.20	5/2659 (0.2%)
All	All	0.83	4/12759 (0.0%)	1.21	49/17369 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	14	ASP	CA-C	7.72	1.58	1.53
1	A	14	ASP	CA-C	7.57	1.58	1.53
3	D	111	ILE	CG1-CD1	-5.43	1.30	1.51
3	G	111	ILE	CG1-CD1	-5.28	1.31	1.51

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	117	ASP	CA-CB-CG	8.07	120.67	112.60
1	C	222	VAL	N-CA-C	-7.89	105.80	113.53
4	H	101	SER	N-CA-C	6.99	121.56	112.89
4	E	101	SER	N-CA-C	6.97	121.53	112.89
4	H	100	GLY	N-CA-C	6.60	121.22	112.77
2	B	34	ASP	CA-CB-CG	6.43	119.03	112.60
4	H	117	ASP	CA-CB-CG	6.32	118.92	112.60
4	H	73	GLU	N-CA-C	6.25	119.66	109.59
1	C	175	THR	CA-C-N	6.19	129.72	120.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	175	THR	C-N-CA	6.19	129.72	120.31
1	C	102	GLU	CB-CG-CD	6.03	122.85	112.60
3	G	123	ASP	CA-CB-CG	6.01	118.61	112.60
2	F	4	THR	CB-CA-C	5.89	117.93	109.26
1	A	28	VAL	N-CA-CB	5.89	119.09	111.67
4	E	73	GLU	N-CA-C	5.80	118.92	109.59
3	D	148	ASP	N-CA-C	5.79	118.05	108.49
2	F	33	SER	N-CA-C	5.71	118.44	111.82
1	A	56	PRO	CA-C-N	5.67	128.15	120.38
1	A	56	PRO	C-N-CA	5.67	128.15	120.38
4	E	125	GLU	N-CA-C	-5.58	100.15	109.24
1	C	56	PRO	CA-C-N	5.56	128.00	120.38
1	C	56	PRO	C-N-CA	5.56	128.00	120.38
4	E	97	THR	CB-CA-C	5.54	121.14	109.79
1	A	134	ASN	CA-CB-CG	-5.50	107.10	112.60
4	E	100	GLY	N-CA-C	5.45	119.74	112.77
2	B	33	SER	N-CA-C	5.37	118.05	111.82
3	G	175	SER	CA-C-N	5.34	131.35	121.52
3	G	175	SER	C-N-CA	5.34	131.35	121.52
4	H	125	GLU	N-CA-C	-5.31	100.58	109.24
2	B	21	ASN	CA-CB-CG	5.29	117.89	112.60
2	F	21	ASN	CA-CB-CG	5.26	117.86	112.60
2	B	71	THR	CB-CA-C	5.22	115.65	110.33
4	E	204	ASN	CA-CB-CG	5.21	117.81	112.60
1	A	175	THR	CA-C-N	5.20	128.22	120.31
1	A	175	THR	C-N-CA	5.20	128.22	120.31
1	C	74	LYS	CA-C-N	5.19	127.09	120.56
1	C	74	LYS	C-N-CA	5.19	127.09	120.56
1	A	74	LYS	CA-C-N	5.18	127.08	120.56
1	A	74	LYS	C-N-CA	5.18	127.08	120.56
1	A	31	HIS	N-CA-C	5.17	116.43	109.24
1	C	31	HIS	N-CA-C	5.17	116.42	109.24
3	G	137	PHE	CA-C-N	5.16	128.54	120.75
3	G	137	PHE	C-N-CA	5.16	128.54	120.75
2	F	71	THR	CB-CA-C	5.13	115.57	110.33
3	D	149	SER	CA-C-N	5.13	130.94	121.70
3	D	149	SER	C-N-CA	5.13	130.94	121.70
1	A	133	ASP	CA-CB-CG	5.11	117.71	112.60
3	G	142	ASN	CA-CB-CG	5.10	117.70	112.60
1	C	28	VAL	N-CA-CB	5.01	117.98	111.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2098	0	1979	9	0
1	C	2149	0	2020	9	0
2	B	757	0	694	3	0
2	F	774	0	717	3	0
3	D	1376	0	1259	7	0
3	G	1516	0	1407	6	0
4	E	1851	0	1713	5	0
4	H	1900	0	1762	8	0
5	A	22	0	18	0	0
5	C	22	0	18	0	0
6	H	1	0	0	0	0
7	A	174	0	0	1	0
7	B	56	0	0	0	0
7	C	194	0	0	2	0
7	D	87	0	0	0	0
7	E	139	0	0	0	0
7	F	67	0	0	0	0
7	G	143	0	0	0	0
7	H	180	0	0	0	0
All	All	13506	0	11587	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:149:SER:HA	3:D:151:VAL:H	1.45	0.82
2:F:4:THR:HA	2:F:86:THR:HG21	1.69	0.72
1:C:257:HIS:ND1	7:C:585:HOH:O	2.19	0.71
1:C:198:LEU:HG	1:C:246:LEU:HD11	1.74	0.68
2:F:84:HIS:ND1	2:F:86:THR:HG22	2.10	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:HIS:ND1	7:A:557:HOH:O	2.31	0.61
3:G:159:LEU:HB3	4:H:172:CYS:HB2	1.85	0.58
3:G:164:MET:HE1	4:H:198:SER:HB3	1.86	0.58
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.42	0.54
4:H:21:LEU:HD12	4:H:76:LEU:HD23	1.91	0.53
3:G:28:GLY:HA3	3:G:93:SER:HB3	1.92	0.52
4:E:26:ASP:OD1	4:E:72:ARG:NE	2.36	0.51
3:D:28:GLY:HA3	3:D:93:SER:HB3	1.92	0.51
1:A:215:MET:HG3	1:A:257:HIS:CD2	2.46	0.50
1:A:8:PHE:CE2	1:A:95:MET:HG3	2.47	0.50
1:C:77:LEU:HD13	1:C:92:TYR:HB2	1.95	0.48
4:E:27:MET:HE1	4:E:105:PHE:CD2	2.49	0.48
1:A:46:ARG:HD3	1:A:46:ARG:HA	1.75	0.47
1:C:246:LEU:HD12	1:C:246:LEU:N	2.30	0.47
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.97	0.46
3:G:152:TYR:O	3:G:173:ALA:HA	2.16	0.45
1:C:46:ARG:HA	1:C:46:ARG:HD3	1.78	0.44
4:E:83:PRO:HA	4:E:114:VAL:HB	1.99	0.44
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.53	0.44
3:D:159:LEU:HB3	4:E:172:CYS:HB2	1.99	0.44
1:C:113:ALA:HB2	2:F:60:TRP:CE2	2.52	0.44
1:A:16:ILE:HD13	2:B:34:ASP:OD2	2.18	0.43
1:C:94:ARG:HD3	7:C:416:HOH:O	2.18	0.43
4:H:27:MET:HE1	4:H:105:PHE:CD2	2.53	0.43
1:A:154:LYS:HD3	3:G:51:LEU:HD11	2.01	0.43
3:G:158:VAL:HG22	3:G:169:ASN:OD1	2.19	0.43
3:D:50:VAL:O	3:D:66:ARG:HD3	2.20	0.42
4:H:83:PRO:HA	4:H:114:VAL:HB	2.01	0.42
4:H:122:PHE:CD2	4:H:188:ARG:HD3	2.55	0.42
3:D:134:PHE:O	3:D:170:SER:HA	2.20	0.41
2:B:24:ASN:HB3	2:B:65:LEU:HD11	2.02	0.41
1:C:61:ARG:HG2	3:D:94:ASN:HB3	2.02	0.41
1:A:43:LYS:HD2	1:A:62:TYR:HB3	2.03	0.41
4:E:121:VAL:HG12	4:E:231:PRO:HB2	2.03	0.41
3:D:152:TYR:O	3:D:173:ALA:HA	2.21	0.41
4:H:150:GLY:O	4:H:188:ARG:HD2	2.21	0.41
4:H:128:VAL:HG23	4:H:238:ALA:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/271 (92%)	238 (95%)	11 (4%)	1 (0%)	30	49
1	C	257/271 (95%)	250 (97%)	7 (3%)	0	100	100
2	B	94/99 (95%)	93 (99%)	1 (1%)	0	100	100
2	F	95/99 (96%)	94 (99%)	1 (1%)	0	100	100
3	D	179/203 (88%)	176 (98%)	3 (2%)	0	100	100
3	G	196/203 (97%)	192 (98%)	4 (2%)	0	100	100
4	E	239/245 (98%)	234 (98%)	5 (2%)	0	100	100
4	H	243/245 (99%)	235 (97%)	8 (3%)	0	100	100
All	All	1553/1636 (95%)	1512 (97%)	40 (3%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	220	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/241 (91%)	214 (97%)	6 (3%)	39	67
1	C	225/241 (93%)	217 (96%)	8 (4%)	31	58
2	B	80/94 (85%)	78 (98%)	2 (2%)	42	69
2	F	83/94 (88%)	80 (96%)	3 (4%)	31	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	D	136/180 (76%)	132 (97%)	4 (3%)	37	65
3	G	165/180 (92%)	162 (98%)	3 (2%)	51	77
4	E	195/211 (92%)	192 (98%)	3 (2%)	57	80
4	H	200/211 (95%)	195 (98%)	5 (2%)	42	69
All	All	1304/1452 (90%)	1270 (97%)	34 (3%)	42	68

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	MET
1	A	16	ILE
1	A	73	PHE
1	A	78	LYS
1	A	82	ARG
1	A	183	LEU
2	B	4	THR
2	B	35	ILE
1	C	0	MET
1	C	73	PHE
1	C	78	LYS
1	C	102	GLU
1	C	111	GLN
1	C	183	LEU
1	C	215	MET
1	C	221	ILE
3	D	2	GLN
3	D	85	SER
3	D	111	ILE
3	D	157	CYS
4	E	79	GLU
4	E	80	SER
4	E	237	SER
2	F	4	THR
2	F	16	GLU
2	F	35	ILE
3	G	85	SER
3	G	111	ILE
3	G	163	SER
4	H	140[A]	GLN
4	H	140[B]	GLN
4	H	206[A]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	H	206[B]	ARG
4	H	237	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	137	HIS
1	A	203	HIS
2	B	13	HIS
1	C	64	GLN
1	C	93	GLN
1	C	111	GLN
1	C	239	GLN
3	D	37	GLN
4	E	37	GLN
4	E	62	ASN
4	E	120	ASN
4	E	214	GLN
3	G	19	GLN
3	G	21	ASN
3	G	187	ASN
4	H	62	ASN
4	H	65	ASN
4	H	214	GLN
4	H	226	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	2LJ	C	301	1	22,22,22	1.29	1 (4%)	24,29,29	1.14	2 (8%)
5	2LJ	A	301	1	22,22,22	1.22	1 (4%)	24,29,29	1.39	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2LJ	C	301	1	-	1/18/19/19	0/1/1/1
5	2LJ	A	301	1	-	1/18/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	2LJ	C8A-N8	5.45	1.41	1.32
5	A	301	2LJ	C8A-N8	4.83	1.40	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	301	2LJ	C6-N5-C4A	4.26	127.23	121.17
5	C	301	2LJ	C6-N5-C4A	3.19	125.70	121.17
5	A	301	2LJ	C8-C7-C6	2.67	118.47	113.75
5	A	301	2LJ	C2'-C1'-N8	2.43	118.27	111.53
5	C	301	2LJ	C2'-C1'-N8	2.12	117.40	111.53

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	301	2LJ	N8-C1'-C2'-C3'
5	C	301	2LJ	N8-C1'-C2'-C3'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/271 (94%)	-0.42	2 (0%) 82 80	10, 25, 57, 75	1 (0%)
1	C	263/271 (97%)	-0.54	4 (1%) 72 68	10, 22, 48, 61	0
2	B	96/99 (96%)	0.13	3 (3%) 51 47	18, 43, 66, 85	0
2	F	97/99 (97%)	-0.42	1 (1%) 79 76	12, 28, 55, 64	0
3	D	185/203 (91%)	0.03	14 (7%) 20 17	10, 33, 76, 89	0
3	G	198/203 (97%)	-0.46	1 (0%) 87 85	9, 24, 51, 63	0
4	E	240/245 (97%)	-0.26	4 (1%) 69 65	10, 29, 67, 87	1 (0%)
4	H	243/245 (99%)	-0.53	2 (0%) 82 80	11, 23, 42, 67	2 (0%)
All	All	1577/1636 (96%)	-0.35	31 (1%) 65 61	9, 25, 62, 89	4 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	176	ASN	6.4
1	C	16	ILE	4.6
1	A	223	GLN	3.4
4	E	242	GLY	3.4
2	B	96	ASP	3.3
3	D	184	ASN	3.2
3	G	199	SER	3.1
3	D	122	ARG	3.0
1	C	222	VAL	3.0
4	H	1	ASN	2.8
4	H	185	ASN	2.8
3	D	190	ILE	2.8
2	B	1	ILE	2.8
1	C	246	LEU	2.7
3	D	198	PRO	2.7
1	A	221	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	E	227	ASP	2.6
2	F	97	ARG	2.5
4	E	225	THR	2.3
3	D	146	SER	2.3
3	D	181	ALA	2.3
3	D	150	ASP	2.3
1	C	252	ASN	2.2
3	D	188	ASN	2.2
3	D	191	ILE	2.2
3	D	187	ASN	2.2
4	E	221	ASN	2.2
3	D	195	THR	2.1
3	D	189	SER	2.1
3	D	182	CYS	2.1
2	B	92	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NA	H	301	1/1	0.91	0.06	40,40,40,40	0
5	2LJ	C	301	22/22	0.98	0.05	11,13,17,18	0
5	2LJ	A	301	22/22	0.98	0.04	12,14,16,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2LJ C 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 2LJ A 301:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.