



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:42 AM UTC

PDB ID : 4NQI / pdb_00004nqi
Title : Structure of the N-terminal I-BAR domain (1-259) of D.Discoideum IBARa
Authors : Witte, G.; Faix, J.; Runge-Wollmann, P.
Deposited on : 2013-11-25
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	
1	B	264	
1	C	264	
1	D	264	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SH3 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	Se	83	0	0
			1767	1102	315	345	5			
1	B	223	Total	C	N	O	Se	140	0	0
			1790	1118	313	354	5			
1	C	227	Total	C	N	O	Se	64	0	0
			1828	1140	323	360	5			
1	D	232	Total	C	N	O	Se	61	1	0
			1869	1165	333	366	5			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP C7FZZ0
A	-3	PRO	-	expression tag	UNP C7FZZ0
A	-2	LEU	-	expression tag	UNP C7FZZ0
A	-1	GLY	-	expression tag	UNP C7FZZ0
A	0	SER	-	expression tag	UNP C7FZZ0
A	1	MSE	-	expression tag	UNP C7FZZ0
A	11	VAL	GLY	SEE REMARK 999	UNP C7FZZ0
B	-4	GLY	-	expression tag	UNP C7FZZ0
B	-3	PRO	-	expression tag	UNP C7FZZ0
B	-2	LEU	-	expression tag	UNP C7FZZ0
B	-1	GLY	-	expression tag	UNP C7FZZ0
B	0	SER	-	expression tag	UNP C7FZZ0
B	1	MSE	-	expression tag	UNP C7FZZ0
B	11	VAL	GLY	SEE REMARK 999	UNP C7FZZ0
C	-4	GLY	-	expression tag	UNP C7FZZ0
C	-3	PRO	-	expression tag	UNP C7FZZ0
C	-2	LEU	-	expression tag	UNP C7FZZ0
C	-1	GLY	-	expression tag	UNP C7FZZ0
C	0	SER	-	expression tag	UNP C7FZZ0
C	1	MSE	-	expression tag	UNP C7FZZ0
C	11	VAL	GLY	SEE REMARK 999	UNP C7FZZ0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-4	GLY	-	expression tag	UNP C7FZZ0
D	-3	PRO	-	expression tag	UNP C7FZZ0
D	-2	LEU	-	expression tag	UNP C7FZZ0
D	-1	GLY	-	expression tag	UNP C7FZZ0
D	0	SER	-	expression tag	UNP C7FZZ0
D	1	MSE	-	expression tag	UNP C7FZZ0
D	11	VAL	GLY	SEE REMARK 999	UNP C7FZZ0

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total	O	0	0
			40	40		
4	B	20	Total	O	0	0
			20	20		
4	C	117	Total	O	0	0
			117	117		
4	D	99	Total	O	0	0
			99	99		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SH3 domain-containing protein




- Molecule 1: SH3 domain-containing protein




- Molecule 1: SH3 domain-containing protein



- Molecule 1: SH3 domain-containing protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.10Å 60.21Å 147.62Å 90.00° 100.38° 90.00°	Depositor
Resolution (Å)	48.40 – 2.21 48.40 – 2.21	Depositor EDS
% Data completeness (in resolution range)	98.5 (48.40-2.21) 99.4 (48.40-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.194 , 0.230 0.197 , 0.232	Depositor DCC
R_{free} test set	2562 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.319	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 61.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7559	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1779	0.80	2/2377 (0.1%)
1	B	0.33	0/1803	0.78	3/2415 (0.1%)
1	C	0.34	0/1841	0.71	0/2464
1	D	0.34	0/1886	0.74	2/2524 (0.1%)
All	All	0.34	0/7309	0.76	7/9780 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	137	LYS	N-CA-C	9.57	121.66	111.03
1	A	155	THR	N-CA-C	-6.86	103.92	112.90
1	B	9	ASN	CA-C-N	6.46	125.69	118.97
1	B	9	ASN	C-N-CA	6.46	125.69	118.97
1	B	137	LYS	N-CA-C	-5.94	105.89	113.02
1	D	9	ASN	CA-C-N	5.02	124.19	118.97
1	D	9	ASN	C-N-CA	5.02	124.19	118.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1767	0	1811	18	0
1	B	1790	0	1827	18	0
1	C	1828	0	1869	7	0
1	D	1869	0	1922	11	0
2	B	14	0	20	5	0
2	C	7	0	10	0	0
3	D	8	0	6	1	0
4	A	40	0	0	0	0
4	B	20	0	0	1	0
4	C	117	0	0	3	0
4	D	99	0	0	5	0
All	All	7559	0	7465	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:108:ASP:OD2	4:C:485:HOH:O	2.01	0.78
3:D:301:ACT:O	4:D:491:HOH:O	2.02	0.77
1:D:207:LYS:NZ	4:D:471:HOH:O	1.82	0.73
1:D:41:GLN:HB3	1:D:94:LEU:HD23	1.73	0.70
1:D:49:GLN:NE2	4:D:435:HOH:O	2.06	0.69
1:B:41:GLN:OE1	1:B:45:GLN:NE2	2.26	0.69
1:A:131:ILE:HD11	1:A:161:ILE:HG13	1.75	0.68
1:D:43:GLN:NE2	4:D:401:HOH:O	2.25	0.67
1:A:93:VAL:HG21	1:A:196:LYS:HG3	1.79	0.64
1:A:225:LEU:HD22	1:B:173:ARG:HG2	1.80	0.64
1:C:216:ARG:NH1	4:C:432:HOH:O	2.19	0.64
1:A:21:ASN:ND2	1:A:179:GLU:OE2	2.33	0.62
1:B:185:THR:HB	2:B:302:PEG:H22	1.80	0.61
1:B:51:GLY:HA3	1:B:84:GLU:HG3	1.81	0.61
1:B:226:PRO:O	1:B:229:MSE:HB2	2.01	0.60
1:D:93:VAL:HG21	1:D:196:LYS:HG3	1.83	0.60
1:B:182:LYS:HA	2:B:302:PEG:H21	1.84	0.59
1:C:146:THR:HG23	1:C:149:VAL:H	1.67	0.58
1:A:128:ARG:HA	1:A:131:ILE:HG22	1.86	0.57
1:A:138:THR:HA	1:A:141:ALA:HB3	1.87	0.57
1:A:173:ARG:HG2	1:B:225:LEU:HD11	1.88	0.55
1:D:197:GLU:OE2	4:D:493:HOH:O	2.18	0.54
1:A:104:GLN:NE2	1:A:108:ASP:OD1	2.37	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:ASP:OD1	1:A:68:ASP:N	2.41	0.52
1:A:34:ILE:HD11	1:A:102:ASN:HB3	1.92	0.52
1:A:88:ASP:O	1:A:92:LYS:HG2	2.11	0.51
1:B:103:LYS:NZ	4:B:404:HOH:O	2.40	0.51
1:B:226:PRO:HD2	1:B:229:MSE:HG3	1.93	0.50
1:B:152:GLN:HA	1:B:155:THR:HG22	1.92	0.49
1:C:43:GLN:NE2	4:C:439:HOH:O	2.35	0.47
1:D:51:GLY:HA3	1:D:84:GLU:HG3	1.97	0.47
1:C:51:GLY:HA3	1:C:84:GLU:HG3	1.96	0.46
1:D:14:ILE:HG22	1:D:172:LEU:HD11	1.97	0.46
1:A:87:ARG:HH12	2:B:301:PEG:H42	1.80	0.46
1:B:93:VAL:HG13	1:B:97:GLU:HG3	1.96	0.46
1:B:146:THR:O	1:B:149:VAL:N	2.49	0.45
1:A:68:ASP:HB2	1:A:218:LEU:HD21	1.98	0.45
1:A:156:GLU:O	1:A:160:LYS:HB2	2.17	0.45
1:D:134:LEU:O	1:D:138:THR:HG23	2.17	0.44
1:C:68:ASP:OD1	1:C:68:ASP:N	2.50	0.44
1:A:90:VAL:HG11	1:A:197:GLU:OE2	2.18	0.44
1:D:134:LEU:HD23	1:D:134:LEU:HA	1.88	0.43
1:B:93:VAL:HG21	1:B:196:LYS:HG3	2.00	0.43
1:A:51:GLY:HA3	1:A:84:GLU:HG3	1.99	0.43
1:A:93:VAL:O	1:A:97:GLU:HB2	2.18	0.43
1:B:82:ASP:O	1:B:86:ARG:HG3	2.19	0.43
1:B:123:ASP:HB3	1:B:167:LEU:HD21	2.00	0.42
1:D:140:LYS:C	1:D:142:GLY:H	2.28	0.41
1:B:87:ARG:NH1	2:B:301:PEG:H21	2.35	0.41
1:A:15:THR:O	1:A:19:ILE:HG13	2.21	0.41
1:B:185:THR:CB	2:B:302:PEG:H22	2.50	0.41
1:B:229:MSE:HE2	1:B:229:MSE:HA	2.03	0.41
1:C:93:VAL:HG21	1:C:196:LYS:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/264 (82%)	211 (98%)	5 (2%)	0	100	100
1	B	219/264 (83%)	211 (96%)	8 (4%)	0	100	100
1	C	223/264 (84%)	222 (100%)	1 (0%)	0	100	100
1	D	231/264 (88%)	226 (98%)	5 (2%)	0	100	100
All	All	889/1056 (84%)	870 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/228 (86%)	187 (95%)	10 (5%)	21	26
1	B	202/228 (89%)	190 (94%)	12 (6%)	18	20
1	C	206/228 (90%)	201 (98%)	5 (2%)	43	56
1	D	210/228 (92%)	203 (97%)	7 (3%)	33	44
All	All	815/912 (89%)	781 (96%)	34 (4%)	26	34

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	GLU
1	A	56	ASP
1	A	134	LEU
1	A	137	LYS
1	A	138	THR
1	A	155	THR
1	A	156	GLU
1	A	158	ASN
1	A	161	ILE
1	A	231	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	13	GLU
1	B	118	LYS
1	B	128	ARG
1	B	131	ILE
1	B	132	LEU
1	B	138	THR
1	B	145	THR
1	B	154	ILE
1	B	165	GLU
1	B	166	GLN
1	B	227	LEU
1	B	230	GLU
1	C	42	GLN
1	C	134	LEU
1	C	137	LYS
1	C	138	THR
1	C	206	SER
1	D	8	GLN
1	D	11	VAL
1	D	118	LYS
1	D	138	THR
1	D	145	THR
1	D	225	LEU
1	D	227	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	38	ASN
1	A	45	GLN
1	A	183	HIS
1	B	64	HIS
1	B	104	GLN
1	B	220	ASN
1	C	102	ASN
1	C	211	ASN
1	D	43	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	C	301	-	6,6,6	0.61	0	5,5,5	1.43	0
2	PEG	B	302	-	6,6,6	0.62	0	5,5,5	1.46	1 (20%)
3	ACT	D	301	-	3,3,3	0.82	0	3,3,3	1.33	0
2	PEG	B	301	-	6,6,6	0.62	0	5,5,5	1.47	0
3	ACT	D	302	-	3,3,3	0.81	0	3,3,3	1.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	301	-	-	3/4/4/4	-
2	PEG	C	301	-	-	3/4/4/4	-
2	PEG	B	302	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	302	PEG	O2-C2-C1	2.06	119.17	110.11

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	301	PEG	O1-C1-C2-O2
2	B	301	PEG	O2-C3-C4-O4
2	B	302	PEG	O1-C1-C2-O2
2	B	302	PEG	C4-C3-O2-C2
2	B	301	PEG	O1-C1-C2-O2
2	C	301	PEG	C1-C2-O2-C3
2	B	301	PEG	C4-C3-O2-C2
2	C	301	PEG	C4-C3-O2-C2

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	PEG	3	0
3	D	301	ACT	1	0
2	B	301	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/264 (81%)	0.52	19 (8%) 15 13	24, 55, 87, 115	28 (13%)
1	B	218/264 (82%)	0.50	14 (6%) 25 22	28, 54, 88, 117	40 (18%)
1	C	222/264 (84%)	-0.14	7 (3%) 50 47	23, 36, 74, 99	21 (9%)
1	D	227/264 (85%)	0.09	9 (3%) 42 39	19, 41, 69, 113	22 (9%)
All	All	882/1056 (83%)	0.24	49 (5%) 30 27	19, 47, 79, 117	111 (12%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	138	THR	7.8
1	A	154	ILE	5.3
1	D	142	GLY	3.9
1	D	141	ALA	3.5
1	D	239	THR	3.5
1	A	141	ALA	3.2
1	A	137	LYS	3.1
1	A	138	THR	3.1
1	C	139	ARG	3.0
1	B	130	ASP	2.9
1	D	147	PRO	2.9
1	D	144	LYS	2.8
1	A	134	LEU	2.7
1	A	140	LYS	2.6
1	D	146	THR	2.6
1	A	11	VAL	2.6
1	A	133	LYS	2.6
1	D	237	GLU	2.6
1	B	145	THR	2.6
1	D	145	THR	2.6
1	B	149	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	225	LEU	2.5
1	A	126	GLN	2.5
1	A	160	LYS	2.5
1	B	155	THR	2.4
1	B	134	LEU	2.4
1	A	136	ALA	2.4
1	A	155	THR	2.4
1	A	132	LEU	2.4
1	C	145	THR	2.4
1	B	146	THR	2.3
1	C	149	VAL	2.3
1	B	230	GLU	2.3
1	B	233	ILE	2.3
1	B	227	LEU	2.3
1	A	10	PRO	2.3
1	B	147	PRO	2.3
1	A	166	GLN	2.2
1	B	133	LYS	2.2
1	A	142	GLY	2.2
1	C	146	THR	2.2
1	A	164	SER	2.1
1	D	138	THR	2.1
1	A	167	LEU	2.1
1	C	134	LEU	2.1
1	C	150	LEU	2.1
1	C	7	GLN	2.0
1	A	123	ASP	2.0
1	B	226	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	B	302	7/7	0.74	0.22	56,76,82,83	0
2	PEG	B	301	7/7	0.83	0.16	65,71,75,77	0
2	PEG	C	301	7/7	0.85	0.16	59,72,77,78	0
3	ACT	D	302	4/4	0.85	0.20	51,57,62,63	0
3	ACT	D	301	4/4	0.87	0.15	50,51,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.