



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:03 AM UTC

PDB ID : 3NR0 / pdb_00003nr0
Title : Optimization of the in silico designed Kemp eliminase KE70 by computational design and directed evolution R6 6/10A
Authors : Khersonsky, O.; Rothlisberge, D.; Wollacott, A.M.; Dym, O.; Baker, D.; Tawfik, D.S.; Israel Structural Proteomics Center (ISPC)
Deposited on : 2010-06-30
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
1	B	271	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called deoxyribose phosphate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	0	0
			1796	1138	305	344	9			
1	B	245	Total	C	N	O	S	0	0	0
			1814	1148	311	346	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	29	Total	O	0	0
			29	29		
2	B	37	Total	O	0	0
			37	37		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: deoxyribose phosphate aldolase



- Molecule 1: deoxyribose phosphate aldolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.10Å 41.22Å 139.44Å 90.00° 94.96° 90.00°	Depositor
Resolution (Å)	50.00 – 2.19 50.00 – 2.19	Depositor EDS
% Data completeness (in resolution range)	95.0 (50.00-2.19) 95.0 (50.00-2.19)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.4.0067	Depositor
R, R_{free}	0.207 , 0.268 0.206 , 0.265	Depositor DCC
R_{free} test set	1416 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 26.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3676	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.48	11/1822 (0.6%)	1.34	10/2474 (0.4%)
1	B	1.54	11/1840 (0.6%)	1.41	11/2496 (0.4%)
All	All	1.51	22/3662 (0.6%)	1.37	21/4970 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	53	ILE	CA-CB	10.54	1.60	1.54
1	B	3	THR	CA-CB	7.73	1.65	1.53
1	A	3	THR	CA-CB	6.88	1.64	1.53
1	B	243	ALA	CA-CB	6.71	1.64	1.53
1	B	209	ALA	CA-CB	6.22	1.63	1.53
1	A	101	VAL	CA-CB	6.13	1.62	1.54
1	B	228	ALA	C-O	-6.08	1.17	1.23
1	B	102	ALA	CA-CB	6.03	1.61	1.53
1	A	104	VAL	CA-CB	6.00	1.61	1.54
1	B	229	ASP	CA-C	5.95	1.61	1.53
1	A	118	GLY	N-CA	5.92	1.53	1.45
1	B	161	ALA	C-O	-5.65	1.16	1.24
1	A	181	SER	N-CA	5.63	1.53	1.46
1	B	198	VAL	C-O	5.46	1.29	1.24
1	B	178	THR	CA-CB	5.44	1.60	1.54
1	A	72	THR	CA-CB	5.34	1.62	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	GLU	C-O	-5.33	1.17	1.24
1	A	229	ASP	CA-C	5.33	1.60	1.53
1	A	109	ALA	CA-CB	-5.24	1.45	1.53
1	A	235	PHE	CA-C	-5.12	1.46	1.52
1	A	76	PHE	C-O	-5.06	1.17	1.24
1	B	107	TYR	CA-C	5.02	1.59	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	229	ASP	N-CA-C	-11.46	93.19	109.69
1	B	228	ALA	N-CA-C	10.98	126.85	109.39
1	B	229	ASP	N-CA-C	-10.52	94.80	109.95
1	A	228	ALA	N-CA-C	8.88	123.52	109.39
1	B	147	ASP	N-CA-C	8.20	122.77	110.64
1	A	147	ASP	N-CA-C	6.49	119.78	110.24
1	A	147	ASP	CB-CA-C	-6.28	100.42	109.84
1	A	112	ALA	N-CA-C	-6.26	105.22	112.92
1	B	147	ASP	CB-CA-C	-6.07	102.34	109.80
1	B	43	THR	N-CA-C	-5.88	102.73	110.43
1	A	204	GLY	N-CA-C	-5.76	104.28	112.81
1	B	107	TYR	N-CA-C	5.70	118.43	111.82
1	A	35	GLN	CB-CG-CD	-5.63	103.03	112.60
1	B	78	HIS	N-CA-C	5.55	118.26	111.82
1	A	78	HIS	N-CA-C	5.34	118.02	111.82
1	B	15	MET	N-CA-C	5.18	117.83	110.10
1	B	180	GLU	N-CA-C	5.13	116.68	111.14
1	A	43	THR	N-CA-C	-5.10	103.75	110.43
1	A	222	LEU	N-CA-C	5.06	116.88	111.36
1	B	240	SER	N-CA-CB	5.03	117.61	110.16
1	B	225	ALA	N-CA-C	5.02	118.50	112.38

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18	ALA	Peptide
1	A	2	ALA	Peptide
1	A	228	ALA	Peptide
1	B	18	ALA	Peptide
1	B	228	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1796	0	1796	19	0
1	B	1814	0	1835	19	0
2	A	29	0	0	3	0
2	B	37	0	0	1	0
All	All	3676	0	3631	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:LYS:HE2	2:A:329:HOH:O	1.75	0.84
1:A:214:LYS:CE	2:A:329:HOH:O	2.36	0.71
1:B:202:PRO:HD2	1:B:234:ALA:O	1.93	0.68
1:A:196:LYS:HG3	2:A:321:HOH:O	1.93	0.68
1:B:10:ARG:HD3	1:B:216:LEU:HD13	1.75	0.68
1:A:202:PRO:HD2	1:A:234:ALA:O	1.96	0.65
1:B:15:MET:HE3	1:B:237:ALA:HB3	1.83	0.60
1:A:179:PRO:HG3	1:A:214:LYS:HE3	1.83	0.60
1:B:247:LYS:HD2	1:B:247:LYS:C	2.29	0.57
1:B:107:TYR:CE1	1:B:108:ARG:CD	2.88	0.56
1:A:85:ILE:O	1:A:89:GLU:HG3	2.06	0.55
1:A:179:PRO:CG	1:A:214:LYS:HE3	2.37	0.55
1:B:55:ILE:HD12	1:B:55:ILE:O	2.08	0.54
1:A:221:GLU:HG2	1:A:221:GLU:O	2.03	0.54
1:B:85:ILE:O	1:B:89:GLU:HG3	2.09	0.52
1:A:195:GLU:HB2	1:A:227:TRP:NE1	2.24	0.52
1:A:10:ARG:HD3	1:A:216:LEU:HD13	1.92	0.50
1:B:100:SER:HA	1:B:134:VAL:HG13	1.93	0.49
1:B:108:ARG:HD3	2:B:302:HOH:O	2.13	0.48
1:A:179:PRO:HG3	1:A:214:LYS:CE	2.42	0.48
1:B:207:ARG:HE	1:B:207:ARG:HB2	1.40	0.47
1:B:129:CYS:HB3	1:B:134:VAL:O	2.15	0.46
1:A:2:ALA:HA	1:A:5:LYS:HB2	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:LEU:HD21	1:B:210:GLU:OE2	2.17	0.45
1:B:107:TYR:CE1	1:B:108:ARG:NE	2.86	0.44
1:A:195:GLU:HB2	1:A:227:TRP:HE1	1.83	0.43
1:A:167:VAL:HG13	1:A:201:ILE:HB	1.99	0.43
1:A:172:LYS:HD2	1:A:172:LYS:HA	1.78	0.43
1:B:57:ARG:NH2	1:B:97:GLY:O	2.51	0.43
1:B:216:LEU:HA	1:B:216:LEU:HD23	1.79	0.42
1:B:126:LYS:HE2	1:B:130:ALA:HB2	2.01	0.42
1:A:106:PRO:CG	1:A:118:GLY:HA2	2.49	0.41
1:A:100:SER:HA	1:A:134:VAL:HG13	2.03	0.41
1:B:107:TYR:O	1:B:111:MET:HG3	2.20	0.41
1:A:107:TYR:O	1:A:111:MET:HG3	2.20	0.41
1:B:172:LYS:HD2	1:B:172:LYS:HA	1.71	0.40
1:B:13:LYS:O	1:B:44:ASP:HB2	2.21	0.40
1:A:203:VAL:HG22	1:A:236:GLY:HA3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/271 (89%)	234 (97%)	7 (3%)	0	100	100
1	B	241/271 (89%)	233 (97%)	8 (3%)	0	100	100
All	All	482/542 (89%)	467 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/207 (86%)	168 (95%)	9 (5%)	21	27
1	B	183/207 (88%)	171 (93%)	12 (7%)	15	18
All	All	360/414 (87%)	339 (94%)	21 (6%)	18	22

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	19	THR
1	A	55	ILE
1	A	108	ARG
1	A	147	ASP
1	A	214	LYS
1	A	221	GLU
1	A	238	SER
1	A	240	SER
1	B	3	THR
1	B	19	THR
1	B	55	ILE
1	B	147	ASP
1	B	153	LYS
1	B	156	GLU
1	B	172	LYS
1	B	181	SER
1	B	207	ARG
1	B	238	SER
1	B	240	SER
1	B	247	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	HIS
1	B	16	HIS
1	B	251	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/271 (90%)	0.09	6 (2%) 59 56	19, 31, 46, 62	0
1	B	245/271 (90%)	-0.10	4 (1%) 70 67	20, 29, 42, 54	0
All	All	490/542 (90%)	-0.01	10 (2%) 65 62	19, 30, 44, 62	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	4.2
1	A	4	LEU	3.8
1	B	2	ALA	3.8
1	A	251	HIS	3.6
1	A	3	THR	3.3
1	A	19	THR	3.1
1	B	19	THR	2.7
1	A	204	GLY	2.3
1	B	250	GLY	2.3
1	B	108	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.