



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:49 AM UTC

PDB ID : 1NSZ / pdb_00001nsz
Title : Crystal structure of galactose mutarotase from *Lactococcus lactis* mutant H170N complexed with glucose
Authors : Holden, H.M.; Thoden, J.B.
Deposited on : 2003-01-28
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	 80% 16% . .
1	B	347	 83% 14% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5822 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GALACTOSE MUTAROTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	0	1	0
			2656	1674	447	532	3			
1	B	346	Total	C	N	O	S	0	1	0
			2720	1712	465	540	3			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	SER	GLU	cloning artifact	UNP Q9ZB17
A	170	ASN	HIS	engineered mutation	UNP Q9ZB17
A	340	LEU	-	expression tag	UNP Q9ZB17
A	341	GLU	-	expression tag	UNP Q9ZB17
A	342	HIS	-	expression tag	UNP Q9ZB17
A	343	HIS	-	expression tag	UNP Q9ZB17
A	344	HIS	-	expression tag	UNP Q9ZB17
A	345	HIS	-	expression tag	UNP Q9ZB17
A	346	HIS	-	expression tag	UNP Q9ZB17
A	347	HIS	-	expression tag	UNP Q9ZB17
B	2	SER	GLU	cloning artifact	UNP Q9ZB17
B	170	ASN	HIS	engineered mutation	UNP Q9ZB17
B	340	LEU	-	expression tag	UNP Q9ZB17
B	341	GLU	-	expression tag	UNP Q9ZB17
B	342	HIS	-	expression tag	UNP Q9ZB17
B	343	HIS	-	expression tag	UNP Q9ZB17
B	344	HIS	-	expression tag	UNP Q9ZB17
B	345	HIS	-	expression tag	UNP Q9ZB17
B	346	HIS	-	expression tag	UNP Q9ZB17
B	347	HIS	-	expression tag	UNP Q9ZB17

- Molecule 2 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	196	Total	O	0	0
			196	196		
4	B	225	Total	O	0	0
			225	225		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GALACTOSE MUTAROTASE



• Molecule 1: GALACTOSE MUTAROTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.90Å 76.50Å 210.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.75 30.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.1 (30.00-1.75) 97.9 (30.00-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 1.72Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.165 , 0.218 0.160 , 0.165	Depositor DCC
R_{free} test set	7456 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 75.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5822	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	0/2711	1.51	18/3669 (0.5%)
1	B	1.13	1/2781 (0.0%)	1.44	19/3763 (0.5%)
All	All	1.12	1/5492 (0.0%)	1.48	37/7432 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	309	PRO	CA-C	5.17	1.58	1.52

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340	LEU	N-CA-CB	7.50	121.59	110.49
1	A	227	ASN	CA-CB-CG	-7.35	105.25	112.60
1	A	33	ARG	N-CA-CB	-6.99	99.13	111.08
1	A	210	VAL	N-CA-C	6.59	117.40	108.17
1	A	244	HIS	N-CA-C	6.58	115.88	108.25
1	A	76	LEU	N-CA-C	6.54	119.91	109.24
1	A	307	VAL	N-CA-C	-6.41	100.49	109.21
1	A	33	ARG	N-CA-C	6.41	119.98	109.72
1	A	244	HIS	CB-CA-C	-6.38	100.08	109.97
1	A	38	GLN	N-CA-C	6.36	119.09	109.23
1	A	340	LEU	O-C-N	6.32	130.64	122.43
1	B	33	ARG	N-CA-CB	-6.24	100.41	111.08
1	B	307	VAL	N-CA-C	-6.04	101.00	109.21
1	A	63	ALA	N-CA-C	6.00	118.64	110.55
1	B	63	ALA	N-CA-C	5.95	118.59	110.55
1	B	246	PHE	CA-CB-CG	-5.90	107.90	113.80
1	B	244	HIS	O-C-N	5.83	125.44	121.71
1	B	189	ARG	CB-CA-C	5.81	120.35	109.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	337	HIS	CA-CB-CG	-5.79	108.01	113.80
1	A	125	VAL	N-CA-CB	5.79	118.75	111.46
1	A	244	HIS	O-C-N	5.69	125.80	121.65
1	B	210	VAL	N-CA-CB	-5.54	101.75	111.39
1	B	43	HIS	CB-CA-C	-5.54	101.03	109.89
1	B	154	HIS	CA-CB-CG	-5.53	108.27	113.80
1	A	340	LEU	CA-C-N	5.47	131.54	121.70
1	A	340	LEU	C-N-CA	5.47	131.54	121.70
1	B	232	ASN	N-CA-CB	5.39	119.55	110.92
1	B	33	ARG	N-CA-C	5.37	118.30	109.72
1	B	347	HIS	CA-CB-CG	-5.32	108.48	113.80
1	A	223	LYS	CB-CA-C	5.28	119.81	110.16
1	B	76	LEU	N-CA-C	5.20	117.20	108.73
1	B	211	ASP	CA-CB-CG	-5.18	107.42	112.60
1	A	194	ARG	NE-CZ-NH1	5.09	126.59	121.50
1	B	186	HIS	CA-CB-CG	5.06	118.86	113.80
1	B	244	HIS	N-CA-C	5.05	116.43	108.55
1	B	345	HIS	CA-CB-CG	-5.04	108.76	113.80
1	B	326	GLU	CB-CA-C	5.01	117.99	109.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2656	0	2592	33	0
1	B	2720	0	2635	33	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
3	A	1	0	0	0	0
4	A	196	0	0	3	0
4	B	225	0	0	2	0
All	All	5822	0	5251	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:294:LYS:HB2	1:B:294:LYS:NZ	1.99	0.76
1:B:78:LYS:NZ	1:B:78:LYS:HB3	2.01	0.75
1:B:189:ARG:HG3	1:B:189:ARG:HH11	1.53	0.74
1:B:218:ASP:O	1:B:223:LYS:HE2	1.89	0.72
1:B:338:THR:HG22	1:B:339:LYS:HG3	1.70	0.71
1:A:201:GLN:NE2	1:A:201:GLN:H	1.88	0.71
1:A:218:ASP:OD1	1:A:220:ARG:HD3	1.92	0.69
1:B:189:ARG:HG3	1:B:189:ARG:NH1	2.05	0.69
1:B:338:THR:CG2	1:B:339:LYS:HG3	2.31	0.60
1:B:78:LYS:HG3	1:B:83:ASP:OD1	2.02	0.59
1:B:294:LYS:HB2	1:B:294:LYS:HZ2	1.65	0.59
1:A:76:LEU:HD23	1:A:77:VAL:N	2.18	0.59
1:A:201:GLN:H	1:A:201:GLN:CD	2.11	0.57
1:B:78:LYS:HB3	1:B:78:LYS:HZ3	1.69	0.56
1:A:20:LYS:HD2	1:A:147:ASP:OD2	2.07	0.55
1:A:39:LYS:HG3	4:A:1499:HOH:O	2.09	0.53
1:B:337:HIS:HD2	4:B:2613:HOH:O	1.92	0.53
1:B:78:LYS:HB3	1:B:78:LYS:HZ2	1.72	0.53
1:A:218:ASP:OD2	1:A:220:ARG:NH1	2.43	0.51
1:A:82:LYS:HD3	1:A:84:TYR:CE1	2.46	0.51
1:A:152:LYS:HG3	1:A:333:ILE:HG12	1.94	0.49
1:B:52:LYS:HE2	1:B:56:GLU:OE2	2.13	0.49
1:B:323:LYS:O	1:B:326:GLU:HG2	2.13	0.48
1:A:221:GLN:O	1:A:222:GLU:C	2.54	0.48
1:B:157:ALA:C	1:B:158:ILE:HG12	2.39	0.47
1:A:337:HIS:HB3	4:A:1482:HOH:O	2.14	0.47
1:A:280:THR:O	1:A:298:HIS:HA	2.15	0.47
1:A:127:ASN:HB2	1:B:130:THR:HG23	1.98	0.46
1:A:212:ILE:O	1:A:215:THR:HG23	2.14	0.46
1:A:289:LEU:HD23	1:A:294:LYS:HA	1.97	0.46
1:A:232:ASN:O	1:A:237:GLN:NE2	2.49	0.46
1:A:60:TYR:N	1:A:61:PRO:CD	2.80	0.45
1:A:291:HIS:O	1:A:292:GLU:HB2	2.16	0.45
1:B:60:TYR:N	1:B:61:PRO:CD	2.79	0.45
1:B:158:ILE:HD13	1:B:327:LYS:HA	1.99	0.45
1:B:189:ARG:HH11	1:B:189:ARG:CG	2.25	0.45
1:A:79:ILE:HB	1:A:84:TYR:CE1	2.52	0.45
1:A:253:LEU:HB2	4:A:1594:HOH:O	2.15	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:GLY:O	1:B:32:ALA:C	2.60	0.45
1:B:197:PRO:HD2	1:B:208:ASP:O	2.17	0.45
1:A:197:PRO:HD2	1:A:208:ASP:O	2.17	0.45
1:A:193:SER:O	1:A:212:ILE:HG13	2.17	0.44
1:A:180:SER:HA	1:A:296:VAL:HG13	2.00	0.44
1:A:297:HIS:O	1:A:298:HIS:HB2	2.18	0.44
1:A:43:HIS:O	1:A:177:GLY:HA2	2.18	0.44
1:B:335:SER:HB2	4:B:2498:HOH:O	2.18	0.43
1:B:43:HIS:O	1:B:177:GLY:HA2	2.17	0.43
1:A:239:VAL:O	1:A:240:LYS:HB2	2.19	0.42
1:B:340:LEU:HD12	1:B:340:LEU:HA	1.62	0.42
1:A:19:ASN:HB2	1:A:147:ASP:OD1	2.19	0.42
1:B:158:ILE:CD1	1:B:327:LYS:HA	2.49	0.42
1:B:219:PHE:CD2	1:B:223:LYS:HG2	2.54	0.42
1:B:317:LEU:HA	1:B:317:LEU:HD23	1.79	0.42
1:A:244:HIS:HA	1:A:245:PRO:HD3	1.82	0.41
1:B:280:THR:O	1:B:298:HIS:HA	2.19	0.41
1:A:339:LYS:O	1:A:339:LYS:HG3	2.20	0.41
1:A:270:PHE:HB2	1:A:333:ILE:HB	2.03	0.41
1:A:285:ASP:O	1:A:286:LEU:C	2.61	0.41
1:B:294:LYS:NZ	1:B:294:LYS:CB	2.79	0.41
1:A:29:ASN:HA	1:A:122:PHE:CE2	2.56	0.41
1:B:244:HIS:HA	1:B:245:PRO:HD3	1.88	0.40
1:A:277:VAL:HB	1:A:304:GLU:HB2	2.03	0.40
1:B:135:GLY:HA3	1:B:136:LYS:HA	1.88	0.40
1:B:152:LYS:HG2	1:B:153:ILE:N	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/347 (98%)	318 (94%)	21 (6%)	0	100	100
1	B	345/347 (99%)	326 (94%)	19 (6%)	0	100	100
All	All	684/694 (99%)	644 (94%)	40 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/300 (98%)	283 (96%)	11 (4%)	30	11
1	B	300/300 (100%)	285 (95%)	15 (5%)	22	5
All	All	594/600 (99%)	568 (96%)	26 (4%)	25	7

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LYS
1	A	39	LYS
1	A	76	LEU
1	A	170	ASN
1	A	195	PHE
1	A	201	GLN
1	A	212	ILE
1	A	221	GLN
1	A	223	LYS
1	A	224	GLN
1	A	339	LYS
1	B	4	LYS
1	B	20	LYS
1	B	78	LYS
1	B	152	LYS
1	B	170	ASN
1	B	189	ARG
1	B	195	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	201	GLN
1	B	212	ILE
1	B	213	LYS
1	B	255	LYS
1	B	294	LYS
1	B	323	LYS
1	B	327	LYS
1	B	338	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	201	GLN
1	A	221	GLN
1	A	227	ASN
1	A	295	GLN
1	B	38	GLN
1	B	201	GLN
1	B	227	ASN
1	B	250	GLN
1	B	306	GLN
1	B	344	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	A	1400	-	12,12,12	0.51	0	17,17,17	1.24	2 (11%)
2	GLC	B	2400	-	12,12,12	0.66	0	17,17,17	1.25	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	1400	-	-	2/2/22/22	0/1/1/1
2	GLC	B	2400	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2400	GLC	C3-C4-C5	-3.10	104.62	110.23
2	B	2400	GLC	C4-C3-C2	-2.65	106.18	110.83
2	A	1400	GLC	O4-C4-C3	2.20	115.57	110.38
2	A	1400	GLC	C3-C4-C5	-2.19	106.27	110.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1400	GLC	C4-C5-C6-O6
2	A	1400	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/347 (97%)	-0.44	1 (0%) 90 92	15, 25, 56, 95	1 (0%)
1	B	346/347 (99%)	-0.60	0 100 100	15, 23, 49, 66	1 (0%)
All	All	686/694 (98%)	-0.52	1 (0%) 92 93	15, 24, 54, 95	2 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	340	LEU	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NA	A	1401	1/1	0.86	0.13	41,41,41,41	0
2	GLC	B	2400	12/12	0.96	0.06	25,32,36,43	0
2	GLC	A	1400	12/12	0.97	0.05	20,26,32,40	0

6.5 Other polymers ⓘ

There are no such residues in this entry.