



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:57 AM UTC

PDB ID : 2NS8 / pdb_00002ns8
Title : How an in vitro selected peptide mimics the antibiotic tetracycline to induce TET repressor
Authors : Luckner, S.R.; Klotzsche, M.; Berens, C.; Hillen, W.; Muller, Y.A.
Deposited on : 2006-11-03
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	7% 75% 18% . .
1	B	208	5% 71% 23% . .
1	C	208	7% 74% 22% . .
1	D	208	4% 73% 20% . 5%
2	E	17	6% 59% 12% 29%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	17	
2	G	17	
2	H	17	
2	Z	17	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetracycline repressor protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	200	Total	C	N	O	S	0	0	0
			1599	1009	285	303	2			
1	B	204	Total	C	N	O	S	0	0	0
			1626	1029	287	308	2			
1	C	206	Total	C	N	O	S	0	0	0
			1643	1038	292	311	2			
1	D	198	Total	C	N	O	S	0	0	0
			1577	996	280	299	2			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	68	SER	CYS	engineered mutation	UNP P04483
A	88	ASN	CYS	engineered mutation	UNP P04483
A	121	THR	CYS	engineered mutation	UNP P04483
A	144	SER	CYS	engineered mutation	UNP P04483
B	68	SER	CYS	engineered mutation	UNP P04483
B	88	ASN	CYS	engineered mutation	UNP P04483
B	121	THR	CYS	engineered mutation	UNP P04483
B	144	SER	CYS	engineered mutation	UNP P04483
C	68	SER	CYS	engineered mutation	UNP P04483
C	88	ASN	CYS	engineered mutation	UNP P04483
C	121	THR	CYS	engineered mutation	UNP P04483
C	144	SER	CYS	engineered mutation	UNP P04483
D	68	SER	CYS	engineered mutation	UNP P04483
D	88	ASN	CYS	engineered mutation	UNP P04483
D	121	THR	CYS	engineered mutation	UNP P04483
D	144	SER	CYS	engineered mutation	UNP P04483


- Molecule 2 is a protein called 16 residue peptide Tip (Transcription inducing peptide).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	13	Total	C	N	O	0	0	0
			102	70	15	17			
2	E	12	Total	C	N	O	0	0	0
			96	67	14	15			
2	F	13	Total	C	N	O	0	0	0
			102	70	15	17			
2	G	12	Total	C	N	O	0	0	0
			96	67	14	15			
2	Z	4	Total	C	N	O	0	0	0
			43	30	7	6			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tetracycline repressor protein



- Molecule 1: Tetracycline repressor protein




- Molecule 1: Tetracycline repressor protein




- Molecule 1: Tetracycline repressor protein






- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)




- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	77.89Å 200.59Å 65.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.55 30.00 – 2.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.00-2.55) 99.4 (30.00-2.55)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.206 , 0.270 0.214 , 0.274	Depositor DCC
R_{free} test set	2715 reflections (8.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.1	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6884	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/1631	0.94	2/2206 (0.1%)
1	B	0.87	1/1658 (0.1%)	1.00	3/2244 (0.1%)
1	C	0.78	0/1675	0.93	2/2266 (0.1%)
1	D	0.76	0/1607	0.91	1/2171 (0.0%)
2	E	0.76	0/100	0.82	0/140
2	F	0.67	0/106	0.62	0/148
2	G	0.61	0/100	0.67	0/140
2	H	0.81	0/106	0.83	0/148
2	Z	0.92	0/46	0.72	0/64
All	All	0.81	1/7029 (0.0%)	0.93	8/9527 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	162	THR	CA-CB	6.10	1.60	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	71	GLU	CB-CA-C	-7.86	106.44	115.79
1	C	160	THR	CA-C-N	6.36	127.80	119.84
1	C	160	THR	C-N-CA	6.36	127.80	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	159	GLU	N-CA-C	6.20	118.54	108.26
1	A	68	SER	N-CA-C	5.88	115.07	108.25
1	B	161	PRO	N-CA-C	5.55	119.91	111.19
1	A	45	VAL	CB-CA-C	-5.12	102.89	111.29
1	B	110	TYR	N-CA-C	5.04	117.42	111.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	164	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1599	0	1583	21	0
1	B	1626	0	1616	31	0
1	C	1643	0	1634	29	0
1	D	1577	0	1555	28	0
2	E	96	0	81	1	0
2	F	102	0	86	4	0
2	G	96	0	81	7	0
2	H	102	0	86	2	0
2	Z	43	0	35	0	0
All	All	6884	0	6757	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:0:ACE:H1	2:F:1:TRP:HB2	1.53	0.91
1:C:10:ILE:HG21	1:C:58:GLU:HG3	1.61	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:206:GLN:O	1:C:207:ILE:HG23	1.88	0.74
1:A:26:THR:OG1	1:A:29:LYS:HD2	1.90	0.71
1:C:200:GLN:HE21	1:D:188:HIS:HE1	1.39	0.70
1:B:119:PHE:CZ	1:B:123:GLN:NE2	2.62	0.68
1:B:70:LEU:O	1:B:73:GLU:HB3	1.93	0.67
1:B:4:LEU:HD13	1:B:4:LEU:O	1.95	0.66
1:C:132:TYR:OH	1:D:180:GLN:NE2	2.30	0.65
2:F:0:ACE:CH3	2:F:1:TRP:HB2	2.26	0.64
1:B:206:GLN:HE22	1:C:46:LYS:H	1.46	0.63
1:D:97:ALA:CB	1:D:146:LEU:HD13	2.29	0.62
1:A:163:THR:HG22	1:A:164:ASP:H	1.65	0.61
1:A:14:LEU:HD13	1:A:55:LEU:HG	1.83	0.61
1:B:119:PHE:CE1	1:B:123:GLN:NE2	2.69	0.61
1:A:26:THR:OG1	1:A:29:LYS:CD	2.47	0.61
1:D:10:ILE:HG21	1:D:58:GLU:HG3	1.82	0.61
1:B:106:THR:H	1:B:109:GLN:NE2	1.99	0.60
1:A:159:GLU:O	1:A:160:THR:O	2.20	0.60
1:A:59:MET:HE3	1:A:92:SER:OG	2.03	0.59
1:A:4:LEU:HD21	1:A:45:VAL:HG22	1.83	0.58
1:B:83:ALA:HB2	1:B:137:VAL:CG1	2.34	0.58
1:D:60:LEU:CD1	2:G:0:ACE:H3	2.35	0.57
1:A:146:LEU:HD13	1:B:146:LEU:HD13	1.87	0.56
1:D:66:HIS:CE1	1:D:70:LEU:HD11	2.40	0.56
1:B:14:LEU:HD13	1:B:55:LEU:HG	1.87	0.55
1:B:121:THR:HG22	1:B:125:PHE:O	2.06	0.55
1:C:28:ARG:NH1	1:C:38:GLN:HG3	2.22	0.54
1:A:59:MET:CE	1:A:89:ALA:O	2.56	0.54
1:D:152:GLN:NE2	1:D:152:GLN:HA	2.23	0.54
2:E:1:TRP:CD1	2:E:2:THR:H	2.26	0.54
1:C:177:PHE:CZ	1:D:131:LEU:HD22	2.43	0.54
1:D:60:LEU:HD11	2:G:0:ACE:H3	1.90	0.54
1:C:206:GLN:O	1:C:207:ILE:HG12	2.08	0.53
1:B:106:THR:OG1	1:B:109:GLN:NE2	2.39	0.53
1:A:171:ARG:O	1:A:175:GLU:OE2	2.26	0.53
1:D:151:HIS:CD2	2:F:7:ALA:HA	2.44	0.52
1:B:10:ILE:CG2	1:B:58:GLU:HG3	2.40	0.51
1:D:97:ALA:HB3	1:D:146:LEU:HD13	1.92	0.51
1:A:16:LEU:HD23	1:A:25:LEU:HD12	1.91	0.51
1:C:129:ASN:HD21	1:C:206:GLN:HE22	1.59	0.51
1:D:118:ALA:O	1:D:122:GLN:HG2	2.10	0.50
1:B:151:HIS:CD2	2:H:7:ALA:HA	2.46	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:MET:SD	1:B:174:ILE:HG13	2.51	0.50
1:C:150:GLU:HG2	1:D:101:LEU:HB3	1.93	0.50
1:D:97:ALA:HB1	1:D:146:LEU:HD13	1.93	0.50
1:D:148:ASP:O	1:D:152:GLN:HG2	2.12	0.50
1:B:59:MET:HE3	1:B:92:SER:OG	2.12	0.50
1:D:104:ARG:HD3	2:G:7:ALA:O	2.12	0.49
1:A:101:LEU:HB3	1:B:150:GLU:HG2	1.95	0.49
1:B:10:ILE:HG21	1:B:58:GLU:CG	2.42	0.49
1:D:159:GLU:CD	1:D:159:GLU:C	2.81	0.49
1:D:59:MET:HE2	1:D:89:ALA:O	2.13	0.49
1:A:26:THR:HG1	1:A:29:LYS:HD2	1.78	0.49
1:B:96:GLY:O	1:B:100:HIS:HD2	1.97	0.48
1:C:70:LEU:O	1:C:71:GLU:C	2.58	0.47
1:C:206:GLN:O	1:C:207:ILE:CG2	2.61	0.47
1:A:97:ALA:HB1	1:A:146:LEU:HG	1.96	0.47
1:C:90:LEU:HD22	1:C:97:ALA:HA	1.97	0.47
1:A:106:THR:OG1	1:A:109:GLN:NE2	2.48	0.47
1:B:36:VAL:HG13	1:B:40:THR:OG1	2.15	0.47
1:B:97:ALA:HB2	1:B:145:VAL:HG12	1.97	0.46
1:C:38:GLN:N	1:C:39:PRO:HD2	2.31	0.46
1:A:32:GLN:O	1:A:33:LYS:C	2.58	0.46
1:C:68:SER:HB2	1:C:69:PRO:HD2	1.98	0.46
1:B:10:ILE:HG21	1:B:58:GLU:HG3	1.98	0.46
1:B:13:ALA:HB3	1:B:55:LEU:HD11	1.98	0.46
1:C:110:TYR:CZ	1:D:166:MET:HE3	2.52	0.46
1:C:14:LEU:HD13	1:C:55:LEU:HG	1.97	0.45
1:B:97:ALA:HB1	1:B:146:LEU:HG	1.98	0.45
1:D:28:ARG:HD2	1:D:38:GLN:HE22	1.81	0.45
2:G:0:ACE:H1	2:G:1:TRP:HB2	1.98	0.45
1:C:6:LYS:O	1:C:10:ILE:HG12	2.17	0.45
1:B:177:PHE:CD2	2:H:5:ALA:HA	2.52	0.44
1:D:166:MET:HB3	1:D:167:PRO:CD	2.47	0.44
1:C:10:ILE:HG21	1:C:58:GLU:CG	2.42	0.43
1:B:16:LEU:O	1:B:20:VAL:HG22	2.18	0.43
1:D:10:ILE:HG21	1:D:58:GLU:CG	2.48	0.43
1:C:180:GLN:NE2	1:D:132:TYR:OH	2.51	0.43
1:A:166:MET:HB3	1:B:110:TYR:CE2	2.54	0.43
1:D:166:MET:CB	1:D:167:PRO:CD	2.97	0.43
1:B:47:ASN:OD1	1:B:47:ASN:C	2.62	0.43
1:A:58:GLU:OE1	1:A:62:ARG:NE	2.51	0.42
2:G:1:TRP:HZ2	2:G:7:ALA:CB	2.32	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:LEU:HD23	1:B:71:GLU:HB2	2.01	0.42
1:D:60:LEU:HD13	2:G:0:ACE:H3	2.01	0.42
1:C:42:TYR:OH	1:C:46:LYS:NZ	2.43	0.41
1:C:97:ALA:HB1	1:C:146:LEU:HG	2.02	0.41
2:F:1:TRP:HZ2	2:F:7:ALA:CB	2.33	0.41
1:C:160:THR:OG1	1:C:161:PRO:HD2	2.21	0.41
1:D:90:LEU:CD2	1:D:100:HIS:CD2	3.04	0.41
1:B:15:GLU:O	1:B:19:GLU:HG2	2.21	0.41
1:C:3:ARG:NH1	1:C:34:LEU:O	2.54	0.41
1:C:177:PHE:O	1:C:181:GLY:N	2.48	0.41
1:C:200:GLN:HE21	1:D:188:HIS:CE1	2.29	0.41
1:A:131:LEU:HD22	1:B:177:PHE:CZ	2.56	0.40
1:C:201:LEU:HD23	1:C:201:LEU:HA	1.92	0.40
1:A:163:THR:HA	1:A:166:MET:HE3	2.04	0.40
1:A:167:PRO:CB	1:A:168:PRO:CD	2.99	0.40
1:C:157:GLU:OE2	1:D:49:ARG:NH1	2.54	0.40
1:C:166:MET:HE1	2:G:8:PHE:CE2	2.56	0.40
1:B:14:LEU:HG	1:B:93:HIS:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/208 (95%)	183 (92%)	13 (7%)	2 (1%)	12	17
1	B	202/208 (97%)	197 (98%)	3 (2%)	2 (1%)	12	17
1	C	204/208 (98%)	197 (97%)	5 (2%)	2 (1%)	12	17
1	D	194/208 (93%)	181 (93%)	10 (5%)	3 (2%)	8	10
2	E	10/17 (59%)	9 (90%)	1 (10%)	0	100	100
2	F	11/17 (65%)	9 (82%)	2 (18%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	G	10/17 (59%)	9 (90%)	1 (10%)	0	100	100
2	H	11/17 (65%)	11 (100%)	0	0	100	100
2	Z	2/17 (12%)	2 (100%)	0	0	100	100
All	All	842/917 (92%)	798 (95%)	35 (4%)	9 (1%)	11	15

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	THR
1	B	165	SER
1	A	162	THR
1	C	64	HIS
1	C	161	PRO
1	D	71	GLU
1	D	158	ARG
1	D	166	MET
1	B	69	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/179 (96%)	153 (89%)	19 (11%)	6	6
1	B	175/179 (98%)	157 (90%)	18 (10%)	7	8
1	C	177/179 (99%)	154 (87%)	23 (13%)	4	4
1	D	168/179 (94%)	148 (88%)	20 (12%)	5	5
2	E	7/9 (78%)	7 (100%)	0	100	100
2	F	8/9 (89%)	8 (100%)	0	100	100
2	G	7/9 (78%)	7 (100%)	0	100	100
2	H	8/9 (89%)	8 (100%)	0	100	100
2	Z	4/9 (44%)	3 (75%)	1 (25%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	726/761 (95%)	645 (89%)	81 (11%)	6 6

All (81) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	14	LEU
1	A	23	GLU
1	A	28	ARG
1	A	30	LEU
1	A	32	GLN
1	A	45	VAL
1	A	46	LYS
1	A	55	LEU
1	A	68	SER
1	A	79	LEU
1	A	94	ARG
1	A	104	ARG
1	A	127	LEU
1	A	152	GLN
1	A	162	THR
1	A	176	LEU
1	A	199	VAL
1	A	202	THR
1	B	4	LEU
1	B	14	LEU
1	B	30	LEU
1	B	36	VAL
1	B	40	THR
1	B	45	VAL
1	B	55	LEU
1	B	67	PHE
1	B	68	SER
1	B	79	LEU
1	B	104	ARG
1	B	160	THR
1	B	162	THR
1	B	171	ARG
1	B	176	LEU
1	B	201	LEU
1	B	202	THR
1	B	204	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	14	LEU
1	C	28	ARG
1	C	30	LEU
1	C	32	GLN
1	C	70	LEU
1	C	79	LEU
1	C	94	ARG
1	C	106	THR
1	C	112	THR
1	C	122	GLN
1	C	152	GLN
1	C	153	VAL
1	C	156	GLU
1	C	159	GLU
1	C	160	THR
1	C	164	ASP
1	C	174	ILE
1	C	176	LEU
1	C	183	GLU
1	C	195	ARG
1	C	200	GLN
1	C	201	LEU
1	C	205	LEU
1	D	11	ASN
1	D	14	LEU
1	D	20	VAL
1	D	30	LEU
1	D	32	GLN
1	D	36	VAL
1	D	40	THR
1	D	55	LEU
1	D	61	ASP
1	D	65	THR
1	D	70	LEU
1	D	94	ARG
1	D	104	ARG
1	D	108	LYS
1	D	127	LEU
1	D	146	LEU
1	D	156	GLU
1	D	165	SER
1	D	195	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	200	GLN
2	Z	1	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	HIS
1	A	81	ASN
1	A	109	GLN
1	A	115	ASN
1	A	151	HIS
1	B	11	ASN
1	B	32	GLN
1	B	66	HIS
1	B	88	ASN
1	B	100	HIS
1	B	109	GLN
1	B	151	HIS
1	B	206	GLN
1	C	66	HIS
1	C	100	HIS
1	C	115	ASN
1	C	129	ASN
1	C	180	GLN
1	C	200	GLN
1	D	11	ASN
1	D	38	GLN
1	D	44	HIS
1	D	81	ASN
1	D	100	HIS
1	D	151	HIS
1	D	152	GLN
1	D	180	GLN
1	D	188	HIS
1	D	200	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/208 (96%)	0.45	14 (7%) 22 21	38, 48, 63, 73	0
1	B	204/208 (98%)	0.51	10 (4%) 35 34	42, 47, 56, 60	0
1	C	206/208 (99%)	0.65	15 (7%) 21 20	36, 46, 58, 70	0
1	D	198/208 (95%)	0.56	9 (4%) 38 38	38, 47, 59, 71	0
2	E	11/17 (64%)	0.75	1 (9%) 15 14	48, 50, 57, 59	0
2	F	12/17 (70%)	1.20	2 (16%) 4 3	49, 53, 60, 61	0
2	G	11/17 (64%)	0.94	0 100 100	48, 50, 54, 58	0
2	H	12/17 (70%)	0.58	1 (8%) 17 17	45, 47, 54, 55	0
2	Z	4/17 (23%)	2.03	2 (50%) 0 0	53, 54, 55, 56	0
All	All	858/917 (93%)	0.57	54 (6%) 26 25	36, 47, 59, 73	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	164	ASP	4.5
1	A	161	PRO	4.3
1	C	205	LEU	4.3
1	A	164	ASP	4.3
1	A	162	THR	4.2
1	B	159	GLU	3.7
1	C	160	THR	3.7
1	C	161	PRO	3.6
1	B	162	THR	3.6
1	B	43	TRP	3.5
1	A	157	GLU	3.5
1	D	160	THR	3.5
1	A	163	THR	3.4
1	C	162	THR	3.4
1	B	4	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	163	THR	3.4
1	C	165	SER	3.3
1	B	163	THR	3.2
1	C	164	ASP	3.2
1	D	3	ARG	3.1
1	D	5	ASP	3.0
1	C	5	ASP	2.9
1	C	153	VAL	2.9
1	C	159	GLU	2.9
1	D	159	GLU	2.9
1	A	3	ARG	2.9
1	C	2	SER	2.8
1	A	5	ASP	2.8
2	F	1	TRP	2.8
1	C	88	ASN	2.8
1	D	203	ALA	2.8
2	Z	2	THR	2.7
1	D	165	SER	2.7
2	H	12	SER	2.7
1	A	160	THR	2.7
1	C	3	ARG	2.7
1	D	157	GLU	2.7
1	A	158	ARG	2.6
1	D	164	ASP	2.5
2	E	11	PRO	2.5
1	B	207	ILE	2.5
1	B	67	PHE	2.4
2	Z	4	ASN	2.3
1	A	202	THR	2.3
1	A	156	GLU	2.3
1	C	4	LEU	2.3
1	C	67	PHE	2.3
1	A	24	GLY	2.2
2	F	12	SER	2.2
1	A	159	GLU	2.1
1	B	71	GLU	2.1
1	D	158	ARG	2.1
1	B	70	LEU	2.0
1	A	165	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.