



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:12 AM UTC

PDB ID : 3NSX / pdb_00003nsx
Title : The crystal structure of the D420A mutant of the alpha-glucosidase (FAMILY 31) from Ruminococcus obeum ATCC 29174
Authors : Tan, K.; Tesar, C.; Wilton, R.; Keigher, L.; Babnigg, G.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-07-02
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	666	5% 91% 8%
1	B	666	3% 91% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12583 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called alpha-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	666	Total	C	N	O	S	0	26	0
			5594	3603	905	1049	37			
1	B	666	Total	C	N	O	S	0	17	0
			5543	3559	908	1041	35			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A5ZY13
A	-1	ASN	-	expression tag	UNP A5ZY13
A	0	ALA	-	expression tag	UNP A5ZY13
A	420	ALA	ASP	engineered mutation	UNP A5ZY13
B	-2	SER	-	expression tag	UNP A5ZY13
B	-1	ASN	-	expression tag	UNP A5ZY13
B	0	ALA	-	expression tag	UNP A5ZY13
B	420	ALA	ASP	engineered mutation	UNP A5ZY13

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	4	1	3		
2	A	1	Total	C	N	O	0	0
			8	4	1	3		
2	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	712	Total	O	0	0
			712	712		
3	B	710	Total	O	0	0
			710	710		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha-glucosidase



- Molecule 1: alpha-glucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.83Å 120.02Å 88.25Å 90.00° 109.10° 90.00°	Depositor
Resolution (Å)	39.73 – 1.57 39.73 – 1.57	Depositor EDS
% Data completeness (in resolution range)	95.5 (39.73-1.57) 99.5 (39.73-1.57)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.57Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.153 , 0.179 0.152 , 0.176	Depositor DCC
R_{free} test set	9733 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	15.9	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12583	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/5815	0.70	2/7838 (0.0%)
1	B	0.40	0/5737	0.70	2/7731 (0.0%)
All	All	0.39	0/11552	0.70	4/15569 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	PHE	CA-C-N	5.17	124.97	119.28
1	A	214	PHE	C-N-CA	5.17	124.97	119.28
1	B	394	ASP	CA-C-N	5.01	124.79	119.28
1	B	394	ASP	C-N-CA	5.01	124.79	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5594	0	5431	62	0
1	B	5543	0	5351	46	0
2	A	16	0	24	2	0
2	B	8	0	12	0	0
3	A	712	0	0	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	710	0	0	14	0
All	All	12583	0	10818	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:MET:HE2	1:A:347:MET:HE2	1.19	1.13
1:A:347:MET:HE3	1:B:343:MET:HG2	1.15	1.13
1:A:347:MET:SD	3:B:1130:HOH:O	2.28	0.91
1:A:347:MET:HE3	1:B:343:MET:CG	2.01	0.90
1:B:153[A]:ARG:HD3	3:B:1124:HOH:O	1.71	0.90
1:A:169:TRP:HD1	1:A:169:TRP:O	1.60	0.84
1:A:343:MET:CE	1:A:347:MET:HE2	2.05	0.83
1:B:416:ILE:HD11	1:B:438:LEU:HD22	1.60	0.83
1:A:659[B]:ARG:HH21	1:A:659[B]:ARG:HG3	1.43	0.83
1:B:253[A]:LYS:HG2	3:B:903:HOH:O	1.83	0.77
1:A:1[A]:MET:HE3	1:A:2:ILE:H	1.50	0.77
1:A:169:TRP:CD2	1:A:481:GLU:HB2	2.20	0.75
1:A:117[B]:ARG:HD3	1:A:119:ASP:OD1	1.88	0.73
1:B:492:GLU:HG3	3:B:970:HOH:O	1.89	0.73
1:A:455:ASP:HA	1:A:484:ARG:HD2	1.72	0.70
1:A:343:MET:HE2	1:A:347:MET:CE	2.12	0.66
1:B:166:GLN:NE2	1:B:171:TYR:OH	2.28	0.66
1:A:492:GLU:HG3	1:A:493:ASN:H	1.61	0.65
1:A:416:ILE:HD11	1:A:438:LEU:HD22	1.79	0.65
1:A:169:TRP:O	1:A:169:TRP:CD1	2.47	0.65
1:B:371:HIS:HE1	1:B:375:HIS:ND1	1.94	0.64
1:B:74:ASP:HB2	3:B:1293:HOH:O	1.98	0.62
1:B:659[A]:ARG:HG2	3:B:1216:HOH:O	1.99	0.62
1:A:371:HIS:HE1	1:A:375:HIS:ND1	1.98	0.61
1:A:169:TRP:CE3	1:A:481:GLU:HB2	2.35	0.61
1:A:495[A]:GLU:H	1:A:495[A]:GLU:CD	2.07	0.60
1:A:169:TRP:CD1	1:A:169:TRP:C	2.79	0.59
1:B:77:HIS:HE1	1:B:419:GLY:O	1.86	0.59
1:B:362:HIS:NE2	1:B:371:HIS:HD2	2.01	0.59
1:B:416:ILE:HD11	1:B:438:LEU:CD2	2.33	0.58
1:A:166[B]:GLN:OE1	1:A:487:GLU:HG3	2.02	0.58
1:A:362:HIS:NE2	1:A:371:HIS:HD2	2.03	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:455:ASP:OD1	1:A:484:ARG:HB3	2.05	0.57
1:B:118:MET:HE2	3:B:981:HOH:O	2.05	0.55
1:A:453:PHE:N	1:A:453:PHE:CD2	2.73	0.55
1:A:481:GLU:HA	3:A:1365:HOH:O	2.06	0.55
1:A:659[B]:ARG:HH21	1:A:659[B]:ARG:CG	2.16	0.55
1:A:491:PHE:O	1:A:494:ILE:HG22	2.06	0.54
1:A:117[A]:ARG:HG3	1:A:117[A]:ARG:NH1	2.22	0.54
1:A:371:HIS:CE1	1:A:375:HIS:ND1	2.75	0.54
1:A:1[B]:MET:SD	1:A:3:ARG:HG3	2.48	0.54
1:B:310:GLU:N	1:B:311:PRO:HA	2.22	0.54
1:B:371:HIS:CE1	1:B:375:HIS:ND1	2.75	0.54
1:A:310:GLU:N	1:A:311:PRO:HA	2.23	0.53
1:A:117[A]:ARG:HG3	1:A:117[A]:ARG:HH11	1.72	0.53
1:B:211:GLU:HB2	3:B:1379:HOH:O	2.09	0.53
1:B:38:PHE:CZ	1:B:132:ILE:HD11	2.43	0.53
1:A:38:PHE:CZ	1:A:132:ILE:HD11	2.44	0.52
1:A:118[A]:MET:HE2	3:A:845:HOH:O	2.08	0.52
1:A:287[A]:LYS:HB2	1:A:287[A]:LYS:NZ	2.24	0.52
1:A:348:LYS:HG2	1:B:344:GLN:NE2	2.26	0.50
1:B:171:TYR:CZ	1:B:180:VAL:HG21	2.47	0.50
1:A:659[B]:ARG:HG3	1:A:659[B]:ARG:NH2	2.21	0.50
1:A:416:ILE:HD11	1:A:438:LEU:CD2	2.41	0.50
1:A:368:LYS:HE2	3:A:716:HOH:O	2.10	0.50
1:B:105:TYR:OH	1:B:127[B]:ASN:ND2	2.45	0.49
1:B:310:GLU:HG3	1:B:405[B]:SER:OG	2.13	0.48
1:B:576:ILE:HD12	1:B:586:GLU:HG2	1.96	0.48
1:A:369[A]:ILE:HD13	3:A:1306:HOH:O	2.14	0.47
1:A:605:LEU:C	1:A:605:LEU:HD12	2.40	0.47
1:A:488:CYS:SG	1:A:501[B]:ILE:HD11	2.54	0.47
1:B:24[A]:GLU:HG3	3:B:1004:HOH:O	2.14	0.47
1:B:631:GLN:NE2	1:B:633:ILE:HD11	2.30	0.47
1:A:169:TRP:CE3	2:A:665:TRS:H11	2.50	0.47
1:B:149[B]:ARG:HD2	1:B:522:ASP:OD2	2.16	0.46
1:B:168:ARG:HD3	1:B:479:ALA:HB3	1.95	0.46
1:A:281[B]:LEU:HD13	1:A:362:HIS:CE1	2.51	0.46
1:B:363:ASN:HB2	1:B:368:LYS:HE2	1.96	0.46
1:A:280:MET:O	1:A:286:ARG:HG3	2.17	0.45
1:A:494:ILE:HG23	3:A:1359:HOH:O	2.16	0.45
1:A:453:PHE:HA	1:A:483:THR:HG22	1.98	0.44
1:B:654:LYS:HB3	1:B:654:LYS:HE3	1.64	0.44
1:B:576:ILE:CD1	1:B:586:GLU:HG2	2.47	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:169:TRP:CD2	1:B:481:GLU:HB3	2.53	0.44
1:A:79[A]:GLU:HG2	3:A:1294:HOH:O	2.16	0.44
1:A:487:GLU:HB3	1:A:489:TYR:CE1	2.52	0.44
1:A:367:LYS:HE2	1:A:369[A]:ILE:HD11	2.00	0.44
1:B:436:PRO:O	1:B:440:MET:HG3	2.18	0.44
1:B:153[A]:ARG:HD2	1:B:520:LEU:HA	1.99	0.43
1:B:199:ASP:HB2	3:B:844:HOH:O	2.19	0.43
1:B:456:ASP:HB2	3:B:1060:HOH:O	2.18	0.43
1:A:166[B]:GLN:HE21	1:A:166[B]:GLN:HB3	1.62	0.42
1:B:172:THR:HG22	1:B:213:ASN:ND2	2.33	0.42
1:B:486:GLN:H	1:B:486:GLN:HG2	1.63	0.42
1:B:149[B]:ARG:HD3	3:B:774:HOH:O	2.20	0.42
1:B:73:ASP:HB3	3:B:940:HOH:O	2.20	0.42
1:A:170:GLY:N	3:A:1365:HOH:O	2.49	0.41
1:A:34:GLN:HA	1:A:37:GLY:O	2.21	0.41
1:A:169:TRP:O	3:A:706:HOH:O	2.22	0.41
1:A:580:PRO:HB3	3:A:765:HOH:O	2.20	0.41
1:B:422:LYS:O	1:B:427:HIS:HE1	2.02	0.41
1:A:483:THR:HB	1:A:486:GLN:HE21	1.86	0.41
1:B:270:VAL:HG12	1:B:271:TRP:H	1.86	0.41
1:A:233:PRO:HD2	1:A:303:GLY:O	2.21	0.41
1:A:576[A]:ILE:HD12	1:A:586:GLU:HG2	2.03	0.41
1:B:280:MET:O	1:B:286:ARG:HG3	2.21	0.41
1:B:430:LEU:HG	1:B:434:MET:HE3	2.03	0.41
1:A:492:GLU:HG3	1:A:493:ASN:N	2.33	0.40
1:B:590:GLU:HG3	3:B:1100:HOH:O	2.20	0.40
1:A:341:TRP:CE3	2:A:664:TRS:H12	2.56	0.40
1:B:81:LYS:HD3	1:B:81:LYS:HA	1.83	0.40
1:A:455:ASP:CA	1:A:484:ARG:HD2	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	690/666 (104%)	668 (97%)	20 (3%)	2 (0%)	36	18
1	B	681/666 (102%)	658 (97%)	22 (3%)	1 (0%)	48	26
All	All	1371/1332 (103%)	1326 (97%)	42 (3%)	3 (0%)	43	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	492	GLU
1	A	270	VAL
1	B	270	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	599/574 (104%)	586 (98%)	13 (2%)	45	18
1	B	590/574 (103%)	581 (98%)	9 (2%)	57	31
All	All	1189/1148 (104%)	1167 (98%)	22 (2%)	55	22

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25[A]	GLU
1	A	25[B]	GLU
1	A	36	GLU
1	A	192	ASP
1	A	270	VAL
1	A	287[A]	LYS
1	A	287[B]	LYS
1	A	305	TRP
1	A	348	LYS
1	A	453	PHE
1	A	494	ILE
1	A	495[A]	GLU
1	A	495[B]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	24[A]	GLU
1	B	24[B]	GLU
1	B	74	ASP
1	B	136	GLU
1	B	270	VAL
1	B	422	LYS
1	B	481	GLU
1	B	486	GLN
1	B	590	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	89	ASN
1	A	306	ASN
1	A	339	HIS
1	A	344	GLN
1	A	371	HIS
1	A	486	GLN
1	A	550	ASN
1	B	77	HIS
1	B	146	GLN
1	B	202	GLN
1	B	306	ASN
1	B	371	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TRS	B	664	-	7,7,7	0.32	0	9,9,9	0.21	0
2	TRS	A	664	-	7,7,7	0.32	0	9,9,9	0.70	0
2	TRS	A	665	-	7,7,7	0.20	0	9,9,9	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	B	664	-	-	6/9/9/9	-
2	TRS	A	664	-	-	0/9/9/9	-
2	TRS	A	665	-	-	3/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	665	TRS	C1-C-C2-O2
2	B	664	TRS	C1-C-C2-O2
2	B	664	TRS	C3-C-C2-O2
2	B	664	TRS	N-C-C2-O2
2	B	664	TRS	C1-C-C3-O3
2	A	665	TRS	C3-C-C2-O2
2	B	664	TRS	C2-C-C3-O3
2	B	664	TRS	N-C-C3-O3
2	A	665	TRS	N-C-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	664	TRS	1	0
2	A	665	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	666/666 (100%)	-0.09	30 (4%)	38 46	6, 17, 41, 100	29 (4%)
1	B	666/666 (100%)	-0.14	23 (3%)	47 55	8, 16, 36, 76	23 (3%)
All	All	1332/1332 (100%)	-0.11	53 (3%)	42 50	6, 16, 38, 100	52 (3%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	453	PHE	6.2
1	A	488	CYS	6.1
1	A	491	PHE	5.7
1	B	21	GLU	4.6
1	A	454	GLY	4.5
1	A	494	ILE	4.4
1	A	480	ALA	4.3
1	B	481	GLU	4.2
1	B	365	ASN	4.1
1	A	169	TRP	4.1
1	A	482	GLY	4.0
1	A	486	GLN	4.0
1	A	1[A]	MET	3.8
1	B	171	TYR	3.7
1	A	455	ASP	3.5
1	A	497	PHE	3.4
1	A	35	LYS	3.4
1	A	483	THR	3.4
1	B	-2	SER	3.3
1	A	452	GLY	3.3
1	B	440	MET	3.3
1	B	169	TRP	3.1
1	A	-2	SER	3.1
1	B	480	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	118[A]	MET	3.0
1	B	453	PHE	3.0
1	A	489	TYR	2.9
1	B	172	THR	2.9
1	A	492	GLU	2.9
1	A	493	ASN	2.8
1	B	482	GLY	2.7
1	A	0	ALA	2.7
1	B	216	ASP	2.6
1	A	484	ARG	2.6
1	A	-1	ASN	2.6
1	A	76	ILE	2.5
1	B	452	GLY	2.5
1	A	36	GLU	2.5
1	B	75	PRO	2.5
1	B	76	ILE	2.5
1	B	215	PRO	2.5
1	A	75	PRO	2.4
1	B	455	ASP	2.4
1	B	421	ASN	2.3
1	B	451	GLY	2.3
1	A	117[A]	ARG	2.3
1	B	653	ASP	2.2
1	B	35	LYS	2.1
1	A	481	GLU	2.1
1	A	19	LYS	2.1
1	A	138[A]	GLU	2.0
1	B	211	GLU	2.0
1	B	-1	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TRS	A	665	8/8	0.59	0.22	28,44,51,54	0
2	TRS	A	664	8/8	0.70	0.17	43,47,52,52	0
2	TRS	B	664	8/8	0.72	0.18	48,51,53,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.