



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2026 – 12:53 PM UTC

PDB ID : 5NS6 / pdb_00005ns6
Title : Crystal structure of beta-glucosidase BglM-G1 from marine metagenome
Authors : Mhaindarkar, D.C.; Gasper, R.; Lupilova, N.; Leichert, L.I.; Hofmann, E.
Deposited on : 2017-04-25
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	 93%
1	B	455	 94%
1	C	455	 95%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 22387 atoms, of which 10542 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	438	Total	C	H	N	O	S	0	15	0
			7067	2335	3459	593	667	13			
1	B	438	Total	C	H	N	O	S	0	12	0
			7042	2322	3450	596	661	13			
1	C	438	Total	C	H	N	O	S	0	13	0
			7042	2323	3449	593	664	13			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	264	Total 264	O 264	0	0
4	B	311	Total 311	O 311	0	0
4	C	284	Total 284	O 284	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-glucosidase



- Molecule 1: Beta-glucosidase



- Molecule 1: Beta-glucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	138.02Å 138.02Å 97.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.80 – 1.50 48.80 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.80-1.50) 100.0 (48.80-1.50)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.50Å)	Xtriage
Refinement program	PHENIX dev_2621	Depositor
R, R_{free}	0.150 , 0.172 0.151 , 0.172	Depositor DCC
R_{free} test set	14633 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	22387	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/3764	0.45	0/5108
1	B	0.29	0/3738	0.48	0/5072
1	C	0.28	0/3742	0.48	0/5078
All	All	0.28	0/11244	0.47	0/15258

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3608	3459	3486	13	0
1	B	3592	3450	3473	10	0
1	C	3593	3449	3471	6	0
2	A	25	0	0	1	0
2	B	15	0	0	0	0
2	C	15	0	0	0	0
3	A	42	56	56	3	0
3	B	54	72	72	2	0
3	C	42	56	56	0	0
4	A	264	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	311	0	0	3	0
4	C	284	0	0	1	0
All	All	11845	10542	10614	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212[B]:GLU:OE1	4:A:601:HOH:O	1.88	0.90
1:C:212[B]:GLU:OE2	4:C:601:HOH:O	1.93	0.85
1:B:212[B]:GLU:OE1	4:B:601:HOH:O	2.05	0.74
1:C:210:SER:OG	1:C:212[A]:GLU:OE1	2.05	0.73
1:A:97[A]:ASP:OD1	4:A:602:HOH:O	2.08	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	451/455 (99%)	442 (98%)	9 (2%)	0	100	100
1	B	448/455 (98%)	440 (98%)	8 (2%)	0	100	100
1	C	449/455 (99%)	441 (98%)	8 (2%)	0	100	100
All	All	1348/1365 (99%)	1323 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/379 (100%)	379 (100%)	0	100	100
1	B	376/379 (99%)	376 (100%)	0	100	100
1	C	377/379 (100%)	377 (100%)	0	100	100
All	All	1132/1137 (100%)	1132 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	350	ASN
1	C	158	ASN
1	B	350	ASN
1	B	158	ASN
1	C	127	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

34 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	SO4	B	503	-	4,4,4	0.24	0	6,6,6	0.07	0
3	GOL	C	507	-	5,5,5	1.13	0	5,5,5	1.05	0
3	GOL	B	509	-	5,5,5	1.01	0	5,5,5	1.04	0
3	GOL	A	512	-	5,5,5	0.96	0	5,5,5	0.93	0
2	SO4	C	503	-	4,4,4	0.24	0	6,6,6	0.12	0
3	GOL	A	511	-	5,5,5	1.04	0	5,5,5	1.16	1 (20%)
3	GOL	B	508	-	5,5,5	1.09	0	5,5,5	1.04	0
2	SO4	B	502	-	4,4,4	0.25	0	6,6,6	0.15	0
3	GOL	C	505	-	5,5,5	1.17	0	5,5,5	1.05	1 (20%)
2	SO4	A	503	-	4,4,4	0.21	0	6,6,6	0.12	0
2	SO4	C	501	-	4,4,4	0.23	0	6,6,6	0.25	0
2	SO4	A	504	-	4,4,4	0.24	0	6,6,6	0.09	0
3	GOL	C	509	-	5,5,5	1.10	0	5,5,5	0.93	0
3	GOL	C	510	-	5,5,5	0.83	0	5,5,5	0.96	0
3	GOL	A	509	-	5,5,5	1.12	0	5,5,5	1.04	0
3	GOL	B	510	-	5,5,5	1.11	0	5,5,5	0.97	0
2	SO4	A	505	-	4,4,4	0.24	0	6,6,6	0.08	0
3	GOL	A	507	-	5,5,5	1.02	0	5,5,5	1.02	0
2	SO4	B	501	-	4,4,4	0.31	0	6,6,6	0.53	0
3	GOL	B	506	-	5,5,5	0.94	0	5,5,5	1.10	0
3	GOL	C	508	-	5,5,5	1.09	0	5,5,5	0.92	0
3	GOL	B	504	-	5,5,5	1.00	0	5,5,5	0.98	0
3	GOL	A	508	-	5,5,5	0.83	0	5,5,5	1.07	0
3	GOL	B	507	-	5,5,5	0.94	0	5,5,5	1.17	1 (20%)
2	SO4	C	502	-	4,4,4	0.22	0	6,6,6	0.16	0
3	GOL	B	512	-	5,5,5	0.96	0	5,5,5	0.85	0
2	SO4	A	502	-	4,4,4	0.25	0	6,6,6	0.06	0
3	GOL	C	506	-	5,5,5	0.98	0	5,5,5	1.02	0
3	GOL	A	506	-	5,5,5	1.31	1 (20%)	5,5,5	0.97	0
3	GOL	C	504	-	5,5,5	1.04	0	5,5,5	0.97	0
3	GOL	B	511	-	5,5,5	1.02	0	5,5,5	0.87	0
3	GOL	B	505	-	5,5,5	1.29	1 (20%)	5,5,5	0.92	0
3	GOL	A	510	-	5,5,5	1.01	0	5,5,5	0.93	0
2	SO4	A	501	-	4,4,4	0.15	0	6,6,6	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	507	-	-	0/4/4/4	-
3	GOL	B	509	-	-	2/4/4/4	-
3	GOL	A	512	-	-	0/4/4/4	-
3	GOL	A	511	-	-	1/4/4/4	-
3	GOL	C	505	-	-	0/4/4/4	-
3	GOL	A	510	-	-	1/4/4/4	-
3	GOL	C	509	-	-	2/4/4/4	-
3	GOL	C	510	-	-	1/4/4/4	-
3	GOL	A	509	-	-	1/4/4/4	-
3	GOL	B	510	-	-	2/4/4/4	-
3	GOL	A	507	-	-	0/4/4/4	-
3	GOL	B	506	-	-	0/4/4/4	-
3	GOL	C	508	-	-	3/4/4/4	-
3	GOL	B	504	-	-	0/4/4/4	-
3	GOL	A	508	-	-	0/4/4/4	-
3	GOL	B	507	-	-	0/4/4/4	-
3	GOL	B	512	-	-	2/4/4/4	-
3	GOL	C	506	-	-	0/4/4/4	-
3	GOL	A	506	-	-	0/4/4/4	-
3	GOL	C	504	-	-	0/4/4/4	-
3	GOL	B	511	-	-	0/4/4/4	-
3	GOL	B	505	-	-	0/4/4/4	-
3	GOL	B	508	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	506	GOL	C1-C2	2.17	1.60	1.51
3	B	505	GOL	C1-C2	2.12	1.59	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	511	GOL	C3-C2-C1	-2.30	103.35	111.80
3	B	507	GOL	C3-C2-C1	-2.06	104.23	111.80
3	C	505	GOL	C3-C2-C1	-2.03	104.36	111.80

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	509	GOL	O1-C1-C2-C3
3	B	512	GOL	C1-C2-C3-O3
3	C	508	GOL	C1-C2-C3-O3
3	B	512	GOL	O2-C2-C3-O3
3	C	508	GOL	O2-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	511	GOL	2	0
2	A	504	SO4	1	0
3	B	510	GOL	1	0
3	B	512	GOL	1	0
3	A	510	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/455 (96%)	-0.29	6 (1%) 73 77	11, 27, 50, 64	15 (3%)
1	B	438/455 (96%)	-0.70	2 (0%) 87 89	9, 20, 35, 59	12 (2%)
1	C	438/455 (96%)	-0.61	2 (0%) 87 89	10, 21, 39, 69	13 (2%)
All	All	1314/1365 (96%)	-0.53	10 (0%) 82 86	9, 22, 42, 69	40 (3%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	335[A]	ARG	4.4
1	A	88[A]	TYR	4.1
1	A	314	ASN	3.0
1	A	312	ALA	3.0
1	C	338	LYS	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	503	5/5	0.68	0.17	72,96,100,172	0
2	SO4	A	504	5/5	0.77	0.17	59,74,92,128	0
3	GOL	A	510	6/6	0.77	0.24	67,85,111,113	0
3	GOL	B	509	6/6	0.77	0.24	61,79,107,108	0
3	GOL	B	512	6/6	0.77	0.17	55,84,102,123	0
3	GOL	C	510	6/6	0.79	0.22	33,50,79,92	0
2	SO4	C	502	5/5	0.81	0.22	69,70,90,127	0
3	GOL	C	508	6/6	0.82	0.20	55,80,92,104	0
2	SO4	A	502	5/5	0.85	0.20	63,75,92,123	0
3	GOL	A	511	6/6	0.87	0.25	25,68,85,85	0
3	GOL	C	507	6/6	0.88	0.18	23,37,85,85	0
2	SO4	A	505	5/5	0.88	0.11	55,62,94,98	0
2	SO4	B	502	5/5	0.88	0.12	54,56,63,88	0
3	GOL	A	506	6/6	0.89	0.23	28,77,93,107	0
3	GOL	A	507	6/6	0.89	0.17	43,63,77,92	0
3	GOL	A	509	6/6	0.89	0.16	28,43,72,81	0
2	SO4	A	503	5/5	0.89	0.23	48,75,109,143	0
2	SO4	C	503	5/5	0.89	0.14	60,64,79,88	0
3	GOL	B	511	6/6	0.91	0.16	21,33,64,70	0
3	GOL	B	508	6/6	0.91	0.13	21,30,48,58	0
3	GOL	C	505	6/6	0.91	0.15	28,44,62,66	0
3	GOL	C	509	6/6	0.92	0.19	18,57,71,85	0
3	GOL	B	510	6/6	0.93	0.15	18,43,70,70	0
3	GOL	A	512	6/6	0.93	0.12	23,35,64,77	0
3	GOL	A	508	6/6	0.93	0.15	37,62,76,83	0
2	SO4	C	501	5/5	0.93	0.11	29,31,65,66	0
3	GOL	B	506	6/6	0.94	0.13	25,48,74,76	0
3	GOL	B	507	6/6	0.94	0.15	23,52,74,74	0
3	GOL	B	505	6/6	0.94	0.15	26,59,75,75	0
3	GOL	C	504	6/6	0.95	0.09	29,37,52,52	0
3	GOL	C	506	6/6	0.95	0.12	30,62,80,90	0
2	SO4	A	501	5/5	0.98	0.08	29,29,36,38	0
2	SO4	B	501	5/5	0.98	0.06	22,23,29,33	0
3	GOL	B	504	6/6	0.98	0.05	20,24,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.