



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:15 PM UTC

PDB ID : 5NSK / pdb_00005nsk
Title : apo Structure of Leucyl aminopeptidase from Trypanosoma brucei
Authors : Timm, J.; Wilson, K.S.
Deposited on : 2017-04-26
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	4% 92% ...
1	B	521	5% 91% 5% ..
1	C	521	3% 93% ...
1	D	521	4% 93% ..
1	E	521	5% 91% 5% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	521	4% 92% ...

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 22652 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	507	Total	C	N	O	S	0	0	0
			3683	2329	627	707	20			
1	B	506	Total	C	N	O	S	0	0	0
			3683	2327	628	709	19			
1	C	506	Total	C	N	O	S	0	0	0
			3700	2335	632	714	19			
1	D	506	Total	C	N	O	S	0	0	0
			3675	2320	627	709	19			
1	E	506	Total	C	N	O	S	0	0	0
			3704	2342	633	709	20			
1	F	508	Total	C	N	O	S	0	0	0
			3707	2342	632	713	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	ALA	THR	conflict	UNP Q385B0
A	140	THR	ALA	conflict	UNP Q385B0
B	33	ALA	THR	conflict	UNP Q385B0
B	140	THR	ALA	conflict	UNP Q385B0
C	33	ALA	THR	conflict	UNP Q385B0
C	140	THR	ALA	conflict	UNP Q385B0
D	33	ALA	THR	conflict	UNP Q385B0
D	140	THR	ALA	conflict	UNP Q385B0
E	33	ALA	THR	conflict	UNP Q385B0
E	140	THR	ALA	conflict	UNP Q385B0
F	33	ALA	THR	conflict	UNP Q385B0
F	140	THR	ALA	conflict	UNP Q385B0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	83	Total 83	O 83	0	0
2	B	62	Total 62	O 62	0	0
2	C	119	Total 119	O 119	0	0
2	D	71	Total 71	O 71	0	0
2	E	87	Total 87	O 87	0	0
2	F	78	Total 78	O 78	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aminopeptidase, putative




- Molecule 1: Aminopeptidase, putative




- Molecule 1: Aminopeptidase, putative





- Molecule 1: Aminopeptidase, putative







- Molecule 1: Aminopeptidase, putative



- Molecule 1: Aminopeptidase, putative



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	162.33Å 162.45Å 176.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.61 – 2.60 72.61 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (72.61-2.60) 100.0 (72.61-2.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.62Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.191 , 0.211 0.196 , 0.213	Depositor DCC
R_{free} test set	7257 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	38.1	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 25.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.013 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	22652	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.18	3/3757 (0.1%)	0.98	1/5134 (0.0%)
1	B	1.20	4/3757 (0.1%)	1.02	6/5134 (0.1%)
1	C	1.21	3/3774 (0.1%)	0.99	2/5155 (0.0%)
1	D	1.16	4/3748 (0.1%)	1.00	3/5122 (0.1%)
1	E	1.22	7/3778 (0.2%)	1.02	5/5159 (0.1%)
1	F	1.21	6/3782 (0.2%)	1.01	3/5169 (0.1%)
All	All	1.20	27/22596 (0.1%)	1.00	20/30873 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	E	0	1
All	All	0	3

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	36	GLU	N-CA	7.67	1.55	1.46
1	F	36	GLU	N-CA	7.23	1.55	1.46
1	A	36	GLU	N-CA	7.22	1.55	1.46
1	F	294	TYR	C-O	-7.08	1.15	1.24
1	B	36	GLU	N-CA	7.02	1.54	1.46
1	D	36	GLU	N-CA	6.89	1.54	1.46
1	D	198	LYS	N-CA	6.09	1.53	1.46
1	B	310	LYS	N-CA	6.05	1.53	1.46
1	C	310	LYS	N-CA	6.02	1.53	1.46
1	F	196	PRO	CA-C	5.95	1.55	1.51
1	D	278	LYS	N-CA	5.90	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	280	THR	CA-C	5.81	1.58	1.52
1	E	136	LYS	N-CA	5.60	1.53	1.46
1	F	234	ALA	CA-C	5.43	1.59	1.52
1	C	14	SER	CA-C	5.43	1.60	1.52
1	E	40	LYS	CA-C	5.31	1.59	1.53
1	E	136	LYS	CA-C	5.29	1.59	1.52
1	B	280	THR	N-CA	5.27	1.53	1.46
1	A	39	GLN	N-CA	5.14	1.52	1.45
1	A	11	LYS	N-CA	5.13	1.52	1.46
1	E	310	LYS	N-CA	5.11	1.52	1.46
1	F	120	ARG	C-O	-5.10	1.21	1.23
1	E	63	GLN	N-CA	5.10	1.52	1.46
1	B	235	LYS	N-CA	5.05	1.52	1.46
1	C	40	LYS	CA-C	5.04	1.59	1.53
1	F	278	LYS	N-CA	5.01	1.52	1.46
1	D	11	LYS	N-CA	5.00	1.52	1.46

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	96	ALA	N-CA-C	-8.37	98.80	110.35
1	D	96	ALA	N-CA-C	-7.92	99.41	110.35
1	B	135	VAL	N-CA-CB	7.54	120.59	110.26
1	D	280	THR	N-CA-CB	6.68	120.20	110.32
1	B	269	ARG	CG-CD-NE	-6.14	98.50	112.00
1	D	135	VAL	N-CA-CB	6.07	117.14	110.53
1	C	519	ARG	NE-CZ-NH2	5.81	124.43	119.20
1	E	163	LYS	CB-CA-C	5.81	119.58	109.65
1	F	151	VAL	N-CA-C	5.67	116.40	110.62
1	E	135	VAL	N-CA-C	-5.64	100.30	108.36
1	C	14	SER	CB-CA-C	5.38	121.01	110.67
1	B	58	ARG	CG-CD-NE	-5.30	100.33	112.00
1	B	36	GLU	N-CA-CB	5.30	117.91	110.12
1	F	160	ARG	CB-CA-C	-5.18	99.48	110.31
1	A	184	ARG	CB-CA-C	5.16	118.47	109.65
1	E	136	LYS	N-CA-C	5.15	119.69	113.41
1	B	160	ARG	CB-CA-C	-5.14	99.56	110.31
1	F	196	PRO	N-CA-C	5.13	115.39	110.47
1	B	454	LYS	N-CA-CB	5.09	117.38	109.48
1	E	486	LYS	CA-CB-CG	5.04	124.17	114.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	96	ALA	Peptide
1	B	96	ALA	Peptide
1	E	97	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3683	0	3559	12	0
1	B	3683	0	3563	12	0
1	C	3700	0	3586	10	0
1	D	3675	0	3539	6	0
1	E	3704	0	3611	10	0
1	F	3707	0	3583	11	0
2	A	83	0	0	0	0
2	B	62	0	0	1	0
2	C	119	0	0	0	0
2	D	71	0	0	0	0
2	E	87	0	0	1	0
2	F	78	0	0	0	0
All	All	22652	0	21441	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:516:GLU:OE1	1:F:519:ARG:NH1	2.26	0.68
1:B:114:ARG:NH2	1:C:419:VAL:O	2.28	0.66
1:A:340:CYS:HB3	1:A:377:VAL:HG13	1.80	0.64
1:B:419:VAL:O	1:C:114:ARG:NH2	2.30	0.64
1:C:352:ARG:HG2	1:F:295:ASP:HA	1.82	0.61
1:F:340:CYS:HB3	1:F:377:VAL:HG13	1.82	0.61
1:E:340:CYS:HB3	1:E:377:VAL:HG13	1.83	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:340:CYS:HB3	1:B:377:VAL:HG13	1.81	0.60
1:A:114:ARG:NH2	1:E:419:VAL:O	2.33	0.60
1:D:340:CYS:HB3	1:D:377:VAL:HG13	1.84	0.60
1:A:419:VAL:O	1:E:114:ARG:NH2	2.35	0.59
1:D:114:ARG:NH2	1:F:419:VAL:O	2.35	0.59
1:B:89:VAL:HG11	1:B:92:LEU:HD22	1.84	0.59
1:C:340:CYS:HB3	1:C:377:VAL:HG13	1.85	0.57
1:C:516:GLU:OE1	1:C:519:ARG:NH2	2.36	0.57
1:B:96:ALA:O	1:B:97:GLY:C	2.52	0.52
1:F:464:MET:HE1	1:F:471:GLY:HA2	1.92	0.51
1:C:360:LYS:HG3	1:C:383:PHE:CD2	2.45	0.51
1:B:360:LYS:HG3	1:B:383:PHE:CD2	2.46	0.51
1:B:464:MET:HE1	1:B:471:GLY:HA2	1.93	0.51
1:D:464:MET:HE1	1:D:471:GLY:HA2	1.95	0.49
1:A:360:LYS:HG3	1:A:383:PHE:CD2	2.48	0.49
1:A:96:ALA:O	1:A:97:GLY:C	2.55	0.49
1:C:464:MET:HE1	1:C:471:GLY:HA2	1.94	0.48
1:E:360:LYS:HG3	1:E:383:PHE:CD2	2.48	0.48
1:B:132:LEU:O	1:B:135:VAL:HG22	2.14	0.48
1:A:464:MET:HE1	1:A:471:GLY:HA2	1.95	0.48
1:D:360:LYS:HG3	1:D:383:PHE:CD2	2.48	0.47
1:B:96:ALA:O	1:B:97:GLY:O	2.31	0.47
1:F:360:LYS:HG3	1:F:383:PHE:CD2	2.50	0.47
1:A:120:ARG:NH1	1:A:122:ASP:OD2	2.48	0.46
1:A:96:ALA:O	1:A:97:GLY:O	2.35	0.46
1:E:464:MET:HE1	1:E:471:GLY:HA2	1.97	0.45
1:E:141:THR:OG1	1:E:184:ARG:NH1	2.52	0.43
1:D:120:ARG:NH1	1:D:122:ASP:OD2	2.51	0.43
1:A:74:GLU:HG3	1:A:78:ARG:HD2	2.01	0.43
1:E:182:VAL:HG11	2:E:679:HOH:O	2.19	0.43
1:B:375:ARG:HD2	1:B:462:ASN:O	2.18	0.43
1:F:120:ARG:NH1	1:F:122:ASP:OD2	2.51	0.42
1:F:375:ARG:HD2	1:F:462:ASN:O	2.19	0.42
1:E:375:ARG:HD2	1:E:462:ASN:O	2.19	0.41
1:C:254:TYR:CE1	1:C:360:LYS:HG2	2.55	0.41
1:F:69:ASN:OD1	1:F:69:ASN:C	2.64	0.41
1:B:52:GLY:HA2	2:B:635:HOH:O	2.20	0.41
1:D:375:ARG:HD2	1:D:462:ASN:O	2.21	0.41
1:F:259:SER:HA	1:F:262:LYS:HE2	2.03	0.41
1:C:259:SER:HA	1:C:262:LYS:HE2	2.03	0.41
1:F:464:MET:H	1:F:464:MET:HG2	1.69	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:TYR:CE1	1:A:360:LYS:HG2	2.56	0.41
1:E:259:SER:HA	1:E:262:LYS:HE2	2.02	0.41
1:C:375:ARG:HD2	1:C:462:ASN:O	2.22	0.40
1:E:94:ASN:O	1:E:96:ALA:O	2.40	0.40
1:A:263:ALA:CB	1:A:350:SER:HB3	2.52	0.40
1:A:375:ARG:HD2	1:A:462:ASN:O	2.21	0.40
1:B:263:ALA:CB	1:B:350:SER:HB3	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	503/521 (96%)	492 (98%)	10 (2%)	1 (0%)	43	66
1	B	502/521 (96%)	488 (97%)	12 (2%)	2 (0%)	30	51
1	C	502/521 (96%)	491 (98%)	11 (2%)	0	100	100
1	D	502/521 (96%)	490 (98%)	12 (2%)	0	100	100
1	E	502/521 (96%)	491 (98%)	11 (2%)	0	100	100
1	F	504/521 (97%)	493 (98%)	11 (2%)	0	100	100
All	All	3015/3126 (96%)	2945 (98%)	67 (2%)	3 (0%)	48	70

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	97	GLY
1	A	97	GLY
1	B	280	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/411 (91%)	370 (99%)	4 (1%)	65	84
1	B	376/411 (92%)	369 (98%)	7 (2%)	50	75
1	C	380/411 (92%)	374 (98%)	6 (2%)	55	79
1	D	373/411 (91%)	369 (99%)	4 (1%)	65	84
1	E	381/411 (93%)	376 (99%)	5 (1%)	61	82
1	F	379/411 (92%)	373 (98%)	6 (2%)	55	79
All	All	2263/2466 (92%)	2231 (99%)	32 (1%)	59	81

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	262	LYS
1	A	274	LEU
1	A	351	TYR
1	A	377	VAL
1	B	81	GLU
1	B	110	THR
1	B	135	VAL
1	B	262	LYS
1	B	274	LEU
1	B	351	TYR
1	B	377	VAL
1	C	98	SER
1	C	110	THR
1	C	262	LYS
1	C	274	LEU
1	C	351	TYR
1	C	377	VAL
1	D	262	LYS
1	D	274	LEU
1	D	351	TYR
1	D	377	VAL
1	E	5	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	262	LYS
1	E	274	LEU
1	E	351	TYR
1	E	377	VAL
1	F	154	ILE
1	F	160	ARG
1	F	262	LYS
1	F	274	LEU
1	F	351	TYR
1	F	377	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	HIS
1	A	186	GLN
1	A	311	HIS
1	B	186	GLN
1	C	54	HIS
1	C	115	ASN
1	C	186	GLN
1	D	54	HIS
1	D	311	HIS
1	E	115	ASN
1	E	186	GLN
1	E	211	GLN
1	F	54	HIS
1	F	115	ASN
1	F	211	GLN
1	F	413	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	507/521 (97%)	0.01	22 (4%)	40	34	26, 35, 68, 115	0
1	B	506/521 (97%)	0.27	27 (5%)	32	26	28, 40, 70, 130	0
1	C	506/521 (97%)	-0.24	16 (3%)	50	44	22, 30, 63, 121	0
1	D	506/521 (97%)	0.01	19 (3%)	44	38	28, 37, 70, 116	0
1	E	506/521 (97%)	0.01	26 (5%)	33	28	24, 34, 63, 133	0
1	F	508/521 (97%)	0.12	20 (3%)	43	38	25, 36, 64, 116	0
All	All	3039/3126 (97%)	0.03	130 (4%)	40	34	22, 36, 68, 133	0

All (130) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	294	TYR	8.1
1	E	346	ILE	7.7
1	F	295	ASP	7.4
1	A	294	TYR	7.2
1	B	294	TYR	7.1
1	C	346	ILE	7.1
1	F	346	ILE	6.9
1	A	306	MET	6.8
1	F	306	MET	6.8
1	D	294	TYR	6.8
1	E	5	LEU	6.5
1	E	294	TYR	6.4
1	A	346	ILE	5.7
1	B	348	PRO	5.2
1	D	346	ILE	5.2
1	D	306	MET	5.2
1	F	307	LYS	5.1
1	E	306	MET	5.0
1	A	4	THR	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	5	LEU	4.8
1	A	348	PRO	4.7
1	B	306	MET	4.7
1	B	346	ILE	4.6
1	D	139	ASN	4.6
1	E	349	LYS	4.5
1	F	294	TYR	4.3
1	C	347	GLY	4.3
1	C	306	MET	4.3
1	A	307	LYS	4.2
1	C	348	PRO	4.2
1	B	136	LYS	4.1
1	E	348	PRO	4.0
1	D	349	LYS	4.0
1	D	265	PHE	4.0
1	C	136	LYS	4.0
1	C	307	LYS	3.9
1	A	347	GLY	3.8
1	A	139	ASN	3.7
1	B	137	GLU	3.7
1	C	265	PHE	3.6
1	D	5	LEU	3.6
1	D	348	PRO	3.6
1	C	5	LEU	3.6
1	F	4	THR	3.6
1	D	345	ALA	3.5
1	D	307	LYS	3.5
1	C	349	LYS	3.5
1	A	349	LYS	3.5
1	E	139	ASN	3.4
1	E	347	GLY	3.4
1	D	137	GLU	3.4
1	E	137	GLU	3.4
1	B	139	ASN	3.4
1	B	307	LYS	3.3
1	E	40	LYS	3.3
1	F	5	LEU	3.3
1	A	136	LYS	3.3
1	B	265	PHE	3.3
1	E	350	SER	3.2
1	B	292	ILE	3.2
1	B	138	PRO	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	278	LYS	3.1
1	E	293	VAL	3.1
1	B	351	TYR	3.1
1	E	36	GLU	3.1
1	B	349	LYS	3.0
1	D	136	LYS	3.0
1	B	41	GLY	3.0
1	F	348	PRO	3.0
1	E	138	PRO	3.0
1	D	135	VAL	2.9
1	E	92	LEU	2.9
1	D	293	VAL	2.8
1	F	349	LYS	2.8
1	A	5	LEU	2.8
1	C	293	VAL	2.8
1	E	136	LYS	2.7
1	B	347	GLY	2.7
1	D	347	GLY	2.7
1	B	6	PRO	2.7
1	B	280	THR	2.7
1	B	345	ALA	2.7
1	B	135	VAL	2.7
1	E	307	LYS	2.7
1	E	351	TYR	2.6
1	A	137	GLU	2.6
1	C	91	GLY	2.6
1	A	97	GLY	2.5
1	A	278	LYS	2.5
1	B	293	VAL	2.5
1	A	138	PRO	2.4
1	F	91	GLY	2.4
1	D	351	TYR	2.4
1	C	350	SER	2.4
1	F	36	GLU	2.4
1	B	93	ALA	2.4
1	C	345	ALA	2.4
1	B	64	ARG	2.4
1	A	345	ALA	2.4
1	F	280	THR	2.3
1	A	38	ASN	2.3
1	F	292	ILE	2.3
1	A	265	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	292	ILE	2.3
1	A	309	MET	2.2
1	D	138	PRO	2.2
1	F	133	ASP	2.2
1	B	42	VAL	2.2
1	E	345	ALA	2.2
1	E	41	GLY	2.2
1	A	293	VAL	2.2
1	E	309	MET	2.2
1	F	293	VAL	2.2
1	B	36	GLU	2.2
1	F	136	LYS	2.2
1	E	280	THR	2.1
1	E	292	ILE	2.1
1	E	265	PHE	2.1
1	E	266	GLU	2.1
1	B	278	LYS	2.1
1	C	40	LYS	2.1
1	F	278	LYS	2.1
1	D	352	ARG	2.0
1	F	63	GLN	2.0
1	B	35	TYR	2.0
1	A	266	GLU	2.0
1	E	134	GLU	2.0
1	F	345	ALA	2.0
1	F	140	THR	2.0
1	C	292	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.