



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:43 AM UTC

PDB ID : 5NST / pdb_00005nst
Title : Human monoclonal antibody with a LAIR1 insertion
Authors : Hsieh, F.-L.; Higgins, M.K.
Deposited on : 2017-04-26
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	0% 78% 18% • •
1	C	214	6% 75% 20% • •
2	B	360	4% 74% 17% • 7%
2	D	360	5% 72% 15% • 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Light Chain of antibody MGD21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1618	1005	280	325	8			
1	C	208	Total	C	N	O	S	0	0	0
			1593	989	274	322	8			

- Molecule 2 is a protein called Heavy Chain of antibody MGD21.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	336	Total	C	N	O	S	0	0	0
			2578	1620	439	510	9			
2	D	322	Total	C	N	O	S	0	0	0
			2475	1554	424	489	8			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	O	0	0
			5	5		
5	B	10	Total	O	0	0
			10	10		
5	C	14	Total	O	0	0
			14	14		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	16	Total	O	0	0
			16	16		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Light Chain of antibody MGD21




- Molecule 1: Light Chain of antibody MGD21



- Molecule 2: Heavy Chain of antibody MGD21



- Molecule 2: Heavy Chain of antibody MGD21



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	169.82Å 86.46Å 104.00Å 90.00° 126.65° 90.00°	Depositor
Resolution (Å)	81.90 – 2.52 81.90 – 2.52	Depositor EDS
% Data completeness (in resolution range)	99.7 (81.90-2.52) 99.7 (81.90-2.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.219 , 0.267 0.228 , 0.278	Depositor DCC
R_{free} test set	1968 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	51.9	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 69.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8361	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/1656	1.22	6/2247 (0.3%)
1	C	0.90	1/1630 (0.1%)	1.23	3/2213 (0.1%)
2	B	0.91	1/2639 (0.0%)	1.27	14/3606 (0.4%)
2	D	0.89	1/2533 (0.0%)	1.28	15/3460 (0.4%)
All	All	0.90	3/8458 (0.0%)	1.25	38/11526 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	68	VAL	CA-C	5.56	1.59	1.52
1	C	4	MET	SD-CE	-5.54	1.65	1.79
2	D	107	ASP	CA-C	5.12	1.59	1.52

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	242	ASN	CA-CB-CG	8.48	121.08	112.60
2	D	224	PRO	CA-C-N	6.47	130.30	121.05
2	D	224	PRO	C-N-CA	6.47	130.30	121.05
2	B	30	ASN	CA-CB-CG	6.29	118.89	112.60
2	B	64	LEU	CA-C-N	6.19	128.86	120.38
2	B	64	LEU	C-N-CA	6.19	128.86	120.38
2	B	195	GLU	N-CA-C	-5.92	101.72	110.48
2	B	241	GLY	N-CA-C	-5.85	104.74	112.82
2	B	104	SER	N-CA-C	-5.76	100.44	109.25
2	B	68	VAL	N-CA-C	5.69	117.19	108.71
2	B	65	LYS	N-CA-C	5.68	118.21	111.33
2	D	108	THR	CA-C-N	5.67	128.13	120.65
2	D	108	THR	C-N-CA	5.67	128.13	120.65
2	D	56	GLY	CA-C-N	5.66	128.75	120.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	56	GLY	C-N-CA	5.66	128.75	120.71
1	C	141	PRO	CA-C-N	5.60	128.05	120.38
1	C	141	PRO	C-N-CA	5.60	128.05	120.38
2	D	27	ASP	CA-CB-CG	5.58	118.18	112.60
2	B	243	GLY	N-CA-C	5.57	119.21	112.48
2	D	241	GLY	N-CA-C	-5.57	105.14	112.82
1	A	210	ASN	CA-C-N	5.56	131.71	121.70
1	A	210	ASN	C-N-CA	5.56	131.71	121.70
1	C	188	ARG	N-CA-C	5.36	116.80	109.18
2	B	27	ASP	CA-CB-CG	5.35	117.95	112.60
1	A	17	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	82	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	165	ASP	CA-CB-CG	5.32	117.92	112.60
2	B	167	GLU	N-CA-C	5.20	116.98	109.07
2	B	84	ASP	CA-CB-CG	5.12	117.72	112.60
1	A	145	ASN	CA-CB-CG	5.08	117.68	112.60
2	D	195	GLU	N-CA-C	-5.07	102.78	110.28
2	D	47	GLU	CB-CG-CD	5.06	121.19	112.60
2	B	145	GLU	CA-C-N	5.05	130.18	122.65
2	B	145	GLU	C-N-CA	5.05	130.18	122.65
2	D	298	SER	N-CA-C	5.01	117.13	111.11
2	D	104	SER	N-CA-C	-5.01	100.14	110.80
2	D	44	LYS	CA-C-N	5.00	127.45	122.30
2	D	44	LYS	C-N-CA	5.00	127.45	122.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1618	0	1547	18	0
1	C	1593	0	1527	20	0
2	B	2578	0	2525	27	0
2	D	2475	0	2420	20	0
3	A	12	0	16	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	6	0	8	3	0
3	C	6	0	8	0	0
4	B	14	0	13	0	0
4	D	14	0	13	0	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
5	C	14	0	0	0	0
5	D	16	0	0	0	0
All	All	8361	0	8077	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:224:PRO:HG2	2:B:227:GLU:HB2	1.43	1.01
2:B:169:ARG:HH22	3:B:402:GOL:H12	1.39	0.86
2:D:105:GLN:HG2	2:D:106:ARG:O	1.80	0.81
2:B:224:PRO:HG2	2:B:227:GLU:CB	2.11	0.79
1:C:29:ILE:HG13	1:C:68:GLY:O	1.88	0.74
1:A:59:PRO:HG2	1:A:61:ARG:HH21	1.55	0.72
1:C:48:ILE:HD13	1:C:54:LEU:HD23	1.71	0.70
1:C:48:ILE:CD1	1:C:54:LEU:HD23	2.22	0.68
2:D:105:GLN:HG2	2:D:106:ARG:N	2.09	0.67
2:B:41:ALA:HB3	2:B:44:LYS:HB3	1.79	0.64
2:D:41:ALA:HB3	2:D:44:LYS:HB2	1.80	0.63
1:C:150:ILE:HD11	1:C:179:LEU:HD21	1.82	0.61
2:B:169:ARG:HH22	3:B:402:GOL:C1	2.15	0.59
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.85	0.58
2:D:67:ARG:HD2	2:D:85:SER:HB2	1.85	0.58
1:A:90:GLN:NE2	1:A:96:PRO:HA	2.19	0.57
1:A:195:GLU:HG2	1:A:206:VAL:HG12	1.86	0.57
1:C:195:GLU:HG2	1:C:206:VAL:HG12	1.86	0.57
1:A:61:ARG:HH12	1:A:79:GLN:HB2	1.70	0.57
2:B:44:LYS:HG3	2:B:45:GLY:N	2.19	0.57
2:B:184:ARG:HG3	2:B:199:TYR:CD1	2.40	0.57
2:B:12:MET:HE3	2:B:18:LEU:HA	1.89	0.55
1:C:94:PHE:O	2:D:227:GLU:HB3	2.08	0.53
1:A:150:ILE:HD11	1:A:179:LEU:HD21	1.89	0.53
2:B:136:PRO:O	2:B:166:SER:HB3	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:146:ARG:O	2:B:150:TYR:HA	2.09	0.53
1:C:181:LEU:HD13	1:C:186:TYR:HB2	1.89	0.52
2:D:175:VAL:HA	2:D:179:ASN:HD22	1.73	0.52
2:B:149:ASN:HB3	2:B:151:LEU:HG	1.92	0.52
2:B:146:ARG:HD3	2:B:151:LEU:HD12	1.91	0.51
2:D:39:ARG:HB3	2:D:49:ILE:HD11	1.92	0.51
1:A:12:SER:HB3	1:A:107:LYS:HB2	1.91	0.51
1:A:109:ALA:HA	2:D:298:SER:HB3	1.92	0.51
2:D:146:ARG:HD2	2:D:151:LEU:HD12	1.91	0.51
1:A:29:ILE:HD11	1:A:92:ASN:HB3	1.93	0.51
1:C:12:SER:HB3	1:C:107:LYS:HB2	1.92	0.51
1:A:90:GLN:HE22	1:A:96:PRO:HA	1.77	0.50
1:C:118:PHE:HE1	1:C:135:PHE:CD2	2.29	0.50
1:C:163:TRP:CD1	1:C:175:MET:HG3	2.46	0.50
2:B:64:LEU:HD23	2:B:64:LEU:N	2.27	0.49
2:B:39:ARG:HB3	2:B:49:ILE:HD11	1.94	0.49
1:C:37:GLN:HB2	1:C:47:LEU:HD11	1.93	0.49
2:B:298:SER:HB3	1:C:109:ALA:HA	1.95	0.48
1:A:14:SER:HB3	1:C:169:LYS:HD3	1.96	0.48
2:B:169:ARG:NH2	3:B:402:GOL:H12	2.18	0.47
1:C:18:LYS:HG3	1:C:76:THR:HG22	1.97	0.47
1:A:163:TRP:CD1	1:A:175:MET:HG3	2.49	0.47
2:B:44:LYS:CG	2:B:45:GLY:N	2.78	0.47
2:D:300:VAL:HA	2:D:317:SER:O	2.15	0.47
2:B:175:VAL:HA	2:B:179:ASN:HD22	1.81	0.46
1:A:169:LYS:HD3	1:C:14:SER:HB3	1.98	0.46
2:D:61:ASN:HB3	2:D:64:LEU:HD12	1.97	0.46
1:C:28:HIS:HA	1:C:69:THR:HG23	1.97	0.45
1:A:119:PRO:HB3	1:A:209:PHE:CE1	2.51	0.45
1:A:118:PHE:HE2	1:A:135:PHE:CD2	2.35	0.45
1:C:93:CYS:HB3	2:D:227:GLU:HB2	1.99	0.45
2:D:5:VAL:HG22	2:D:23:ALA:HB3	1.98	0.45
1:C:95:PRO:HG3	2:D:62:PRO:HD2	1.99	0.45
2:B:225:GLN:C	2:B:227:GLU:H	2.24	0.45
1:A:118:PHE:HE2	1:A:135:PHE:CE2	2.35	0.44
2:B:17:THR:HA	2:B:83:VAL:O	2.17	0.44
2:B:64:LEU:HD23	2:B:64:LEU:H	1.82	0.43
2:B:4:LEU:HD22	2:B:24:VAL:HG22	2.00	0.43
2:D:4:LEU:HD22	2:D:24:VAL:HG22	2.01	0.42
2:B:97:ALA:HB1	2:B:237:VAL:HG13	2.00	0.42
2:D:256:PRO:HB3	2:D:282:TYR:HB3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:91:THR:HG23	2:D:247:THR:HA	2.00	0.42
2:D:97:ALA:HB1	2:D:237:VAL:HG13	2.01	0.42
2:B:16:GLY:O	2:B:86:VAL:HG22	2.20	0.42
2:D:186:ILE:HB	2:D:196:GLN:HG3	2.02	0.41
1:A:41:GLY:H	3:A:301:GOL:H2	1.85	0.41
2:B:186:ILE:HB	2:B:196:GLN:HG3	2.01	0.41
1:C:27:GLN:HG2	1:C:28:HIS:H	1.85	0.41
2:B:155:THR:HG22	2:B:170:PHE:CZ	2.56	0.40
2:B:113:PRO:HD2	2:B:195:GLU:O	2.21	0.40
2:D:145:GLU:HB2	2:D:184:ARG:HG3	2.04	0.40
1:A:115:VAL:HA	1:A:135:PHE:O	2.21	0.40
1:C:115:VAL:HA	1:C:135:PHE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/214 (97%)	199 (96%)	9 (4%)	0	100	100
1	C	206/214 (96%)	194 (94%)	12 (6%)	0	100	100
2	B	330/360 (92%)	313 (95%)	17 (5%)	0	100	100
2	D	314/360 (87%)	301 (96%)	13 (4%)	0	100	100
All	All	1058/1148 (92%)	1007 (95%)	51 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/188 (98%)	169 (91%)	16 (9%)	10	19
1	C	182/188 (97%)	164 (90%)	18 (10%)	7	15
2	B	296/316 (94%)	269 (91%)	27 (9%)	9	17
2	D	285/316 (90%)	256 (90%)	29 (10%)	7	14
All	All	948/1008 (94%)	858 (90%)	90 (10%)	8	16

All (90) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	7	SER
1	A	14	SER
1	A	30	ASN
1	A	33	LEU
1	A	65	THR
1	A	67	SER
1	A	76	THR
1	A	90	GLN
1	A	100	GLN
1	A	103	ARG
1	A	105	GLU
1	A	123	GLU
1	A	145	ASN
1	A	165	ASP
1	A	181	LEU
2	B	7	THR
2	B	13	LYS
2	B	25	SER
2	B	33	ARG
2	B	44	LYS
2	B	65	LYS
2	B	66	SER
2	B	71	SER
2	B	76	LYS
2	B	108	THR
2	B	109	ASP
2	B	114	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	145	GLU
2	B	148	ARG
2	B	162	SER
2	B	165	GLU
2	B	196	GLN
2	B	205	LYS
2	B	227	GLU
2	B	254	THR
2	B	287	VAL
2	B	295	SER
2	B	300	VAL
2	B	328	GLU
2	B	329	THR
2	B	331	THR
2	B	348	VAL
1	C	7	SER
1	C	14	SER
1	C	24	ARG
1	C	26	SER
1	C	29	ILE
1	C	31	ASP
1	C	32	SER
1	C	33	LEU
1	C	65	THR
1	C	103	ARG
1	C	105	GLU
1	C	123	GLU
1	C	145	ASN
1	C	151	ASP
1	C	181	LEU
1	C	183	LYS
1	C	187	GLU
1	C	188	ARG
2	D	5	VAL
2	D	7	THR
2	D	21	THR
2	D	25	SER
2	D	51	GLU
2	D	64	LEU
2	D	69	THR
2	D	71	SER
2	D	76	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	80	SER
2	D	83	VAL
2	D	108	THR
2	D	109	ASP
2	D	151	LEU
2	D	162	SER
2	D	165	GLU
2	D	175	VAL
2	D	184	ARG
2	D	196	GLN
2	D	205	LYS
2	D	227	GLU
2	D	228	LEU
2	D	270	ASN
2	D	273	VAL
2	D	287	VAL
2	D	295	SER
2	D	328	GLU
2	D	346	LYS
2	D	348	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	92	ASN
1	A	124	GLN
1	A	189	HIS
1	A	198	HIS
2	B	32	ASN
2	B	105	GLN
2	B	149	ASN
2	B	179	ASN
2	B	308	GLN
1	C	28	HIS
1	C	92	ASN
1	C	124	GLN
1	C	145	ASN
1	C	161	ASN
1	C	198	HIS
2	D	32	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	401	2	14,14,15	0.42	0	17,19,21	1.65	4 (23%)
3	GOL	A	301	-	5,5,5	0.08	0	5,5,5	0.32	0
3	GOL	A	302	-	5,5,5	0.07	0	5,5,5	0.26	0
3	GOL	C	301	-	5,5,5	0.12	0	5,5,5	0.39	0
4	NAG	D	401	2	14,14,15	0.35	0	17,19,21	0.38	0
3	GOL	B	402	-	5,5,5	0.07	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	401	2	-	3/6/23/26	0/1/1/1
3	GOL	A	301	-	-	1/4/4/4	-
3	GOL	A	302	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	301	-	-	0/4/4/4	-
4	NAG	D	401	2	-	0/6/23/26	0/1/1/1
3	GOL	B	402	-	-	2/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	NAG	C1-C2-N2	4.44	117.43	110.43
4	B	401	NAG	C2-N2-C7	3.45	127.53	122.90
4	B	401	NAG	C1-O5-C5	2.54	115.59	112.19
4	B	401	NAG	O5-C1-C2	-2.46	107.49	111.29

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	402	GOL	O1-C1-C2-C3
4	B	401	NAG	C1-C2-N2-C7
4	B	401	NAG	O5-C5-C6-O6
4	B	401	NAG	C4-C5-C6-O6
3	A	301	GOL	O1-C1-C2-C3
3	A	302	GOL	O1-C1-C2-C3
3	A	302	GOL	C1-C2-C3-O3
3	B	402	GOL	O1-C1-C2-O2
3	A	302	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	GOL	1	0
3	B	402	GOL	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/214 (98%)	0.26	2 (0%) 79 77	42, 66, 98, 118	0
1	C	208/214 (97%)	0.40	12 (5%) 29 26	40, 65, 111, 129	0
2	B	336/360 (93%)	0.35	14 (4%) 40 37	40, 64, 103, 136	0
2	D	322/360 (89%)	0.55	18 (5%) 30 27	43, 74, 111, 142	0
All	All	1076/1148 (93%)	0.40	46 (4%) 40 36	40, 67, 107, 142	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	213	LEU	4.8
1	A	29	ILE	4.0
2	B	224	PRO	3.5
2	D	10	GLY	3.4
2	B	351	ASP	3.2
2	B	222	ALA	3.1
2	B	64	LEU	3.0
2	D	108	THR	2.9
2	D	150	TYR	2.9
2	D	9	PRO	2.9
1	A	65	THR	2.8
2	D	85	SER	2.7
1	C	1	ALA	2.7
2	D	351	ASP	2.7
1	C	186	TYR	2.7
2	B	42	PRO	2.6
2	D	42	PRO	2.6
2	D	172	ILE	2.6
1	C	142	LYS	2.4
1	C	66	GLY	2.4
2	D	205	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	150	TYR	2.4
1	C	190	ASN	2.4
2	B	212	ALA	2.4
1	C	191	SER	2.4
1	C	189	HIS	2.4
2	D	86	VAL	2.3
2	D	222	ALA	2.3
1	C	126	THR	2.3
1	C	68	GLY	2.3
2	D	177	ALA	2.2
1	C	163	TRP	2.2
2	B	107	ASP	2.2
2	B	211	TRP	2.2
1	C	183	LYS	2.2
2	B	65	LYS	2.2
2	D	263	PRO	2.1
2	B	108	THR	2.1
2	B	226	GLY	2.1
2	D	245	THR	2.1
2	D	83	VAL	2.1
2	D	248	VAL	2.1
2	D	18	LEU	2.1
2	B	308	GLN	2.0
2	D	296	LEU	2.0
1	C	188	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	401	14/15	0.57	0.16	118,122,125,126	0
4	NAG	D	401	14/15	0.59	0.17	126,129,133,134	0
3	GOL	B	402	6/6	0.76	0.15	99,100,100,102	0
3	GOL	C	301	6/6	0.83	0.18	76,77,77,78	0
3	GOL	A	301	6/6	0.84	0.13	78,79,81,81	0
3	GOL	A	302	6/6	0.90	0.13	75,79,81,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.