



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 01:25 AM UTC

PDB ID : 6NSF / pdb_00006nsf
Title : Crystal structure of the A/Brisbane/10/2007 (H3N2) influenza virus hemagglutinin G186V/L194P mutant in complex with 3'-SLNLN
Authors : Wu, N.C.; Wilson, I.A.
Deposited on : 2019-01-24
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	18% 89% 9%
2	B	176	3% 94%
3	C	5	20% 80%
4	D	3	33% 67%
5	E	3	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	2	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 4541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	7	0
			2518	1578	446	482	12			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	PRO	-	expression tag	UNP C3PR70
A	10	GLY	-	expression tag	UNP C3PR70
A	190	ASP	ASN	conflict	UNP C3PR70
A	194	PRO	LEU	engineered mutation	UNP C3PR70

- Molecule 2 is a protein called Hemagglutinin HA2 chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	173	Total	C	N	O	S	0	6	0
			1426	894	247	279	6			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is N-acetyl-alpha-neuraminic acid (CCD ID: SIA) (formula: $C_{11}H_{19}NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			21	11	1	9		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	187	Total 187	O 187	0	0
9	B	181	Total 181	O 181	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemagglutinin HA1 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2
BGA3

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50%  50%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	100.17Å 100.17Å 382.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.10) 99.9 (50.00-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.182 , 0.214 0.189 , 0.220	Depositor DCC
R_{free} test set	2185 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4541	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, BMA, FUL, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2588	0.80	0/3522
2	B	0.74	0/1465	0.86	0/1969
All	All	0.72	0/4053	0.82	0/5491

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2468	13	0
2	B	1426	0	1375	9	0
3	C	61	0	52	0	0
4	D	38	0	34	0	0
5	E	39	0	34	0	0
6	F	28	0	25	0	0
7	A	28	0	26	1	0
7	B	14	0	13	0	0
8	A	21	0	18	0	0
9	A	187	0	0	0	0
9	B	181	0	0	2	0
All	All	4541	0	4045	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:60[B]:ASN:HD22	2:B:60[B]:ASN:H	1.08	0.98
2:B:60[B]:ASN:H	2:B:60[B]:ASN:ND2	1.70	0.86
2:B:15:GLU:OE2	9:B:301:HOH:O	2.07	0.73
2:B:60[B]:ASN:HD22	2:B:60[B]:ASN:N	1.88	0.67
1:A:141:ARG:NH1	1:A:147:PHE:O	2.38	0.56
1:A:193:PHE:HB3	1:A:194:PRO:HD3	1.90	0.54
1:A:50[A]:GLU:HG2	1:A:273:PRO:HG2	1.91	0.51
1:A:188:ASP:HA	1:A:191:GLN:HB3	1.95	0.48
1:A:264:LYS:HD2	2:B:63:PHE:CE2	2.49	0.48
1:A:141:ARG:HB3	1:A:146:SER:CB	2.45	0.46
1:A:24:THR:OG1	7:A:401:NAG:H81	2.17	0.45
1:A:169:PRO:HA	1:A:242:ILE:HA	2.00	0.44
1:A:141:ARG:HB3	1:A:146:SER:HB2	2.01	0.43
1:A:111:LEU:C	1:A:111:LEU:HD12	2.44	0.42
2:B:121:LYS:NZ	9:B:308:HOH:O	2.52	0.41
1:A:166:VAL:O	1:A:244:LEU:HD12	2.20	0.41
2:B:110:LEU:C	2:B:110:LEU:HD12	2.46	0.41
1:A:264:LYS:HB3	2:B:63:PHE:CG	2.56	0.41
2:B:115:MET:HA	2:B:115:MET:HE3	2.03	0.41
1:A:102:VAL:HG22	1:A:232:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	322/321 (100%)	308 (96%)	14 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	177/176 (101%)	168 (95%)	9 (5%)	0	100	100
All	All	499/497 (100%)	476 (95%)	23 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/285 (101%)	275 (96%)	13 (4%)	24	25
2	B	152/148 (103%)	148 (97%)	4 (3%)	40	46
All	All	440/433 (102%)	423 (96%)	17 (4%)	32	31

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	128	THR
1	A	142	ARG
1	A	143	SER
1	A	144	ASN
1	A	167	THR
1	A	187	THR
1	A	201	ARG
1	A	236	ILE
1	A	246	ASN
1	A	276	LYS
1	A	296[A]	ASN
1	A	296[B]	ASN
1	A	325	GLU
2	B	38	LEU
2	B	60[A]	ASN
2	B	60[B]	ASN
2	B	168	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	GLN
1	A	211	GLN
1	A	246	ASN
1	A	250	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,1	14,14,15	0.48	0	17,19,21	1.31	1 (5%)
3	NAG	C	2	3	14,14,15	0.33	0	17,19,21	0.83	1 (5%)
3	BMA	C	3	3	11,11,12	0.35	0	15,15,17	0.56	0
3	MAN	C	4	3	11,11,12	0.40	0	15,15,17	0.86	1 (6%)
3	MAN	C	5	3	11,11,12	0.69	0	15,15,17	1.90	2 (13%)
4	NAG	D	1	1,4	14,14,15	0.40	0	17,19,21	0.91	0
4	NAG	D	2	4	14,14,15	0.30	0	17,19,21	1.00	1 (5%)
4	FUL	D	3	4	10,10,11	0.49	0	14,14,16	0.97	1 (7%)
5	NAG	E	1	1,5	14,14,15	0.43	0	17,19,21	1.09	1 (5%)
5	NAG	E	2	5	14,14,15	0.63	0	17,19,21	1.30	3 (17%)
5	BMA	E	3	5	11,11,12	0.73	0	15,15,17	1.90	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	F	1	1,6	14,14,15	0.53	0	17,19,21	0.99	2 (11%)
6	NAG	F	2	6	14,14,15	0.27	0	17,19,21	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
4	FUL	D	3	4	-	-	0/1/1/1
5	NAG	E	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
6	NAG	F	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	F	2	6	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	MAN	C1-C2-C3	5.13	117.11	109.64
5	E	3	BMA	C1-O5-C5	4.85	118.69	112.19
5	E	3	BMA	C1-C2-C3	4.64	116.40	109.64
3	C	5	MAN	C1-O5-C5	4.56	118.30	112.19
3	C	1	NAG	C1-O5-C5	4.49	118.21	112.19
5	E	1	NAG	C1-O5-C5	3.60	117.01	112.19
5	E	2	NAG	C1-O5-C5	2.74	115.86	112.19
4	D	2	NAG	C4-C3-C2	2.54	114.74	111.02
3	C	4	MAN	C1-O5-C5	2.47	115.50	112.19
4	D	3	FUL	C1-C2-C3	2.35	113.06	109.64
3	C	2	NAG	C1-O5-C5	2.32	115.29	112.19
6	F	1	NAG	O5-C5-C4	-2.22	105.42	110.83
6	F	1	NAG	C1-O5-C5	-2.15	109.30	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	2	NAG	C2-N2-C7	2.15	125.78	122.90
5	E	2	NAG	C3-C4-C5	-2.01	106.59	110.23

There are no chirality outliers.


All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1	NAG	O5-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6
3	C	3	BMA	O5-C5-C6-O6
3	C	4	MAN	C4-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
5	E	2	NAG	O5-C5-C6-O6
5	E	2	NAG	C4-C5-C6-O6
5	E	3	BMA	C4-C5-C6-O6
5	E	3	BMA	O5-C5-C6-O6
6	F	2	NAG	C4-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
6	F	2	NAG	O5-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	A	401	1	14,14,15	0.61	0	17,19,21	1.14	1 (5%)
7	NAG	B	201	2	14,14,15	0.33	0	17,19,21	0.62	0
8	SIA	A	416	-	21,21,21	0.93	1 (4%)	24,31,31	1.05	2 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	A	415	1	14,14,15	0.45	0	17,19,21	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	401	1	-	2/6/23/26	0/1/1/1
7	NAG	B	201	2	-	2/6/23/26	0/1/1/1
8	SIA	A	416	-	-	5/20/38/38	0/1/1/1
7	NAG	A	415	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	416	SIA	O2-C2	3.09	1.44	1.39

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	401	NAG	C1-O5-C5	2.94	116.13	112.19
8	A	416	SIA	O1A-C1-C2	-2.92	118.98	123.85
8	A	416	SIA	O6-C6-C5	2.63	112.25	109.84

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	A	401	NAG	O5-C5-C6-O6
7	A	401	NAG	C4-C5-C6-O6
7	B	201	NAG	O5-C5-C6-O6
8	A	416	SIA	C7-C8-C9-O9
8	A	416	SIA	O8-C8-C9-O9
8	A	416	SIA	O1B-C1-C2-C3
8	A	416	SIA	O1A-C1-C2-O2
8	A	416	SIA	O1A-C1-C2-C3
7	B	201	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	401	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/321 (98%)	0.72	57 (17%) 3 3	13, 46, 144, 182	7 (2%)
2	B	173/176 (98%)	-0.22	5 (2%) 53 56	15, 31, 48, 91	6 (3%)
All	All	490/497 (98%)	0.39	62 (12%) 8 8	13, 37, 119, 182	13 (2%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	193	PHE	6.4
1	A	214	ILE	5.7
1	A	194	PRO	5.5
1	A	157	LEU	5.3
2	B	173	ILE	5.3
1	A	195	TYR	5.2
1	A	156	HIS	5.1
1	A	213	VAL	5.1
1	A	198	ALA	4.9
1	A	196	ALA	4.8
1	A	186	VAL	4.6
1	A	217	ILE	4.5
1	A	159	PHE	4.3
1	A	192	ILE	4.2
1	A	212	THR	4.2
2	B	169	ASN	4.0
1	A	129	GLY	3.9
2	B	170	ARG	3.7
1	A	202	ILE	3.7
1	A	162	PRO	3.7
1	A	250	ASN	3.6
1	A	223	VAL	3.5
1	A	221	PRO	3.5
1	A	200	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	161	TYR	3.5
1	A	163	ALA	3.4
2	B	171	PHE	3.4
1	A	9	PRO	3.4
1	A	154	LEU	3.4
1	A	204	VAL	3.4
1	A	226	ILE	3.3
1	A	187	THR	3.2
1	A	220	ARG	3.1
1	A	251	LEU	3.1
1	A	190	ASP	3.1
1	A	218	GLY	3.0
1	A	246	ASN	3.0
1	A	155	THR	2.8
1	A	199	SER	2.8
1	A	219	SER	2.8
1	A	191	GLN	2.8
1	A	143	SER	2.7
1	A	128	THR	2.6
1	A	206	THR	2.6
1	A	160	LYS	2.6
1	A	130	VAL	2.5
1	A	158	LYS	2.5
1	A	201	ARG	2.5
1	A	164	LEU	2.5
1	A	166	VAL	2.4
1	A	205	SER	2.4
1	A	243	LEU	2.4
1	A	216	ASN	2.4
1	A	188	ASP	2.3
1	A	215	PRO	2.3
1	A	127	TRP	2.3
1	A	167	THR	2.2
1	A	237	VAL	2.1
2	B	172	GLN	2.1
1	A	248	THR	2.1
1	A	197	GLN	2.0
1	A	131	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	BMA	E	3	11/12	0.28	0.15	136,140,142,142	0
3	MAN	C	4	11/12	0.46	0.15	103,107,114,114	0
4	FUL	D	3	10/11	0.57	0.16	92,95,97,98	0
3	BMA	C	3	11/12	0.58	0.13	92,98,101,105	0
4	NAG	D	2	14/15	0.58	0.14	88,99,103,105	0
5	NAG	E	1	14/15	0.59	0.15	93,101,104,110	0
6	NAG	F	2	14/15	0.69	0.13	72,78,81,84	0
5	NAG	E	2	14/15	0.75	0.14	116,119,122,128	0
6	NAG	F	1	14/15	0.80	0.13	42,50,54,64	0
3	MAN	C	5	11/12	0.80	0.14	96,101,102,102	0
3	NAG	C	1	14/15	0.81	0.13	47,51,53,60	0
4	NAG	D	1	14/15	0.84	0.12	58,66,85,86	0
3	NAG	C	2	14/15	0.89	0.11	65,71,74,83	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	A	401	14/15	0.55	0.16	65,72,76,77	0
8	SIA	A	416	21/21	0.62	0.17	96,104,106,106	0
7	NAG	B	201	14/15	0.65	0.15	72,81,82,85	0
7	NAG	A	415	14/15	0.74	0.14	77,87,90,92	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SIA A 416:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.