



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 08:33 AM UTC

PDB ID : 6NSY / pdb_00006nsy
Title : X-ray reduced Catalase 3 From N.Crassa in Cpd I state (0.263 MGy)
Authors : Zarate-Romero, A.; Rudino-Pinera, E.; Stojanoff, V.
Deposited on : 2019-01-27
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	719	 87% 8% 5%
1	B	719	 85% 9% 6%
1	C	719	 86% 8% 6%
1	D	719	 88% 6% 6%

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 23858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	681	Total	C	N	O	S	0	6	0
			5393	3408	956	1023	6			
1	B	677	Total	C	N	O	S	0	6	0
			5359	3392	951	1010	6			
1	C	679	Total	C	N	O	S	0	9	0
			5394	3411	957	1020	6			
1	D	678	Total	C	N	O	S	0	9	0
			5385	3408	956	1015	6			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		
4	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	C	1	Total	C	O	0	0
			4	2	2		
6	C	1	Total	C	O	0	0
			4	2	2		
6	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	C	O	0	0
			4	2	2		
6	D	1	Total	C	O	0	0
			4	2	2		
6	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is ETHANOL (CCD ID: EOH) (formula: C_2H_6O).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			3	2	1		
7	B	1	Total	C	O	0	0
			3	2	1		
7	B	1	Total	C	O	0	0
			3	2	1		
7	D	1	Total	C	O	0	0
			3	2	1		
7	D	1	Total	C	O	0	0
			3	2	1		
7	D	1	Total	C	O	0	0
			3	2	1		

- Molecule 8 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			16	10	6		

- Molecule 9 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	C	1	Total	C	O	0	0
			13	8	5		

- Molecule 10 is OXYGEN ATOM (CCD ID: O) (formula: O) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	C	1	Total 1	O 1	0	0
10	D	1	Total 1	O 1	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	485	Total 485	O 485	0	0
11	B	493	Total 493	O 493	0	0
11	C	503	Total 503	O 503	0	0
11	D	485	Total 485	O 485	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Catalase-3



• Molecule 1: Catalase-3



• Molecule 1: Catalase-3





● Molecule 1: Catalase-3

Chain D: 88% 6% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	131.10Å 154.50Å 160.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.60 – 2.20 29.60 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.7 (29.60-2.20) 95.6 (29.60-2.20)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.12 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.12 _2829	Depositor
R, R_{free}	0.154 , 0.202 0.155 , 0.203	Depositor DCC
R_{free} test set	7944 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	23858	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, EOH, EDO, PEG, HEM, 1PE, PGE, PG4, O

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.20	0/5524	0.44	0/7488
1	B	0.21	0/5495	0.45	0/7452
1	C	0.21	0/5527	0.43	0/7494
1	D	0.20	0/5521	0.44	0/7485
All	All	0.21	0/22067	0.44	0/29919

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5393	0	5201	50	0
1	B	5359	0	5187	47	0
1	C	5394	0	5210	49	0
1	D	5385	0	5213	42	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0
2	C	43	0	30	1	0
2	D	43	0	30	1	0
3	A	8	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	8	0	6	0	0
3	C	8	0	6	2	0
3	D	8	0	6	2	0
4	A	10	0	14	0	0
4	B	10	0	14	0	0
4	D	10	0	14	1	0
5	A	14	0	20	1	0
5	B	7	0	10	0	0
5	C	21	0	30	0	0
6	A	4	0	6	0	0
6	B	8	0	12	1	0
6	C	12	0	18	2	0
6	D	12	0	18	1	0
7	A	3	0	6	0	0
7	B	6	0	12	0	0
7	D	9	0	18	0	0
8	B	16	0	22	5	0
9	C	13	0	18	6	0
10	C	1	0	0	0	0
10	D	1	0	0	0	0
11	A	485	0	0	0	0
11	B	493	0	0	0	0
11	C	503	0	0	5	0
11	D	485	0	0	0	0
All	All	23858	0	21187	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:717:ASP:HA	1:D:638:LYS:HZ3	1.19	1.02
1:A:717:ASP:HB2	1:D:638:LYS:HE2	1.53	0.89
1:A:717:ASP:HA	1:D:638:LYS:NZ	1.88	0.87
1:C:342:GLN:HG2	9:C:804:PG4:H32	1.65	0.77
1:A:717:ASP:HB2	1:D:638:LYS:CE	2.19	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	683/719 (95%)	661 (97%)	22 (3%)	0	100	100
1	B	681/719 (95%)	661 (97%)	20 (3%)	0	100	100
1	C	686/719 (95%)	661 (96%)	25 (4%)	0	100	100
1	D	685/719 (95%)	667 (97%)	18 (3%)	0	100	100
All	All	2735/2876 (95%)	2650 (97%)	85 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	564/585 (96%)	559 (99%)	5 (1%)	70	84
1	B	561/585 (96%)	557 (99%)	4 (1%)	76	87
1	C	565/585 (97%)	562 (100%)	3 (0%)	81	90
1	D	564/585 (96%)	561 (100%)	3 (0%)	81	90
All	All	2254/2340 (96%)	2239 (99%)	15 (1%)	76	87

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	582	LEU
1	D	315	ILE
1	B	592	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	709	GLU
1	C	706	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	395	ASN
1	C	550	ASN
1	D	512	GLN
1	C	542	GLN
1	C	551	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 40 ligands modelled in this entry, 2 are monoatomic - leaving 38 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	D	805	-	3,3,3	0.46	0	2,2,2	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	C	801	10,1	50,50,50	1.43	8 (16%)	67,82,82	1.16	6 (8%)
7	EOH	D	809	-	2,2,2	0.47	0	1,1,1	0.15	0
6	EDO	D	806	-	3,3,3	0.42	0	2,2,2	0.45	0
4	PGE	A	804	-	9,9,9	0.34	0	8,8,8	0.30	0
7	EOH	A	808	-	2,2,2	0.47	0	1,1,1	0.11	0
3	ACT	C	802	-	3,3,3	1.58	1 (33%)	3,3,3	1.33	0
3	ACT	D	802	-	3,3,3	1.36	0	3,3,3	1.45	0
3	ACT	A	803	-	3,3,3	1.82	1 (33%)	3,3,3	1.20	0
6	EDO	D	807	-	3,3,3	0.46	0	2,2,2	0.39	0
6	EDO	C	808	-	3,3,3	0.44	0	2,2,2	0.28	0
6	EDO	B	807	-	3,3,3	0.43	0	2,2,2	0.33	0
5	PEG	C	805	-	6,6,6	0.49	0	5,5,5	0.43	0
6	EDO	C	810	-	3,3,3	0.42	0	2,2,2	0.39	0
6	EDO	C	809	-	3,3,3	0.44	0	2,2,2	0.47	0
3	ACT	C	803	-	3,3,3	1.56	1 (33%)	3,3,3	1.53	0
5	PEG	C	806	-	6,6,6	0.48	0	5,5,5	0.45	0
2	HEM	D	801	10,1	50,50,50	1.48	10 (20%)	67,82,82	1.15	5 (7%)
3	ACT	A	802	-	3,3,3	1.54	1 (33%)	3,3,3	1.36	0
5	PEG	C	807	-	6,6,6	0.50	0	5,5,5	0.45	0
4	PGE	D	804	-	9,9,9	0.38	0	8,8,8	0.24	0
7	EOH	D	810	-	2,2,2	0.46	0	1,1,1	0.13	0
7	EOH	D	811	-	2,2,2	0.42	0	1,1,1	0.21	0
2	HEM	B	801	11,1	50,50,50	1.42	9 (18%)	67,82,82	1.15	5 (7%)
3	ACT	B	803	-	3,3,3	1.43	1 (33%)	3,3,3	1.22	0
5	PEG	B	806	-	6,6,6	0.46	0	5,5,5	0.62	0
2	HEM	A	801	11,1	50,50,50	1.45	9 (18%)	67,82,82	1.15	5 (7%)
9	PG4	C	804	-	12,12,12	0.50	0	11,11,11	0.38	0
7	EOH	B	810	-	2,2,2	0.45	0	1,1,1	0.20	0
4	PGE	B	805	-	9,9,9	0.30	0	8,8,8	0.37	0
6	EDO	A	807	-	3,3,3	0.41	0	2,2,2	0.44	0
7	EOH	B	809	-	2,2,2	0.43	0	1,1,1	0.24	0
3	ACT	B	802	-	3,3,3	1.33	0	3,3,3	1.33	0
8	1PE	B	804	-	15,15,15	0.52	0	14,14,14	0.45	0
5	PEG	A	806	-	6,6,6	0.48	0	5,5,5	0.58	0
6	EDO	B	808	-	3,3,3	0.45	0	2,2,2	0.33	0
5	PEG	A	805	-	6,6,6	0.49	0	5,5,5	0.56	0
3	ACT	D	803	-	3,3,3	1.64	1 (33%)	3,3,3	1.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	D	805	-	-	0/1/1/1	-
2	HEM	C	801	10,1	-	2/14/54/54	-
6	EDO	D	806	-	-	1/1/1/1	-
4	PGE	A	804	-	-	5/7/7/7	-
6	EDO	D	807	-	-	1/1/1/1	-
6	EDO	C	808	-	-	0/1/1/1	-
6	EDO	B	807	-	-	1/1/1/1	-
5	PEG	C	805	-	-	3/4/4/4	-
6	EDO	C	810	-	-	0/1/1/1	-
6	EDO	C	809	-	-	1/1/1/1	-
5	PEG	C	806	-	-	2/4/4/4	-
2	HEM	D	801	10,1	-	2/14/54/54	-
5	PEG	C	807	-	-	0/4/4/4	-
4	PGE	D	804	-	-	4/7/7/7	-
2	HEM	B	801	11,1	-	4/14/54/54	-
5	PEG	B	806	-	-	1/4/4/4	-
2	HEM	A	801	11,1	-	2/14/54/54	-
9	PG4	C	804	-	-	4/10/10/10	-
6	EDO	A	807	-	-	0/1/1/1	-
4	PGE	B	805	-	-	5/7/7/7	-
8	1PE	B	804	-	-	8/13/13/13	-
5	PEG	A	806	-	-	4/4/4/4	-
6	EDO	B	808	-	-	0/1/1/1	-
5	PEG	A	805	-	-	2/4/4/4	-

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	FE-ND	4.37	2.08	1.94
2	B	801	HEM	FE-ND	4.29	2.08	1.94
2	D	801	HEM	FE-ND	3.99	2.07	1.94
2	D	801	HEM	FE-NB	3.95	2.07	1.94
2	C	801	HEM	FE-NB	3.77	2.06	1.94

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	CAA-CBA-CGA	-3.18	105.24	113.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	801	HEM	CAA-CBA-CGA	-2.79	106.27	113.67
2	D	801	HEM	C4D-ND-C1D	2.77	108.48	105.21
2	A	801	HEM	CAA-CBA-CGA	-2.74	106.39	113.67
2	A	801	HEM	C2A-C1A-NA	-2.61	107.25	110.15

There are no chirality outliers.

5 of 52 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	804	PGE	C3-C4-O3-C5
4	A	804	PGE	O2-C3-C4-O3
9	C	804	PG4	C1-C2-O2-C3
4	D	804	PGE	O3-C5-C6-O4
9	C	804	PG4	O1-C1-C2-O2


There are no ring outliers.


14 monomers are involved in 23 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	801	HEM	1	0
3	C	802	ACT	1	0
3	D	802	ACT	1	0
6	D	807	EDO	1	0
6	C	808	EDO	1	0
6	C	810	EDO	1	0
3	C	803	ACT	1	0
2	D	801	HEM	1	0
4	D	804	PGE	1	0
9	C	804	PG4	6	0
8	B	804	1PE	5	0
5	A	806	PEG	1	0
6	B	808	EDO	1	0
3	D	803	ACT	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	681/719 (94%)	-0.54	1 (0%) 92 90	4, 14, 34, 61	6 (0%)
1	B	677/719 (94%)	-0.60	0 100 100	4, 13, 30, 53	6 (0%)
1	C	679/719 (94%)	-0.60	4 (0%) 85 83	3, 13, 31, 61	9 (1%)
1	D	678/719 (94%)	-0.55	3 (0%) 88 87	4, 15, 32, 59	9 (1%)
All	All	2715/2876 (94%)	-0.57	8 (0%) 90 88	3, 14, 32, 61	30 (1%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	716	ASP	3.5
1	A	540	ALA	3.4
1	C	715	GLY	3.2
1	C	540	ALA	3.2
1	D	681	GLU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	C	809	4/4	0.78	0.15	29,29,30,32	0
3	ACT	D	803	4/4	0.79	0.17	20,21,23,28	0
7	EOH	B	809	3/3	0.79	0.14	23,23,26,29	0
6	EDO	B	808	4/4	0.80	0.15	32,32,32,34	0
6	EDO	D	805	4/4	0.81	0.15	31,32,32,34	0
3	ACT	C	803	4/4	0.81	0.13	20,21,22,29	0
3	ACT	A	803	4/4	0.82	0.14	18,18,20,22	0
4	PGE	D	804	10/10	0.83	0.14	28,30,31,31	0
7	EOH	D	811	3/3	0.83	0.12	21,21,23,24	0
6	EDO	C	808	4/4	0.84	0.13	30,31,32,32	0
8	1PE	B	804	16/16	0.84	0.12	29,30,33,34	0
7	EOH	D	809	3/3	0.86	0.12	28,28,31,36	0
7	EOH	D	810	3/3	0.86	0.11	25,25,27,29	0
6	EDO	D	807	4/4	0.86	0.13	29,29,29,30	0
4	PGE	A	804	10/10	0.86	0.14	24,27,29,31	0
10	O	C	811	1/1	0.86	0.22	8,8,8,8	1
5	PEG	B	806	7/7	0.87	0.18	24,25,27,27	0
10	O	D	808	1/1	0.87	0.17	9,9,9,9	1
5	PEG	A	805	7/7	0.88	0.10	26,26,27,28	0
5	PEG	C	806	7/7	0.88	0.12	27,27,29,31	0
5	PEG	A	806	7/7	0.89	0.12	21,23,24,27	0
9	PG4	C	804	13/13	0.89	0.10	26,27,28,29	0
6	EDO	B	807	4/4	0.89	0.08	27,27,28,28	0
7	EOH	B	810	3/3	0.89	0.10	23,23,23,26	0
4	PGE	B	805	10/10	0.90	0.12	24,26,28,29	0
3	ACT	B	803	4/4	0.91	0.08	15,16,16,18	0
5	PEG	C	807	7/7	0.92	0.08	25,27,29,31	0
5	PEG	C	805	7/7	0.92	0.09	30,30,31,31	0
6	EDO	D	806	4/4	0.93	0.09	29,31,32,33	0
6	EDO	C	810	4/4	0.93	0.07	27,28,28,31	0
7	EOH	A	808	3/3	0.93	0.08	22,22,23,25	0
6	EDO	A	807	4/4	0.94	0.08	27,29,29,29	0
3	ACT	D	802	4/4	0.98	0.04	11,11,12,13	0
2	HEM	A	801	43/43	0.98	0.05	3,7,12,15	0
3	ACT	B	802	4/4	0.98	0.05	9,9,9,9	0
2	HEM	B	801	43/43	0.98	0.06	2,7,11,17	0
3	ACT	C	802	4/4	0.98	0.05	12,13,14,14	0
3	ACT	A	802	4/4	0.98	0.05	11,12,12,13	0
2	HEM	C	801	43/43	0.99	0.05	2,7,10,14	0
2	HEM	D	801	43/43	0.99	0.05	2,7,13,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.




Electron density around O C 811:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around O D 808:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM B 801:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.