



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 08:50 PM EDT

PDB ID : 3NT1 / pdb_00003nt1
Title : High resolution structure of naproxen:COX-2 complex.
Authors : Duggan, K.C.; Musee, J.; Walters, M.J.; Harp, J.M.; Kiefer, J.R.; Oates, J.A.;
Marnett, L.J.
Deposited on : 2010-07-02
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	4% 89% 6%
1	B	587	4% 88% 5% 6%
2	C	2	50% 50%
2	D	2	50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	661	X	-	-	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 10338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin-endoperoxide synthase 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	3	0
			4499	2899	756	819	25			
1	B	552	Total	C	N	O	S	0	0	0
			4474	2885	750	814	25			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is octyl beta-D-glucopyranoside (CCD ID: BOG) (formula: $C_{14}H_{28}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			20	14	6		
4	A	1	Total	C	O	0	0
			20	14	6		
4	A	1	Total	C	O	0	0
			20	14	6		
4	B	1	Total	C	O	0	0
			20	14	6		

- Molecule 5 is (2S)-2-(6-methoxynaphthalen-2-yl)propanoic acid (CCD ID: NPS) (formula: $C_{14}H_{14}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			17	14	3		
5	B	1	Total	C	O	0	0
			17	14	3		

- Molecule 6 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Prostaglandin-endoperoxide synthase 2



- Molecule 1: Prostaglandin-endoperoxide synthase 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	122.30Å 133.23Å 181.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.73 30.00 – 1.73	Depositor EDS
% Data completeness (in resolution range)	98.2 (30.00-1.73) 98.1 (30.00-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.167 , 0.186 0.182 , 0.201	Depositor DCC
R_{free} test set	7592 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10338	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BOG, NAG, NPS, HEM, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/4629	0.68	0/6276
1	B	0.45	0/4601	0.68	0/6239
All	All	0.45	0/9230	0.68	0/12515

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4499	0	4396	20	0
1	B	4474	0	4373	21	0
2	C	28	0	25	1	0
2	D	28	0	25	1	0
3	A	28	0	26	0	0
3	B	28	0	26	0	0
4	A	60	0	84	3	0
4	B	20	0	28	0	0
5	A	17	0	13	0	0
5	B	17	0	13	0	0
6	A	43	0	30	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	43	0	30	1	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
8	A	527	0	0	0	0
8	B	524	0	0	1	0
All	All	10338	0	9069	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:LEU:HD21	1:A:415:LEU:HD12	1.48	0.95
1:A:87:ASN:HD22	4:A:620:BOG:H62	1.37	0.89
1:A:581[B]:ASN:HD22	1:A:582:VAL:N	1.88	0.72
1:A:283:LEU:CD2	1:A:415:LEU:HD12	2.21	0.69
1:B:382:ASN:HD21	6:B:619:HEM:HAD2	1.57	0.68
1:A:271:VAL:HG22	1:A:286:ALA:HB1	1.78	0.65
1:B:81:LEU:HD22	1:B:82:LEU:HD12	1.78	0.65
1:A:271:VAL:CG2	1:A:286:ALA:HB1	2.32	0.59
1:B:375:ASN:ND2	8:B:832:HOH:O	2.21	0.58
1:A:87:ASN:HD22	4:A:620:BOG:C6	2.12	0.58
1:B:274:ILE:HD12	1:B:291:VAL:HG12	1.88	0.55
1:B:581:ASN:HD22	1:B:582:VAL:N	2.07	0.52
1:A:120:ARG:HG3	1:A:531:LEU:HD12	1.91	0.52
1:A:500:VAL:O	1:A:500:VAL:HG12	2.09	0.51
1:B:196:MET:CE	1:B:431:ALA:HB2	2.41	0.51
1:B:382:ASN:O	1:B:386:HIS:HD2	1.94	0.51
1:A:87:ASN:ND2	4:A:620:BOG:H62	2.16	0.50
1:A:270:GLN:HE21	1:A:270:GLN:N	2.09	0.50
1:B:180:LYS:HD2	1:B:494:LEU:HD11	1.94	0.49
1:B:167:GLY:HA3	1:B:500:VAL:HG21	1.96	0.47
1:B:527:ALA:HB3	1:B:528:PRO:HD3	1.96	0.47
1:A:500:VAL:O	1:A:500:VAL:CG1	2.63	0.47
1:B:194:SER:O	1:B:430:ILE:HD12	2.15	0.47
1:A:527:ALA:HB3	1:A:528:PRO:HD3	1.97	0.46
1:B:196:MET:HE2	1:B:196:MET:HA	1.96	0.46
1:B:382:ASN:C	1:B:382:ASN:HD22	2.24	0.46
1:A:216:ARG:NH1	2:C:2:NAG:O7	2.49	0.46
1:A:382:ASN:O	1:A:386:HIS:HD2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:THR:HB	1:A:271:VAL:HG13	1.99	0.45
1:A:388:HIS:N	1:A:389:PRO:CD	2.81	0.43
1:A:581[B]:ASN:HD22	1:A:581[B]:ASN:C	2.27	0.42
1:B:78:ILE:O	1:B:82:LEU:HD13	2.19	0.42
1:B:404:PHE:O	1:B:408:LEU:HD13	2.20	0.42
1:B:388:HIS:N	1:B:389:PRO:CD	2.83	0.41
1:B:463:LEU:HD22	1:B:506:ALA:CB	2.50	0.41
1:B:167:GLY:CA	1:B:500:VAL:HG21	2.51	0.41
1:A:33:ALA:HB3	1:A:158:ASP:CG	2.46	0.41
1:B:81:LEU:C	1:B:81:LEU:HD23	2.45	0.41
1:A:384:LEU:HD12	1:A:384:LEU:C	2.46	0.41
1:B:216:ARG:NH1	2:D:1:NAG:O4	2.54	0.41
1:B:384:LEU:HD12	1:B:384:LEU:C	2.46	0.41
6:A:619:HEM:HHC	6:A:619:HEM:HBB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/587 (94%)	542 (98%)	11 (2%)	0	100	100
1	B	550/587 (94%)	539 (98%)	11 (2%)	0	100	100
All	All	1103/1174 (94%)	1081 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	496/525 (94%)	488 (98%)	8 (2%)	55	35
1	B	493/525 (94%)	487 (99%)	6 (1%)	63	47
All	All	989/1050 (94%)	975 (99%)	14 (1%)	59	41

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	LEU
1	A	101	ASN
1	A	267	LYS
1	A	270	GLN
1	A	271	VAL
1	A	289	GLN
1	A	385	TYR
1	A	484	GLU
1	B	289	GLN
1	B	385	TYR
1	B	405	LYS
1	B	473	LYS
1	B	484	GLU
1	B	581	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	ASN
1	A	270	GLN
1	A	282	ASN
1	A	327	GLN
1	A	330	GLN
1	A	571	ASN
1	B	54	GLN
1	B	101	ASN
1	B	282	ASN
1	B	369	GLN
1	B	382	ASN
1	B	388	HIS
1	B	571	ASN
1	B	581	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	C	1	1,2	14,14,15	0.54	0	17,19,21	0.91	1 (5%)
2	NAG	C	2	2	14,14,15	0.49	0	17,19,21	0.95	1 (5%)
2	NAG	D	1	1,2	14,14,15	0.56	0	17,19,21	0.85	1 (5%)
2	NAG	D	2	2	14,14,15	0.48	0	17,19,21	1.36	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	NAG	C2-N2-C7	3.32	127.34	122.90
2	C	1	NAG	C1-O5-C5	2.74	115.86	112.19
2	C	2	NAG	C4-C3-C2	2.28	114.36	111.02
2	D	2	NAG	C8-C7-N2	2.28	119.89	116.12
2	D	1	NAG	C1-O5-C5	2.24	115.19	112.19
2	D	2	NAG	C1-C2-N2	2.12	113.77	110.43
2	D	2	NAG	O5-C1-C2	-2.08	108.08	111.29

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	NAG	C1-C2-N2-C7
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	1	0
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	681	1	14,14,15	0.53	0	17,19,21	0.77	0
4	BOG	B	3	-	20,20,20	0.76	1 (5%)	25,25,25	0.86	2 (8%)
3	NAG	A	661	1	14,14,15	0.76	0	17,19,21	1.09	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	HEM	B	619	8,1	50,50,50	1.84	9 (18%)	67,82,82	1.43	7 (10%)
3	NAG	A	681	1	14,14,15	0.48	0	17,19,21	0.99	1 (5%)
5	NPS	B	4	-	18,18,18	1.10	1 (5%)	25,25,25	0.69	0
4	BOG	A	6	-	20,20,20	0.72	1 (5%)	25,25,25	0.82	1 (4%)
4	BOG	A	620	-	20,20,20	0.82	1 (5%)	25,25,25	1.62	4 (16%)
4	BOG	A	3	-	20,20,20	0.62	1 (5%)	25,25,25	0.88	1 (4%)
3	NAG	B	9	1	14,14,15	0.56	0	17,19,21	0.89	0
6	HEM	A	619	8,1	50,50,50	1.83	10 (20%)	67,82,82	1.41	8 (11%)
5	NPS	A	5	-	18,18,18	1.14	1 (5%)	25,25,25	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	681	1	-	0/6/23/26	0/1/1/1
4	BOG	B	3	-	-	7/11/31/31	0/1/1/1
3	NAG	A	661	1	1/1/5/7	5/6/23/26	0/1/1/1
6	HEM	B	619	8,1	-	2/14/54/54	-
3	NAG	A	681	1	-	0/6/23/26	0/1/1/1
5	NPS	B	4	-	-	1/10/10/10	0/2/2/2
4	BOG	A	6	-	-	3/11/31/31	0/1/1/1
4	BOG	A	620	-	-	5/11/31/31	0/1/1/1
4	BOG	A	3	-	-	5/11/31/31	0/1/1/1
3	NAG	B	9	1	-	2/6/23/26	0/1/1/1
6	HEM	A	619	8,1	-	0/14/54/54	-
5	NPS	A	5	-	-	0/10/10/10	0/2/2/2

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	619	HEM	C3D-C2D	7.57	1.53	1.36
6	B	619	HEM	C3D-C2D	7.52	1.53	1.36
6	B	619	HEM	FE-ND	4.28	2.08	1.94
6	A	619	HEM	FE-ND	4.18	2.07	1.94
6	B	619	HEM	FE-NB	4.10	2.07	1.94
6	A	619	HEM	FE-NB	3.97	2.07	1.94
6	A	619	HEM	FE-NA	3.71	2.07	1.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	619	HEM	FE-NA	3.36	2.06	1.95
4	A	620	BOG	O1-C1	2.96	1.45	1.40
6	B	619	HEM	FE-NC	2.93	2.04	1.95
6	B	619	HEM	CAC-C3C	2.92	1.55	1.47
6	A	619	HEM	CAC-C3C	2.88	1.55	1.47
6	B	619	HEM	CAB-C3B	2.73	1.54	1.47
6	A	619	HEM	CAB-C3B	2.63	1.54	1.47
6	B	619	HEM	CMC-C2C	2.21	1.55	1.50
4	A	6	BOG	O1-C1	2.21	1.43	1.40
4	B	3	BOG	O1-C1	2.15	1.43	1.40
6	A	619	HEM	CMA-C3A	2.11	1.55	1.50
6	A	619	HEM	CMD-C2D	2.05	1.55	1.50
5	A	5	NPS	C6-C1	2.05	1.40	1.37
4	A	3	BOG	O1-C1	2.05	1.43	1.40
5	B	4	NPS	C6-C1	2.04	1.40	1.37
6	A	619	HEM	CMB-C2B	2.02	1.54	1.50
6	B	619	HEM	CMB-C2B	2.01	1.54	1.50
6	A	619	HEM	C2A-C3A	-2.00	1.33	1.38

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	619	HEM	C4D-ND-C1D	5.63	111.88	105.21
6	A	619	HEM	C4D-ND-C1D	5.35	111.54	105.21
4	A	620	BOG	O1-C1-C2	4.88	115.68	108.27
4	A	620	BOG	O5-C1-O1	-4.20	100.12	110.04
6	B	619	HEM	C3B-C4B-NB	-3.14	107.21	109.47
6	A	619	HEM	C3B-C4B-NB	-2.98	107.33	109.47
4	A	620	BOG	O5-C5-C4	2.93	114.98	109.70
4	A	3	BOG	C1'-O1-C1	-2.91	108.71	113.68
6	B	619	HEM	C1B-NB-C4B	2.85	108.58	105.21
6	B	619	HEM	CHA-C4D-ND	2.67	127.67	124.37
6	A	619	HEM	CHC-C1C-NC	2.62	127.31	124.45
3	A	681	NAG	C1-O5-C5	2.59	115.66	112.19
6	A	619	HEM	C1B-NB-C4B	2.54	108.22	105.21
6	A	619	HEM	CHA-C4D-ND	2.52	127.48	124.37
3	A	661	NAG	O5-C5-C4	-2.47	104.82	110.83
6	A	619	HEM	C3B-C2B-C1B	2.44	108.25	106.41
6	B	619	HEM	CBD-CAD-C3D	-2.32	106.11	112.53
4	B	3	BOG	O5-C5-C4	2.31	113.86	109.70
6	B	619	HEM	CMD-C2D-C1D	2.29	128.61	125.03
6	A	619	HEM	CMD-C2D-C1D	2.26	128.56	125.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	3	BOG	O1-C1-C2	-2.21	104.92	108.27
6	B	619	HEM	C3B-C2B-C1B	2.12	108.00	106.41
4	A	6	BOG	O5-C5-C4	2.12	113.51	109.70
6	A	619	HEM	CBD-CAD-C3D	-2.11	106.71	112.53
4	A	620	BOG	C1'-O1-C1	2.07	117.22	113.68

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	661	NAG	C1

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	661	NAG	C3-C2-N2-C7
3	A	661	NAG	O7-C7-N2-C2
4	A	620	BOG	O5-C1-O1-C1'
3	A	661	NAG	C8-C7-N2-C2
3	A	661	NAG	O5-C5-C6-O6
3	B	9	NAG	O5-C5-C6-O6
3	A	661	NAG	C4-C5-C6-O6
3	B	9	NAG	C4-C5-C6-O6
4	A	3	BOG	O1-C1'-C2'-C3'
4	B	3	BOG	C2-C1-O1-C1'
4	B	3	BOG	O5-C1-O1-C1'
4	B	3	BOG	C3'-C4'-C5'-C6'
4	B	3	BOG	C2'-C3'-C4'-C5'
4	A	3	BOG	C1'-C2'-C3'-C4'
4	B	3	BOG	C4'-C5'-C6'-C7'
4	B	3	BOG	C1'-C2'-C3'-C4'
4	A	6	BOG	C3'-C4'-C5'-C6'
4	A	620	BOG	C2-C1-O1-C1'
4	A	620	BOG	O5-C5-C6-O6
4	A	620	BOG	C4'-C5'-C6'-C7'
4	A	3	BOG	C4'-C5'-C6'-C7'
4	A	3	BOG	C3'-C4'-C5'-C6'
4	A	620	BOG	C2'-C3'-C4'-C5'
4	A	3	BOG	C2'-C3'-C4'-C5'
4	A	6	BOG	C4'-C5'-C6'-C7'
6	B	619	HEM	CAA-CBA-CGA-O2A
6	B	619	HEM	CAA-CBA-CGA-O1A
4	A	6	BOG	C5'-C6'-C7'-C8'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	4	NPS	C10-C14-C15-O
4	B	3	BOG	C4-C5-C6-O6


There are no ring outliers.



3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	619	HEM	1	0
4	A	620	BOG	3	0
6	A	619	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	552/587 (94%)	0.11	21 (3%)	44 53	14, 25, 40, 50	3 (0%)
1	B	552/587 (94%)	0.21	25 (4%)	38 47	16, 25, 41, 55	0
All	All	1104/1174 (94%)	0.16	46 (4%)	40 49	14, 25, 40, 55	3 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	577	PHE	4.4
1	B	583	GLN	4.3
1	B	409	TYR	4.2
1	B	33	ALA	4.1
1	A	583	GLN	4.0
1	A	81	LEU	3.8
1	A	75	LEU	3.7
1	B	122	TYR	3.7
1	A	77	ARG	3.2
1	A	409	TYR	3.0
1	B	268	ASP	3.0
1	A	80	LEU	3.0
1	B	80	LEU	2.9
1	B	74	PHE	2.8
1	A	122	TYR	2.8
1	B	281	GLU	2.8
1	B	81	LEU	2.8
1	A	169	LYS	2.7
1	A	158	ASP	2.6
1	A	239	ASP	2.6
1	B	215	LYS	2.5
1	B	101	ASN	2.4
1	A	74	PHE	2.4
1	A	108	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	214	HIS	2.4
1	B	473	LYS	2.4
1	B	54	GLN	2.4
1	B	239	ASP	2.4
1	B	399	ASP	2.4
1	B	496	SER	2.3
1	A	54	GLN	2.3
1	A	214	HIS	2.2
1	A	33	ALA	2.2
1	B	272	GLU	2.2
1	A	415	LEU	2.2
1	A	107	PHE	2.2
1	B	78	ILE	2.2
1	B	82	LEU	2.1
1	B	294	LEU	2.1
1	B	364	GLU	2.1
1	A	399	ASP	2.1
1	A	82	LEU	2.1
1	A	426	PHE	2.1
1	A	421	GLN	2.0
1	B	500	VAL	2.0
1	B	169	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	2	14/15	0.70	0.16	38,42,44,45	0
2	NAG	C	2	14/15	0.81	0.14	35,38,40,40	0
2	NAG	C	1	14/15	0.92	0.10	22,27,29,31	0
2	NAG	D	1	14/15	0.93	0.08	22,26,28,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	661	14/15	0.61	0.19	38,44,46,47	0
4	BOG	A	620	20/20	0.73	0.17	54,58,59,59	0
3	NAG	B	9	14/15	0.81	0.14	37,42,45,45	0
3	NAG	B	681	14/15	0.83	0.11	33,36,38,39	0
4	BOG	B	3	20/20	0.84	0.14	43,45,50,50	0
3	NAG	A	681	14/15	0.87	0.10	32,35,38,38	0
4	BOG	A	6	20/20	0.88	0.12	41,42,46,46	0
4	BOG	A	3	20/20	0.91	0.11	30,33,43,43	0
5	NPS	A	5	17/17	0.95	0.06	20,21,23,24	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NPS	B	4	17/17	0.96	0.05	20,21,24,25	0
6	HEM	A	619	43/43	0.96	0.09	26,28,38,41	0
6	HEM	B	619	43/43	0.96	0.09	26,28,37,42	0
7	CL	B	1	1/1	0.98	0.05	26,26,26,26	0
7	CL	A	1	1/1	0.99	0.03	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around BOG B 3:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG A 6:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG A 3:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 619:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.