



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 08:59 PM UTC

PDB ID : 3NTK / pdb_00003ntk
Title : Crystal structure of Tudor
Authors : Liu, H.P.; Huang, Y.; Li, Z.Z.; Gong, W.M.; Xu, R.M.
Deposited on : 2010-07-05
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	14% 82% 17% .
1	B	169	13% 79% 18% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2852 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maternal protein tudor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	169	Total	C	N	O	S	0	0	0
			1346	848	224	265	9			
1	B	166	Total	C	N	O	S	0	0	0
			1320	835	221	255	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	85	Total	O	0	0
			85	85		
2	B	101	Total	O	0	0
			101	101		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Maternal protein tudor



• Molecule 1: Maternal protein tudor



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.14Å 47.26Å 85.40Å 90.00° 106.24° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.2 (50.00-1.80) 96.2 (50.00-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 1.79Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.210 , 0.238 0.217 , 0.245	Depositor DCC
R_{free} test set	3616 reflections (10.07%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.498	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2852	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.36	0/1371	0.89	6/1851 (0.3%)
1	B	0.36	0/1344	0.92	9/1813 (0.5%)
All	All	0.36	0/2715	0.90	15/3664 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	2459	ASP	N-CA-C	-7.62	98.18	109.15
1	B	2499	ARG	N-CA-C	-6.92	97.93	109.07
1	B	2498	VAL	N-CA-C	6.33	118.03	108.80
1	B	2470	LEU	CA-C-N	-5.92	110.22	121.54
1	B	2470	LEU	C-N-CA	-5.92	110.22	121.54
1	A	2428	ILE	N-CA-C	5.70	118.17	111.05
1	B	2453	SER	N-CA-C	-5.66	100.47	109.59
1	A	2498	VAL	N-CA-C	5.65	117.05	108.80
1	B	2486	GLU	N-CA-C	-5.53	99.50	108.52
1	A	2497	VAL	N-CA-C	-5.51	99.95	107.99
1	B	2480	SER	N-CA-C	-5.37	101.42	109.79
1	B	2497	VAL	N-CA-C	-5.10	100.42	108.23
1	A	2453	SER	N-CA-C	-5.05	101.64	109.72
1	B	2431	GLY	N-CA-C	5.02	122.04	115.36
1	A	2512	GLN	N-CA-C	5.01	117.47	111.71

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	2470	LEU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1346	0	1299	20	0
1	B	1320	0	1277	37	0
2	A	85	0	0	0	0
2	B	101	0	0	1	0
All	All	2852	0	2576	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2470:LEU:HD23	1:B:2471:LEU:N	1.26	1.49
1:B:2470:LEU:HD23	1:B:2471:LEU:CA	1.53	1.38
1:B:2470:LEU:CD2	1:B:2471:LEU:H	1.35	1.35
1:B:2470:LEU:CD2	1:B:2471:LEU:HB3	1.57	1.33
1:B:2470:LEU:HD23	1:B:2471:LEU:CB	1.58	1.31
1:B:2470:LEU:CD2	1:B:2471:LEU:CB	2.15	1.22
1:B:2470:LEU:CG	1:B:2471:LEU:H	1.42	1.20
1:B:2470:LEU:CD2	1:B:2471:LEU:N	1.98	1.07
1:B:2426:HIS:HE2	1:B:2451:ARG:HH22	1.16	0.90
1:B:2470:LEU:HD21	1:B:2471:LEU:HB3	1.55	0.88
1:B:2470:LEU:CG	1:B:2471:LEU:N	2.25	0.86
1:A:2426:HIS:HE1	1:A:2451:ARG:HH22	1.21	0.83
1:B:2359:PRO:HG3	1:B:2471:LEU:HG	1.60	0.81
1:B:2466:CYS:HG	1:B:2514:CYS:HG	0.81	0.80
1:B:2470:LEU:HD21	1:B:2471:LEU:CB	2.07	0.79
1:A:2359:PRO:HB3	1:A:2458:LEU:HD12	1.65	0.77
1:B:2445:GLU:HG2	2:B:36:HOH:O	1.91	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2426:HIS:CE1	1:A:2451:ARG:HH22	2.08	0.70
1:A:2426:HIS:HE1	1:A:2451:ARG:NH2	1.92	0.67
1:A:2426:HIS:HD2	1:A:2433:ASN:OD1	1.78	0.65
1:B:2376:MET:O	1:B:2380:LEU:HD13	1.96	0.65
1:A:2360:MET:HG2	1:B:2479:PHE:O	1.98	0.62
1:B:2359:PRO:CG	1:B:2471:LEU:HG	2.28	0.60
1:A:2360:MET:HG2	1:B:2479:PHE:C	2.26	0.60
1:B:2470:LEU:C	1:B:2472:GLN:N	2.59	0.60
1:B:2470:LEU:HD22	1:B:2471:LEU:HB3	1.73	0.60
1:B:2455:HIS:HD2	1:B:2495:THR:OG1	1.85	0.59
1:B:2470:LEU:HG	1:B:2471:LEU:H	1.53	0.58
1:A:2360:MET:HG3	1:B:2477:THR:O	2.04	0.57
1:B:2470:LEU:CD2	1:B:2471:LEU:HB2	2.28	0.57
1:B:2471:LEU:HD12	1:B:2474:PHE:CD1	2.39	0.57
1:A:2470:LEU:HD11	1:A:2513:GLU:HB2	1.88	0.55
1:A:2462:THR:HG21	1:A:2499:ARG:HG2	1.90	0.54
1:B:2361:SER:OG	1:B:2455:HIS:HE1	1.91	0.54
1:B:2359:PRO:HB3	1:B:2471:LEU:HD21	1.90	0.53
1:B:2370:VAL:O	1:B:2374:GLU:HG3	2.09	0.52
1:B:2470:LEU:C	1:B:2472:GLN:H	2.19	0.51
1:B:2470:LEU:HD21	1:B:2471:LEU:HB2	1.88	0.49
1:B:2426:HIS:ND1	1:B:2433:ASN:ND2	2.61	0.48
1:A:2423:CYS:HB2	1:A:2439:PHE:CZ	2.49	0.47
1:A:2370:VAL:O	1:A:2374:GLU:HG3	2.16	0.46
1:B:2376:MET:SD	1:B:2380:LEU:HD11	2.56	0.46
1:A:2471:LEU:O	1:A:2475:ILE:HG12	2.15	0.45
1:B:2471:LEU:HG	1:B:2471:LEU:O	2.16	0.44
1:A:2393:LEU:HD13	1:A:2423:CYS:SG	2.58	0.44
1:A:2349:HIS:HB2	1:A:2366:MET:HE1	1.98	0.44
1:A:2403:PHE:O	1:A:2407:GLU:N	2.45	0.43
1:B:2502:TYR:CD2	1:B:2503:GLN:HG3	2.54	0.43
1:B:2376:MET:SD	1:B:2380:LEU:CD1	3.07	0.42
1:A:2457:GLU:CD	1:A:2491:LYS:HZ2	2.28	0.41
1:A:2471:LEU:HD21	1:A:2507:ILE:HG12	2.03	0.41
1:A:2457:GLU:CD	1:A:2491:LYS:NZ	2.79	0.40
1:A:2423:CYS:HB2	1:A:2439:PHE:HZ	1.86	0.40
1:B:2423:CYS:HB2	1:B:2439:PHE:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	167/169 (99%)	163 (98%)	4 (2%)	0	100	100
1	B	162/169 (96%)	154 (95%)	8 (5%)	0	100	100
All	All	329/338 (97%)	317 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/149 (100%)	146 (98%)	3 (2%)	48	38
1	B	145/149 (97%)	144 (99%)	1 (1%)	76	73
All	All	294/298 (99%)	290 (99%)	4 (1%)	59	52

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2360	MET
1	A	2470	LEU
1	A	2479	PHE
1	B	2373	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2349	HIS
1	A	2355	GLN
1	A	2365	GLN
1	A	2426	HIS
1	A	2472	GLN
1	A	2506	ASN
1	B	2365	GLN
1	B	2433	ASN
1	B	2437	GLN
1	B	2455	HIS
1	B	2484	GLN
1	B	2512	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	169/169 (100%)	0.97	23 (13%) 7 5	21, 30, 45, 50	0
1	B	166/169 (98%)	0.88	22 (13%) 7 6	19, 28, 41, 49	0
All	All	335/338 (99%)	0.92	45 (13%) 7 5	19, 29, 44, 50	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2466	CYS	10.3
1	B	2470	LEU	5.9
1	A	2406	ASP	5.5
1	B	2419	ASP	5.0
1	A	2382	ASP	4.5
1	B	2465	LYS	4.4
1	A	2419	ASP	4.3
1	B	2392	ASP	4.2
1	A	2346	ALA	4.1
1	B	2346	ALA	4.1
1	B	2471	LEU	4.0
1	B	2461	SER	3.7
1	A	2477	THR	3.5
1	A	2479	PHE	3.5
1	A	2391	SER	3.4
1	B	2514	CYS	3.3
1	B	2382	ASP	3.2
1	A	2378	ASP	3.2
1	B	2378	ASP	3.1
1	B	2420	ASP	3.0
1	A	2509	GLU	3.0
1	B	2445	GLU	2.9
1	A	2476	ASP	2.9
1	A	2360	MET	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2514	CYS	2.5
1	A	2418	LEU	2.4
1	B	2374	GLU	2.4
1	B	2464	SER	2.4
1	B	2473	SER	2.4
1	A	2445	GLU	2.3
1	B	2509	GLU	2.3
1	B	2360	MET	2.3
1	A	2352	VAL	2.3
1	A	2481	GLU	2.3
1	A	2347	GLU	2.2
1	A	2384	GLU	2.2
1	A	2420	ASP	2.2
1	B	2460	ALA	2.2
1	A	2368	SER	2.1
1	A	2392	ASP	2.1
1	A	2478	ARG	2.1
1	A	2508	SER	2.1
1	B	2356	PHE	2.0
1	B	2481	GLU	2.0
1	B	2405	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.