



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:16 PM UTC

PDB ID : 5NT1 / pdb_00005nt1
Title : Complex of influenza A NS1 effector domain with TRIM25 coiled coil
Authors : Koliopoulos, M.G.; Rittinger, K.
Deposited on : 2017-04-27
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	% 71% 17% .. 10%
1	E	193	% 75% 13% • 10%
1	I	193	% 70% 18% .. 10%
2	B	154	% 68% 10% 22%
2	F	154	% 73% 5% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	J	154	 A horizontal bar chart showing the quality of chain J. The bar is divided into three segments: a green segment representing 69%, a yellow segment representing 8%, and a grey segment representing 23%. The percentages are labeled below the bar.

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin/ISG15 ligase TRIM25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	173	Total	C	N	O	S	0	0	0
			1401	869	250	278	4			
1	E	173	Total	C	N	O	S	0	0	0
			1401	869	250	278	4			
1	I	173	Total	C	N	O	S	0	0	0
			1399	868	250	277	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	187	GLY	-	expression tag	UNP Q14258
A	188	PRO	-	expression tag	UNP Q14258
A	189	GLY	-	expression tag	UNP Q14258
A	358	LEU	PRO	variant	UNP Q14258
E	187	GLY	-	expression tag	UNP Q14258
E	188	PRO	-	expression tag	UNP Q14258
E	189	GLY	-	expression tag	UNP Q14258
E	358	LEU	PRO	variant	UNP Q14258
I	187	GLY	-	expression tag	UNP Q14258
I	188	PRO	-	expression tag	UNP Q14258
I	189	GLY	-	expression tag	UNP Q14258
I	358	LEU	PRO	variant	UNP Q14258

- Molecule 2 is a protein called Non-structural protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	120	Total	C	N	O	S	0	0	0
			945	601	162	176	6			
2	F	120	Total	C	N	O	S	0	0	0
			945	601	162	176	6			
2	J	119	Total	C	N	O	S	0	0	0
			939	598	161	174	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	77	GLY	-	expression tag	UNP P03496
B	78	PRO	-	expression tag	UNP P03496
B	79	GLY	-	expression tag	UNP P03496
B	101	GLU	ASP	variant	UNP P03496
F	77	GLY	-	expression tag	UNP P03496
F	78	PRO	-	expression tag	UNP P03496
F	79	GLY	-	expression tag	UNP P03496
F	101	GLU	ASP	variant	UNP P03496
J	77	GLY	-	expression tag	UNP P03496
J	78	PRO	-	expression tag	UNP P03496
J	79	GLY	-	expression tag	UNP P03496
J	101	GLU	ASP	variant	UNP P03496


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	E	1	Total O 1 1	0	0
3	I	2	Total O 2 2	0	0
3	J	4	Total O 4 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin/ISG15 ligase TRIM25




- Molecule 1: E3 ubiquitin/ISG15 ligase TRIM25



- Molecule 1: E3 ubiquitin/ISG15 ligase TRIM25



- Molecule 2: Non-structural protein 1



GLY	PRO	GLY	THR	MET	ALA	SER	VAL	ARG	PRO	D82	M98	I106	D120	I121	I122	I123	K131	I137	F138	D139	R140	I141	E142	V157	R204	SER	SER	ASN	GLU	ASN	GLY	ARG	PRO	PRO	THR	PRO	LYS	GLN	LYS	ARG	GLU	GLU	MET	ALA	GLY	THR	ILE	ARG	SER	GLU
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.38Å 225.23Å 146.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.69 – 2.82 48.69 – 2.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.69-2.82) 97.5 (48.69-2.82)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.195 , 0.224 0.197 , 0.223	Depositor DCC
R_{free} test set	2601 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	70.2	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.399 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.407 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7038	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1413	0.49	0/1892
1	E	0.27	0/1413	0.48	0/1892
1	I	0.27	0/1411	0.47	0/1889
2	B	0.28	0/961	0.47	0/1301
2	F	0.26	0/961	0.47	0/1301
2	J	0.30	0/955	0.46	0/1293
All	All	0.27	0/7114	0.47	0/9568

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1401	0	1450	17	0
1	E	1401	0	1450	15	0
1	I	1399	0	1446	21	0
2	B	945	0	964	9	0
2	F	945	0	964	4	0
2	J	939	0	959	8	0
3	A	1	0	0	0	0
3	E	1	0	0	1	0
3	I	2	0	0	1	0
3	J	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7038	0	7233	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:256:GLU:O	1:E:260:THR:HG23	1.91	0.69
2:B:196:GLU:OE2	2:B:200:ARG:NH1	2.25	0.69
2:F:96:GLU:OE1	2:F:100:ARG:NE	2.29	0.65
1:A:256:GLU:O	1:A:260:THR:HG23	1.99	0.61
1:I:248:MET:HA	1:I:251:LEU:HD12	1.83	0.60
2:B:101:GLU:OE2	1:I:212:GLN:NE2	2.29	0.59
1:I:222:ASP:OD2	1:I:226:ARG:NH1	2.36	0.58
2:J:137:ILE:HG22	2:J:142:GLU:HB2	1.85	0.58
1:I:254:ALA:O	1:I:258:THR:HG23	2.04	0.56
1:A:209:MET:HA	1:A:212:GLN:CD	2.31	0.56
2:F:120:ASP:HB3	2:F:123:ILE:HG12	1.90	0.53
2:B:203:TRP:O	2:B:204:ARG:NH1	2.42	0.53
1:I:284:LYS:NZ	1:I:319:THR:OG1	2.27	0.52
2:J:106:ILE:HG13	2:J:122:ALA:HB2	1.92	0.52
1:I:222:ASP:HB3	3:I:401:HOH:O	2.10	0.51
1:I:240:GLN:O	1:I:243:GLN:HG2	2.11	0.51
2:B:175:LYS:HD3	2:B:202:ALA:O	2.11	0.50
2:F:137:ILE:HG22	2:F:142:GLU:HB2	1.92	0.50
1:E:240:GLN:O	1:E:243:GLN:HG2	2.11	0.49
1:E:306:GLU:OE1	1:E:306:GLU:N	2.36	0.49
1:E:196:ALA:HA	1:E:199:GLU:HB2	1.94	0.49
2:B:98:MET:HE2	1:I:222:ASP:HB2	1.95	0.49
1:I:280:ILE:HD11	1:I:324:ILE:HD12	1.95	0.48
1:A:237:LYS:HE3	1:A:240:GLN:OE1	2.13	0.48
2:B:106:ILE:HG13	2:B:122:ALA:HB2	1.95	0.48
1:A:317:ILE:HG23	1:A:319:THR:HG23	1.96	0.48
2:J:120:ASP:HB3	2:J:123:ILE:HG12	1.96	0.48
2:F:137:ILE:HG12	2:F:138:PHE:CD2	2.50	0.47
1:I:276:THR:O	1:I:280:ILE:HG23	2.14	0.47
1:I:265:GLU:OE2	1:I:266:GLU:N	2.48	0.47
2:B:202:ALA:HB3	2:B:203:TRP:CE3	2.50	0.47
1:E:293:GLU:O	1:E:297:GLN:HG2	2.15	0.47
1:E:318:SER:HB2	1:E:320:LYS:HE3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:321:PRO:O	1:A:322:VAL:HG12	2.15	0.46
1:A:321:PRO:HB2	1:A:322:VAL:H	1.58	0.46
2:B:98:MET:HE3	2:B:98:MET:HB3	1.84	0.46
2:B:182:ILE:HG12	2:B:192:VAL:HG11	1.96	0.46
1:A:293:GLU:O	1:A:297:GLN:HG2	2.16	0.45
1:I:318:SER:HB2	1:I:320:LYS:HE3	1.98	0.45
1:I:295:ILE:HG12	1:I:314:LEU:HD13	1.98	0.45
1:A:205:LYS:HE2	1:A:209:MET:HE3	1.99	0.45
1:A:273:LYS:HD3	1:A:327:VAL:HG22	1.97	0.45
1:E:332:LYS:HA	1:E:332:LYS:HD3	1.80	0.44
1:A:343:ASP:O	1:A:347:GLU:HG2	2.17	0.44
1:E:245:TYR:OH	3:E:401:HOH:O	2.20	0.44
1:E:283:LYS:HE3	1:E:287:GLU:OE2	2.18	0.44
1:I:293:GLU:O	1:I:297:GLN:HG2	2.16	0.44
2:J:137:ILE:HG13	2:J:138:PHE:CD2	2.53	0.44
1:A:222:ASP:HB2	2:J:98:MET:HE2	2.00	0.44
1:E:345:LYS:HE2	1:E:345:LYS:HB3	1.79	0.44
1:E:284:LYS:O	1:E:288:ILE:HG12	2.18	0.44
2:J:140:ARG:NH1	2:J:204:ARG:HD2	2.33	0.44
1:A:308:LEU:HD12	1:A:308:LEU:H	1.83	0.43
1:E:319:THR:C	1:E:321:PRO:HD3	2.43	0.43
1:I:319:THR:C	1:I:321:PRO:HD3	2.43	0.43
1:I:306:GLU:OE1	1:I:306:GLU:N	2.35	0.43
2:J:92:ASP:HB3	2:J:131:LYS:HB3	2.01	0.43
1:A:209:MET:O	1:A:212:GLN:HG2	2.19	0.43
1:I:321:PRO:HB2	1:I:322:VAL:H	1.62	0.43
1:A:302:ARG:HG2	1:A:303:ASP:N	2.34	0.42
1:E:343:ASP:O	1:E:347:GLU:HG2	2.19	0.42
1:I:245:TYR:HA	1:I:248:MET:HE3	2.02	0.42
1:I:191:LEU:HD23	1:I:191:LEU:HA	1.88	0.42
1:E:301:LYS:HB3	1:E:304:GLU:CD	2.45	0.41
2:J:123:ILE:HG13	2:J:157:VAL:HB	2.02	0.41
1:A:222:ASP:OD2	1:A:226:ARG:NH1	2.50	0.41
1:I:268:LYS:HD2	1:I:268:LYS:HA	1.86	0.41
1:A:198:LEU:O	1:A:202:LEU:HG	2.21	0.41
1:A:298:SER:OG	1:A:310:LYS:NZ	2.54	0.41
1:E:321:PRO:HB2	1:E:322:VAL:H	1.64	0.40
1:I:301:LYS:HB3	1:I:304:GLU:CD	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/193 (89%)	167 (98%)	2 (1%)	2 (1%)	10	31
1	E	171/193 (89%)	165 (96%)	4 (2%)	2 (1%)	10	31
1	I	171/193 (89%)	165 (96%)	4 (2%)	2 (1%)	10	31
2	B	118/154 (77%)	116 (98%)	2 (2%)	0	100	100
2	F	118/154 (77%)	115 (98%)	3 (2%)	0	100	100
2	J	117/154 (76%)	112 (96%)	5 (4%)	0	100	100
All	All	866/1041 (83%)	840 (97%)	20 (2%)	6 (1%)	18	45

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	PRO
1	E	321	PRO
1	I	321	PRO
1	A	322	VAL
1	I	322	VAL
1	E	322	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/175 (91%)	148 (92%)	12 (8%)	12	35
1	E	160/175 (91%)	155 (97%)	5 (3%)	35	69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	159/175 (91%)	150 (94%)	9 (6%)	18	47
2	B	104/132 (79%)	101 (97%)	3 (3%)	37	71
2	F	104/132 (79%)	104 (100%)	0	100	100
2	J	103/132 (78%)	103 (100%)	0	100	100
All	All	790/921 (86%)	761 (96%)	29 (4%)	30	63

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	199	GLU
1	A	201	THR
1	A	230	VAL
1	A	241	LEU
1	A	264	LYS
1	A	270	VAL
1	A	282	LEU
1	A	289	GLN
1	A	292	LYS
1	A	301	LYS
1	A	302	ARG
1	A	322	VAL
2	B	96	GLU
2	B	99	SER
2	B	136	VAL
1	E	202	LEU
1	E	230	VAL
1	E	260	THR
1	E	324	ILE
1	E	349	LYS
1	I	198	LEU
1	I	202	LEU
1	I	221	ASP
1	I	230	VAL
1	I	238	VAL
1	I	270	VAL
1	I	280	ILE
1	I	289	GLN
1	I	322	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	331	HIS
2	B	176	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	173/193 (89%)	-0.99	1 (0%) 85 80	45, 73, 183, 250	0
1	E	173/193 (89%)	-0.98	1 (0%) 85 80	44, 75, 180, 245	0
1	I	173/193 (89%)	-1.00	2 (1%) 76 68	45, 77, 174, 222	0
2	B	120/154 (77%)	-1.43	0 100 100	41, 60, 97, 196	0
2	F	120/154 (77%)	-1.45	0 100 100	43, 59, 97, 215	0
2	J	119/154 (77%)	-1.38	0 100 100	38, 59, 99, 152	0
All	All	878/1041 (84%)	-1.17	4 (0%) 87 82	38, 66, 172, 250	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	362	SER	3.1
1	E	321	PRO	2.7
1	A	305	PHE	2.6
1	I	319	THR	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.