



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2026 – 06:53 PM UTC

PDB ID : 6NTI / pdb\_00006nti  
Title : Neutron/X-ray crystal structure of AAC-VIa bound to kanamycin b  
Authors : Cuneo, M.J.; Kumar, P.  
Deposited on : 2019-01-29  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

|                                |   |                                                                  |
|--------------------------------|---|------------------------------------------------------------------|
| MolProbity                     | : | 4-5-2 with Phenix2.0                                             |
| Mogul                          | : | 2022.3.0, CSD as543be (2022)                                     |
| Xtriage (Phenix)               | : | 2.0                                                              |
| EDS                            | : | <b>FAILED</b>                                                    |
| Percentile statistics          | : | 20250101.v01 (using entries in the PDB archive January 1st 2025) |
| Ideal geometry (proteins)      | : | Engh & Huber (2001)                                              |
| Ideal geometry (DNA, RNA)      | : | Parkinson et al. (1996)                                          |
| Validation Pipeline (wwPDB-VP) | : | 2.49                                                             |


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION, NEUTRON DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|-------------------------------------------------------|
| Clashscore            | 190562                      | 6919 (2.30-2.30)                                      |
| Ramachandran outliers | 187476                      | 6854 (2.30-2.30)                                      |
| Sidechain outliers    | 187428                      | 6854 (2.30-2.30)                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS failed to run properly.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 302    | <br>83% 5% 11%   |

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4726 atoms, of which 2014 are hydrogens and 581 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aminoglycoside N(3)-acetyltransferase.

| Mol | Chain | Residues | Atoms |      |     |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|------|-----|-----|---|---------|---------|-------|
|     |       |          | Total | C    | D   | H    | N   | O   | S |         |         |       |
| 1   | A     | 268      | 4429  | 1281 | 425 | 1977 | 367 | 373 | 6 | 87      | 245     | 0     |

There are 3 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment        | Reference  |
|-------|---------|----------|--------|----------------|------------|
| A     | -2      | GLY      | -      | expression tag | UNP C4NV15 |
| A     | -1      | SER      | -      | expression tag | UNP C4NV15 |
| A     | 0       | HIS      | -      | expression tag | UNP C4NV15 |

- Molecule 2 is (1R,2S,3S,4R,6S)-4,6-DIAMINO-3-[(3-AMINO-3-DEOXY-ALPHA-D-GLUCOPYRANOSYL)OXY]-2-HYDROXYCYCLOHEXYL 2,6-DIAMINO-2,6-DIDEOXY-ALPHA-D-GLUCOPYRANOSIDE (CCD ID: 9CS) (formula: C<sub>18</sub>H<sub>37</sub>N<sub>5</sub>O<sub>10</sub>).



| Mol | Chain | Residues | Atoms |    |    |    |   |    | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|----|----|---|----|---------|---------|
|     |       |          | Total | C  | D  | H  | N | O  |         |         |
| 2   | A     | 1        | 86    | 18 | 16 | 37 | 5 | 10 | 10      | 1       |

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms      |         | ZeroOcc | AltConf |
|-----|-------|----------|------------|---------|---------|---------|
| 3   | A     | 1        | Total<br>1 | Mg<br>1 | 0       | 0       |

- Molecule 4 is water.

| Mol | Chain | Residues | Atoms        |          |         | ZeroOcc | AltConf |
|-----|-------|----------|--------------|----------|---------|---------|---------|
| 4   | A     | 70       | Total<br>210 | D<br>140 | O<br>70 | 1       | 0       |


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: Aminoglycoside N(3)-acetyltransferase

Chain A: 



## 4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

| Property                                                 | Value                                                       | Source    |
|----------------------------------------------------------|-------------------------------------------------------------|-----------|
| Space group                                              | I 1 2 1                                                     | Depositor |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$ | 51.36Å 86.06Å 77.60Å<br>90.00° 93.98° 90.00°                | Depositor |
| Resolution (Å)                                           | 14.00 – 2.30                                                | Depositor |
| % Data completeness<br>(in resolution range)             | 86.0 (14.00-2.30)                                           | Depositor |
| $R_{merge}$                                              | 0.06                                                        | Depositor |
| $R_{sym}$                                                | (Not available)                                             | Depositor |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>               | 3.11 (at 2.29Å)                                             | Xtriage   |
| Refinement program                                       | PHENIX 1.13_2998                                            | Depositor |
| R, $R_{free}$                                            | 0.237 , 0.257                                               | Depositor |
| Wilson B-factor (Å <sup>2</sup> )                        | 41.1                                                        | Xtriage   |
| Anisotropy                                               | 0.496                                                       | Xtriage   |
| L-test for twinning <sup>2</sup>                         | $\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$ | Xtriage   |
| Estimated twinning fraction                              | No twinning to report.                                      | Xtriage   |
| Total number of atoms                                    | 4726                                                        | wwPDB-VP  |
| Average B, all atoms (Å <sup>2</sup> )                   | 54.0                                                        | wwPDB-VP  |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 9CS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |                | Bond angles |             |
|-----|-------|--------------|----------------|-------------|-------------|
|     |       | RMSZ         | $\# Z  > 5$    | RMSZ        | $\# Z  > 5$ |
| 1   | A     | 0.63         | 18/3986 (0.5%) | 0.48        | 0/5448      |

The worst 5 of 18 bond length outliers are listed below:

| Mol | Chain | Res    | Type | Atoms | Z     | Observed(Å) | Ideal(Å) |
|-----|-------|--------|------|-------|-------|-------------|----------|
| 1   | A     | 214[A] | LYS  | C-N   | 14.65 | 1.53        | 1.33     |
| 1   | A     | 214[B] | LYS  | C-N   | 14.65 | 1.53        | 1.33     |
| 1   | A     | 22[A]  | SER  | C-N   | 10.80 | 1.47        | 1.33     |
| 1   | A     | 22[B]  | SER  | C-N   | 10.80 | 1.47        | 1.33     |
| 1   | A     | 24[A]  | SER  | C-N   | -7.29 | 1.24        | 1.33     |

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

CLOSE-CONTACTS INFOmissingINFO

### 5.2 Torsion angles ⓘ

#### 5.2.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed       | Favoured  | Allowed | Outliers | Percentiles |    |
|-----|-------|----------------|-----------|---------|----------|-------------|----|
| 1   | A     | 509/302 (168%) | 492 (97%) | 15 (3%) | 2 (0%)   | 30          | 38 |

All (2) Ramachandran outliers are listed below:

| Mol | Chain | Res    | Type |
|-----|-------|--------|------|
| 1   | A     | 237[A] | GLY  |
| 1   | A     | 237[B] | GLY  |

### 5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed       | Rotameric | Outliers | Percentiles |
|-----|-------|----------------|-----------|----------|-------------|
| 1   | A     | 379/226 (168%) | 369 (97%) | 10 (3%)  | 40 59       |

5 of 10 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res    | Type |
|-----|-------|--------|------|
| 1   | A     | 210[B] | LEU  |
| 1   | A     | 215[A] | ARG  |
| 1   | A     | 215[B] | ARG  |
| 1   | A     | 68[B]  | VAL  |
| 1   | A     | 198[A] | GLU  |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.4 Carbohydrates [i](#)

There are no oligosaccharides in this entry.



## 5.5 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Other polymers

There are no such residues in this entry.

## 5.7 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 6.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 6.5 Other polymers

EDS failed to run properly - this section is therefore empty.