



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 03:16 AM UTC

PDB ID : 2NU9 / pdb\_00002nu9  
Title : C123aT Mutant of E. coli Succinyl-CoA Synthetase Orthorhombic Crystal Form  
Authors : Fraser, M.E.  
Deposited on : 2006-11-08  
Resolution : 2.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	 81% 17% ..
1	D	288	 76% 21% ..
1	F	288	 78% 20% ..
1	H	288	 76% 22% ..
2	B	388	 6% 77% 21% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	E	388	 6% 76% 22% ..
2	G	388	 3% 76% 22% ..
2	I	388	 5% 74% 23% ..

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinyl-CoA ligase [ADP-forming] subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	286	2066	1308	345	402	1	10	0	0	0
1	D	286	2066	1308	345	402	1	10	0	0	0
1	F	286	2066	1308	345	402	1	10	0	0	0
1	H	286	2066	1308	345	402	1	10	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	THR	CYS	engineered mutation	UNP P0AGE9
A	246	NEP	HIS	modified residue	UNP P0AGE9
D	123	THR	CYS	engineered mutation	UNP P0AGE9
D	246	NEP	HIS	modified residue	UNP P0AGE9
F	123	THR	CYS	engineered mutation	UNP P0AGE9
F	246	NEP	HIS	modified residue	UNP P0AGE9
H	123	THR	CYS	engineered mutation	UNP P0AGE9
H	246	NEP	HIS	modified residue	UNP P0AGE9

- Molecule 2 is a protein called Succinyl-CoA synthetase beta chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	385	2885	1823	505	544	13	0	0	0
2	E	385	2885	1823	505	544	13	0	0	0
2	G	385	2885	1823	505	544	13	0	0	0
2	I	385	2885	1823	505	544	13	0	0	0

- Molecule 3 is COENZYME A (CCD ID: COA) (formula:  $C_{21}H_{36}N_7O_{16}P_3S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
3	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	D	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	F	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	H	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	G	1	Total O S 5 4 1	0	0
4	G	1	Total O S 5 4 1	0	0
4	I	1	Total O S 5 4 1	0	0
4	I	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O 1 1	0	0
5	E	1	Total O 1 1	0	0
5	F	1	Total O 1 1	0	0
5	H	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Succinyl-CoA ligase [ADP-forming] subunit alpha




- Molecule 1: Succinyl-CoA ligase [ADP-forming] subunit alpha



- Molecule 1: Succinyl-CoA ligase [ADP-forming] subunit alpha





- Molecule 1: Succinyl-CoA ligase [ADP-forming] subunit alpha












● Molecule 2: Succinyl-CoA synthetase beta chain




● Molecule 2: Succinyl-CoA synthetase beta chain



● Molecule 2: Succinyl-CoA synthetase beta chain



- Molecule 2: Succinyl-CoA synthetase beta chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.53Å 154.90Å 240.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.11 – 2.90 20.11 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.11-2.90) 98.3 (20.11-2.90)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.88Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.229 , 0.251 0.228 , 0.248	Depositor DCC
$R_{free}$ test set	4125 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.8	Xtrriage
Anisotropy	0.511	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 30.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	20040	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NEP, COA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.49	0/2084	0.99	7/2822 (0.2%)
1	D	0.47	0/2084	1.00	7/2822 (0.2%)
1	F	0.50	0/2084	1.01	7/2822 (0.2%)
1	H	0.48	0/2084	1.01	7/2822 (0.2%)
2	B	0.51	1/2927 (0.0%)	0.92	7/3961 (0.2%)
2	E	0.51	1/2927 (0.0%)	0.92	9/3961 (0.2%)
2	G	0.50	0/2927	0.92	8/3961 (0.2%)
2	I	0.50	0/2927	0.92	7/3961 (0.2%)
All	All	0.50	2/20044 (0.0%)	0.95	59/27132 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	124	MET	SD-CE	5.38	1.93	1.79
2	B	124	MET	SD-CE	5.09	1.92	1.79

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	223	GLU	N-CA-C	7.40	119.42	111.36
1	H	14	GLY	N-CA-C	-7.17	104.89	115.72
2	I	202	VAL	N-CA-C	7.12	119.69	108.87
2	B	201	LEU	N-CA-C	-7.09	96.08	108.20
1	D	180	ASP	CA-C-N	7.05	126.88	119.05

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2066	0	2118	31	0
1	D	2066	0	2118	45	0
1	F	2066	0	2118	40	0
1	H	2066	0	2118	45	0
2	B	2885	0	2941	62	0
2	E	2885	0	2941	61	0
2	G	2885	0	2941	60	0
2	I	2885	0	2941	67	0
3	A	48	0	32	1	0
3	D	48	0	32	5	0
3	F	48	0	32	0	0
3	H	48	0	32	3	0
4	B	10	0	0	1	0
4	E	10	0	0	0	0
4	G	10	0	0	0	0
4	I	10	0	0	0	0
5	A	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	H	1	0	0	0	0
All	All	20040	0	20364	384	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 384 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:322:ILE:HD13	2:I:322:ILE:H	1.29	0.97
2:E:322:ILE:HD13	2:E:322:ILE:H	1.26	0.97
2:G:322:ILE:HD13	2:G:322:ILE:H	1.27	0.96
2:B:322:ILE:HD13	2:B:322:ILE:H	1.31	0.94
2:E:315:LEU:HB2	2:E:381:VAL:HG11	1.56	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/288 (98%)	272 (96%)	10 (4%)	1 (0%)	30	59
1	D	283/288 (98%)	271 (96%)	11 (4%)	1 (0%)	30	59
1	F	283/288 (98%)	271 (96%)	12 (4%)	0	100	100
1	H	283/288 (98%)	271 (96%)	11 (4%)	1 (0%)	30	59
2	B	383/388 (99%)	373 (97%)	9 (2%)	1 (0%)	36	65
2	E	383/388 (99%)	373 (97%)	9 (2%)	1 (0%)	36	65
2	G	383/388 (99%)	372 (97%)	10 (3%)	1 (0%)	36	65
2	I	383/388 (99%)	373 (97%)	9 (2%)	1 (0%)	36	65
All	All	2664/2704 (98%)	2576 (97%)	81 (3%)	7 (0%)	36	65

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	384	ALA
2	E	384	ALA
2	G	384	ALA
2	I	384	ALA
1	A	285	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/217 (99%)	209 (97%)	6 (3%)	38	72

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	215/217 (99%)	209 (97%)	6 (3%)	38	72
1	F	215/217 (99%)	210 (98%)	5 (2%)	44	76
1	H	215/217 (99%)	210 (98%)	5 (2%)	44	76
2	B	296/298 (99%)	290 (98%)	6 (2%)	48	78
2	E	296/298 (99%)	289 (98%)	7 (2%)	43	75
2	G	296/298 (99%)	288 (97%)	8 (3%)	39	73
2	I	296/298 (99%)	289 (98%)	7 (2%)	43	75
All	All	2044/2060 (99%)	1994 (98%)	50 (2%)	43	75

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	255	LYS
2	G	287	LEU
2	I	348	ARG
1	F	274	ARG
2	G	200	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	342	ASN
2	I	95	GLN
2	G	50	HIS
2	I	342	ASN
2	G	352	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	NEP	D	246	1	11,14,15	1.47	3 (27%)	8,20,22	1.15	1 (12%)
1	NEP	H	246	1	11,14,15	1.59	2 (18%)	8,20,22	1.19	1 (12%)
1	NEP	F	246	1	11,14,15	1.44	2 (18%)	8,20,22	1.18	1 (12%)
1	NEP	A	246	1	11,14,15	1.53	2 (18%)	8,20,22	1.12	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NEP	D	246	1	-	2/5/12/14	0/1/1/1
1	NEP	H	246	1	-	2/5/12/14	0/1/1/1
1	NEP	F	246	1	-	2/5/12/14	0/1/1/1
1	NEP	A	246	1	-	2/5/12/14	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	246	NEP	P-O1P	-3.57	1.47	1.56
1	H	246	NEP	P-O2P	-3.52	1.47	1.56
1	A	246	NEP	P-O1P	-3.39	1.47	1.56
1	H	246	NEP	P-O1P	-3.33	1.48	1.56
1	A	246	NEP	P-O2P	-3.10	1.48	1.56

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	246	NEP	CE1-ND1-CG	2.37	109.59	105.80
1	A	246	NEP	CE1-ND1-CG	2.33	109.54	105.80
1	H	246	NEP	CE1-ND1-CG	2.30	109.48	105.80
1	F	246	NEP	CE1-ND1-CG	2.28	109.46	105.80

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	F	246	NEP	CA-CB-CG-CD2
1	A	246	NEP	CA-CB-CG-CD2
1	D	246	NEP	CA-CB-CG-CD2
1	H	246	NEP	CA-CB-CG-CD2
1	A	246	NEP	CA-CB-CG-ND1

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	246	NEP	1	0
1	H	246	NEP	1	0
1	F	246	NEP	1	0
1	A	246	NEP	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	G	389	-	4,4,4	0.43	0	6,6,6	0.18	0
4	SO4	I	390	-	4,4,4	0.33	0	6,6,6	0.15	0
3	COA	H	401	-	47,50,50	0.84	3 (6%)	69,75,75	1.58	9 (13%)
4	SO4	B	390	-	4,4,4	0.43	0	6,6,6	0.16	0
4	SO4	E	390	-	4,4,4	0.33	0	6,6,6	0.08	0
4	SO4	E	389	-	4,4,4	0.39	0	6,6,6	0.14	0
4	SO4	I	389	-	4,4,4	0.56	0	6,6,6	0.18	0
3	COA	D	301	-	47,50,50	0.88	2 (4%)	69,75,75	1.20	6 (8%)
3	COA	A	300	-	47,50,50	0.87	3 (6%)	69,75,75	1.14	6 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	G	390	-	4,4,4	0.47	0	6,6,6	0.23	0
4	SO4	B	389	-	4,4,4	0.40	0	6,6,6	0.13	0
3	COA	F	400	-	47,50,50	0.95	2 (4%)	69,75,75	1.46	10 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	300	-	-	2/48/64/64	0/3/3/3
3	COA	H	401	-	-	5/48/64/64	0/3/3/3
3	COA	D	301	-	-	4/48/64/64	0/3/3/3
3	COA	F	400	-	-	7/48/64/64	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	300	COA	OAP-CAP	3.25	1.48	1.42
3	F	400	COA	P1A-O3A	2.83	1.62	1.59
3	D	301	COA	P1A-O3A	2.81	1.62	1.59
3	F	400	COA	P2A-O3A	2.73	1.62	1.59
3	D	301	COA	P2A-O3A	2.73	1.62	1.59

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	401	COA	C6P-C5P-N4P	6.38	127.97	116.34
3	H	401	COA	C3P-N4P-C5P	-4.46	114.51	122.82
3	F	400	COA	CDP-CBP-CCP	4.45	115.57	108.22
3	H	401	COA	O5P-C5P-N4P	-4.41	114.38	123.03
3	H	401	COA	C2B-C1B-N9A	4.10	123.50	113.30

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	300	COA	C2B-C3B-O3B-P3B
3	H	401	COA	S1P-C2P-C3P-N4P
3	D	301	COA	C2B-C3B-O3B-P3B

*Continued on next page...*

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	H	401	COA	C2B-C3B-O3B-P3B
3	F	400	COA	C2B-C3B-O3B-P3B


There are no ring outliers.


4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	401	COA	3	0
3	D	301	COA	5	0
3	A	300	COA	1	0
4	B	389	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/288 (98%)	-0.06	2 (0%) 84 79	18, 35, 51, 59	0
1	D	285/288 (98%)	-0.08	1 (0%) 88 85	19, 37, 51, 60	0
1	F	285/288 (98%)	-0.07	2 (0%) 84 79	18, 34, 50, 59	0
1	H	285/288 (98%)	-0.07	2 (0%) 84 79	19, 36, 51, 60	0
2	B	385/388 (99%)	0.28	25 (6%) 25 20	18, 36, 64, 79	0
2	E	385/388 (99%)	0.27	24 (6%) 26 21	19, 37, 64, 78	0
2	G	385/388 (99%)	0.08	12 (3%) 51 42	19, 37, 64, 78	0
2	I	385/388 (99%)	0.24	18 (4%) 36 28	19, 37, 65, 79	0
All	All	2680/2704 (99%)	0.10	86 (3%) 50 41	18, 36, 60, 79	0

The worst 5 of 86 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	322	ILE	6.0
2	B	322	ILE	5.4
2	G	322	ILE	4.9
2	B	138	GLU	4.6
2	G	320	GLY	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	NEP	F	246	14/15	0.96	0.09	27,33,34,34	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	NEP	D	246	14/15	0.97	0.06	30,33,38,38	0
1	NEP	A	246	14/15	0.97	0.06	29,32,35,36	0
1	NEP	H	246	14/15	0.98	0.07	29,32,36,36	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	I	389	5/5	0.79	0.23	68,68,68,69	5
4	SO4	G	389	5/5	0.85	0.19	22,23,24,25	5
4	SO4	E	389	5/5	0.88	0.19	28,29,30,31	5
3	COA	F	400	48/48	0.92	0.10	39,45,47,50	0
4	SO4	B	389	5/5	0.92	0.13	28,29,30,30	5
4	SO4	B	390	5/5	0.92	0.09	53,53,54,55	0
3	COA	A	300	48/48	0.93	0.11	43,48,52,53	0
3	COA	D	301	48/48	0.93	0.10	43,46,53,54	0
4	SO4	I	390	5/5	0.93	0.10	55,56,56,57	0
3	COA	H	401	48/48	0.94	0.10	33,44,58,59	0
4	SO4	G	390	5/5	0.95	0.09	68,68,69,69	0
4	SO4	E	390	5/5	0.96	0.10	56,56,56,56	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around COA F 400:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around COA A 300:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around COA D 301:**

$2mF_o-DF_c$  (at 0.7 rnsd) in gray  
 $mF_o-DF_c$  (at 3 rnsd) in purple (negative)  
and green (positive)

**Electron density around COA H 401:**

$2mF_o-DF_c$  (at 0.7 rnsd) in gray  
 $mF_o-DF_c$  (at 3 rnsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.