



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:29 AM UTC

PDB ID : 3NUM / pdb\_00003num  
Title : Substrate induced remodeling of the active site regulates HtrA1 activity  
Authors : Truebestein, L.; Tennstaedt, A.; Hauske, P.; Krojer, T.; Kaiser, M.; Clausen, T.; Ehrmann, M.  
Deposited on : 2010-07-07  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1407 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine protease HTRA1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
1	A	190	1407	900	236	271	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	157	MET	-	initiating methionine	UNP Q92743
A	328	ALA	SER	engineered mutation	UNP Q92743
A	481	LEU	-	expression tag	UNP Q92743
A	482	GLU	-	expression tag	UNP Q92743
A	483	HIS	-	expression tag	UNP Q92743
A	484	HIS	-	expression tag	UNP Q92743
A	485	HIS	-	expression tag	UNP Q92743
A	486	HIS	-	expression tag	UNP Q92743
A	487	HIS	-	expression tag	UNP Q92743
A	488	HIS	-	expression tag	UNP Q92743



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.48Å 109.48Å 113.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.32 – 2.75 19.32 – 2.75	Depositor EDS
% Data completeness (in resolution range)	95.7 (19.32-2.75) 95.6 (19.32-2.75)	Depositor EDS
$R_{merge}$	0.40	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.76Å)	Xtrriage
Refinement program	REFMAC 5.6.0066	Depositor
R, $R_{free}$	0.242 , 0.287 0.241 , 0.292	Depositor DCC
$R_{free}$ test set	630 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	84.3	Xtrriage
Anisotropy	0.237	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.031 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	1407	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.90	0/1428	1.00	5/1942 (0.3%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	LEU	CA-C-N	7.65	129.40	119.84
1	A	192	LEU	C-N-CA	7.65	129.40	119.84
1	A	282	ILE	N-CA-C	6.32	117.88	108.46
1	A	342	ILE	CB-CA-C	-5.90	102.81	110.84
1	A	320	ASP	CB-CA-C	-5.87	98.74	110.42

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1407	0	1410	34	0
All	All	1407	0	1410	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:THR:HG22	1:A:346:LYS:H	1.52	0.73
1:A:323:ILE:HD13	1:A:343:ASN:HB3	1.73	0.71
1:A:340:ILE:C	1:A:340:ILE:HD12	2.20	0.66
1:A:344:THR:HB	1:A:353:PHE:O	1.98	0.64
1:A:283:GLY:HA3	1:A:327:ASN:HA	1.84	0.59
1:A:259:GLN:OE1	1:A:259:GLN:N	2.34	0.58
1:A:172:ILE:O	1:A:176:VAL:HG12	2.04	0.58
1:A:207:PHE:HE1	1:A:342:ILE:CD1	2.17	0.58
1:A:340:ILE:C	1:A:340:ILE:CD1	2.82	0.53
1:A:207:PHE:CE1	1:A:342:ILE:CD1	2.92	0.53
1:A:258:HIS:CG	1:A:262:LEU:HD21	2.44	0.53
1:A:272:GLU:OE1	1:A:272:GLU:HA	2.09	0.52
1:A:232:LEU:O	1:A:235:GLY:N	2.42	0.51
1:A:272:GLU:OE1	1:A:272:GLU:CA	2.60	0.50
1:A:232:LEU:HD12	1:A:232:LEU:N	2.26	0.49
1:A:357:SER:HA	1:A:360:ILE:HD12	1.95	0.48
1:A:180:ALA:HB3	1:A:181:PRO:HD3	1.95	0.48
1:A:219:ALA:HB3	1:A:250:ASP:OD1	2.14	0.47
1:A:279:VAL:CG2	1:A:332:LEU:HD11	2.44	0.47
1:A:173:ALA:HA	1:A:176:VAL:CG1	2.45	0.47
1:A:340:ILE:HD12	1:A:341:GLY:N	2.30	0.47
1:A:188:LEU:HB3	1:A:202:ALA:HB3	1.97	0.46
1:A:230:VAL:HG23	1:A:240:ALA:HB2	1.98	0.45
1:A:323:ILE:HD13	1:A:343:ASN:CB	2.43	0.44
1:A:334:ASN:OD1	1:A:334:ASN:C	2.62	0.43
1:A:232:LEU:O	1:A:233:LYS:C	2.63	0.42
1:A:317:ILE:HD12	1:A:356:PRO:HG3	2.02	0.42
1:A:179:ILE:HD13	1:A:333:VAL:HG21	2.01	0.42
1:A:299:THR:O	1:A:300:THR:C	2.63	0.41
1:A:232:LEU:N	1:A:232:LEU:CD1	2.84	0.41
1:A:182:ALA:HA	1:A:208:ILE:HD12	2.03	0.41
1:A:334:ASN:HD21	1:A:338:GLU:HB2	1.86	0.41
1:A:360:ILE:HG22	1:A:364:LEU:CD1	2.51	0.41
1:A:207:PHE:CE1	1:A:342:ILE:HD11	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	184/332 (55%)	172 (94%)	8 (4%)	4 (2%)	5 10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	320	ASP
1	A	233	LYS
1	A	225	LYS
1	A	193	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	151/285 (53%)	140 (93%)	11 (7%)	13 24

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	176	VAL
1	A	198	GLU
1	A	201	VAL
1	A	271	SER
1	A	274	ARG
1	A	291	THR
1	A	292	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	294	THR
1	A	340	ILE
1	A	344	THR
1	A	369	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	190/332 (57%)	0.15	4 (2%) 63 61	52, 91, 123, 159	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300	THR	4.8
1	A	161	ASP	2.6
1	A	325	TYR	2.4
1	A	290	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.