



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 05:23 AM UTC

PDB ID : 2NVA / pdb_00002nva
Title : The X-ray crystal structure of the Paramecium bursaria Chlorella virus arginine decarboxylase bound to agmatine
Authors : Shah, R.H.; Akella, R.; Goldsmith, E.; Phillips, M.A.
Deposited on : 2006-11-11
Resolution : 1.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	 6% 91% 8% ..
1	B	372	 5% 92% 7% ..
1	C	372	 7% 87% 10% ..
1	D	372	 7% 89% 9% ..
1	E	372	 6% 88% 10% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	372	 5% 88% 10% ..
1	G	372	 8% 90% 9% .
1	H	372	 6% 88% 8% ..

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 25034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called arginine decarboxylase, A207R protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	369	2939	1884	480	562	13	0	1	0
1	B	369	2933	1880	478	562	13	0	1	0
1	C	367	2916	1867	475	561	13	0	1	0
1	D	371	2941	1884	480	564	13	0	1	0
1	E	370	2945	1886	479	567	13	0	1	0
1	F	368	2917	1869	476	559	13	0	1	0
1	G	368	2902	1858	475	556	13	0	1	0
1	H	364	2882	1845	469	555	13	0	1	0

- Molecule 2 is (4-[[4-[[AMINO(IMINO)METHYL]AMINO]BUTYL)AMINO]METHYL]-5-HYDROXY-6-METHYLPYRIDIN-3-YL)METHYL DIHYDROGEN PHOSPHATE (CCD ID: PL2) (formula: C₁₃H₂₄N₅O₅P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	B	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	C	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	D	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	E	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	F	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	G	1	Total	C	N	O	P	0	0
			24	13	5	5	1		
2	H	1	Total	C	N	O	P	0	0
			24	13	5	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	219	Total	O	0	0
			219	219		
3	B	193	Total	O	0	0
			193	193		
3	C	163	Total	O	0	0
			163	163		
3	D	173	Total	O	0	0
			173	173		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	211	Total 211	O 211	0	0
3	F	174	Total 174	O 174	0	0
3	G	149	Total 149	O 149	0	0
3	H	185	Total 185	O 185	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: arginine decarboxylase, A207R protein




- Molecule 1: arginine decarboxylase, A207R protein




- Molecule 1: arginine decarboxylase, A207R protein




- Molecule 1: arginine decarboxylase, A207R protein




- Molecule 1: arginine decarboxylase, A207R protein




- Molecule 1: arginine decarboxylase, A207R protein



- Molecule 1: arginine decarboxylase, A207R protein



- Molecule 1: arginine decarboxylase, A207R protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	116.21Å 117.28Å 268.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.74 – 1.80 44.74 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.74-1.80) 99.7 (44.74-1.80)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.26 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.219 , 0.238 0.233 , 0.257	Depositor DCC
R_{free} test set	2052 reflections (0.61%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	25034	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.93 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.2439e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PL2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/3014	0.76	0/4095
1	B	0.59	0/3005	0.73	0/4079
1	C	0.59	0/2987	0.77	0/4056
1	D	0.61	0/3015	0.80	2/4095 (0.0%)
1	E	0.59	0/3018	0.73	0/4099
1	F	0.60	0/2988	0.78	0/4055
1	G	0.60	0/2972	0.77	1/4033 (0.0%)
1	H	0.59	0/2951	0.74	0/4004
All	All	0.60	0/23950	0.76	3/32516 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	6	ASN	N-CA-C	13.32	125.52	111.14
1	G	86	ARG	CB-CA-C	-8.24	96.63	110.56
1	D	7	ASN	N-CA-C	7.63	127.06	110.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	6	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2939	0	2834	31	0
1	B	2933	0	2828	33	0
1	C	2916	0	2807	40	0
1	D	2941	0	2832	41	0
1	E	2945	0	2836	37	0
1	F	2917	0	2814	48	0
1	G	2902	0	2786	25	0
1	H	2882	0	2772	39	0
2	A	24	0	21	2	0
2	B	24	0	21	3	0
2	C	24	0	21	4	0
2	D	24	0	21	4	0
2	E	24	0	21	3	0
2	F	24	0	21	4	0
2	G	24	0	21	3	0
2	H	24	0	21	5	0
3	A	219	0	0	3	0
3	B	193	0	0	1	0
3	C	163	0	0	0	0
3	D	173	0	0	0	0
3	E	211	0	0	1	0
3	F	174	0	0	2	0
3	G	149	0	0	0	0
3	H	185	0	0	3	0
All	All	25034	0	22677	281	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 281 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:324[B]:CYS:SG	2:H:2008:PL2:HA2	1.65	1.33
2:G:2007:PL2:HA2	1:H:324[B]:CYS:SG	1.83	1.18
1:G:324[B]:CYS:SG	2:H:2008:PL2:CA	2.36	1.13
1:D:130:MET:HE2	1:D:130:MET:HA	1.30	1.11
1:H:121:ILE:HG13	1:H:130:MET:HE3	1.13	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/372 (98%)	357 (98%)	9 (2%)	0	100	100
1	B	362/372 (97%)	354 (98%)	8 (2%)	0	100	100
1	C	360/372 (97%)	350 (97%)	10 (3%)	0	100	100
1	D	368/372 (99%)	358 (97%)	9 (2%)	1 (0%)	36	25
1	E	365/372 (98%)	355 (97%)	10 (3%)	0	100	100
1	F	361/372 (97%)	352 (98%)	9 (2%)	0	100	100
1	G	359/372 (96%)	350 (98%)	9 (2%)	0	100	100
1	H	357/372 (96%)	348 (98%)	9 (2%)	0	100	100
All	All	2898/2976 (97%)	2824 (97%)	73 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	7	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	326/329 (99%)	320 (98%)	6 (2%)	51 43
1	B	325/329 (99%)	322 (99%)	3 (1%)	70 67
1	C	324/329 (98%)	316 (98%)	8 (2%)	42 30
1	D	325/329 (99%)	319 (98%)	6 (2%)	51 43
1	E	327/329 (99%)	319 (98%)	8 (2%)	43 31
1	F	323/329 (98%)	319 (99%)	4 (1%)	63 57
1	G	320/329 (97%)	316 (99%)	4 (1%)	61 54
1	H	319/329 (97%)	311 (98%)	8 (2%)	42 30
All	All	2589/2632 (98%)	2542 (98%)	47 (2%)	51 43

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	198	PHE
1	G	147	ASN
1	E	244	GLU
1	F	67	ASP
1	G	249	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 68 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	7	ASN
1	G	147	ASN
1	H	63	ASN
1	C	199	ASN
1	C	183	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PL2	G	2007	-	24,24,24	1.28	2 (8%)	30,32,32	1.75	6 (20%)
2	PL2	H	2008	-	24,24,24	1.29	3 (12%)	30,32,32	1.57	6 (20%)
2	PL2	C	2003	-	24,24,24	1.27	4 (16%)	30,32,32	1.65	7 (23%)
2	PL2	D	2004	-	24,24,24	1.18	3 (12%)	30,32,32	1.53	4 (13%)
2	PL2	F	2006	-	24,24,24	1.28	3 (12%)	30,32,32	1.61	6 (20%)
2	PL2	B	2002	-	24,24,24	1.24	3 (12%)	30,32,32	1.52	6 (20%)
2	PL2	E	2005	-	24,24,24	1.39	3 (12%)	30,32,32	1.69	6 (20%)
2	PL2	A	2001	-	24,24,24	1.37	4 (16%)	30,32,32	1.68	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PL2	G	2007	-	-	3/16/16/16	0/1/1/1
2	PL2	H	2008	-	-	6/16/16/16	0/1/1/1
2	PL2	C	2003	-	-	4/16/16/16	0/1/1/1
2	PL2	D	2004	-	-	7/16/16/16	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PL2	F	2006	-	-	3/16/16/16	0/1/1/1
2	PL2	B	2002	-	-	5/16/16/16	0/1/1/1
2	PL2	E	2005	-	-	3/16/16/16	0/1/1/1
2	PL2	A	2001	-	-	4/16/16/16	0/1/1/1

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2005	PL2	P-OP2	3.86	1.62	1.50
2	H	2008	PL2	C2A-C2	3.37	1.55	1.50
2	A	2001	PL2	P-OP2	3.37	1.60	1.50
2	G	2007	PL2	C2A-C2	3.29	1.55	1.50
2	F	2006	PL2	P-OP2	3.21	1.60	1.50

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	PL2	OP4-C5M-C5	5.38	119.44	109.36
2	H	2008	PL2	OP4-C5M-C5	4.80	118.36	109.36
2	E	2005	PL2	OP4-C5M-C5	4.41	117.62	109.36
2	G	2007	PL2	OP4-C5M-C5	4.40	117.61	109.36
2	G	2007	PL2	C6-C5-C4	4.31	121.32	118.06

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2004	PL2	C4-C5-C5M-OP4
2	H	2008	PL2	C3-C4-C4A-N
2	H	2008	PL2	C5-C4-C4A-N
2	A	2001	PL2	C5-C4-C4A-N
2	B	2002	PL2	C5-C4-C4A-N

There are no ring outliers.

8 monomers are involved in 28 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2007	PL2	3	0
2	H	2008	PL2	5	0
2	C	2003	PL2	4	0



Continued on next page...



Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2004	PL2	4	0
2	F	2006	PL2	4	0
2	B	2002	PL2	3	0
2	E	2005	PL2	3	0
2	A	2001	PL2	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/372 (99%)	0.48	23 (6%) 26 25	7, 13, 16, 17	1 (0%)
1	B	369/372 (99%)	0.42	19 (5%) 33 32	7, 13, 16, 18	1 (0%)
1	C	367/372 (98%)	0.45	25 (6%) 23 22	7, 14, 17, 19	1 (0%)
1	D	371/372 (99%)	0.46	25 (6%) 24 22	7, 14, 17, 19	1 (0%)
1	E	370/372 (99%)	0.40	21 (5%) 29 28	7, 13, 16, 18	1 (0%)
1	F	368/372 (98%)	0.49	19 (5%) 33 31	7, 14, 16, 19	1 (0%)
1	G	368/372 (98%)	0.55	31 (8%) 17 15	7, 14, 17, 20	1 (0%)
1	H	364/372 (97%)	0.38	23 (6%) 26 24	8, 14, 16, 19	1 (0%)
All	All	2946/2976 (98%)	0.45	186 (6%) 26 24	7, 14, 16, 20	8 (0%)

The worst 5 of 186 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	229	TYR	6.5
1	F	311	ASP	5.0
1	F	224	GLY	4.9
1	H	2	ASN	4.7
1	A	36	ILE	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PL2	F	2006	24/24	0.90	0.11	13,16,17,18	0
2	PL2	G	2007	24/24	0.90	0.12	14,17,18,18	0
2	PL2	H	2008	24/24	0.90	0.13	14,16,17,17	0
2	PL2	D	2004	24/24	0.91	0.11	15,16,17,18	0
2	PL2	A	2001	24/24	0.91	0.10	13,16,17,17	0
2	PL2	B	2002	24/24	0.91	0.11	13,16,17,17	0
2	PL2	C	2003	24/24	0.91	0.11	14,16,18,18	0
2	PL2	E	2005	24/24	0.92	0.10	12,16,17,17	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around PL2 H 2008:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around PL2 D 2004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PL2 A 2001:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PL2 C 2003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.