



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:18 AM UTC

PDB ID : 2NVD / pdb_00002nvd
Title : Human Aldose Reductase complexed with novel naphtho[1,2-d]isothiazole acetic acid derivative (2)
Authors : Steuber, H.; Heine, A.; Klebe, G.
Deposited on : 2006-11-12
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	15	0
			2575	1653	431	478	13			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ILE	LEU	SEE REMARK 999	UNP P15121

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 3 is 2-(CARBOXYMETHYL)-1-OXO-1,2-DIHYDRONAPHTHO[1,2-D]ISOTHIAZOLE-4-CARBOXYLIC ACID 3,3-DIOXIDE (CCD ID: ITB) (formula: $C_{14}H_9NO_7S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			23	14	1	7	1		
3	A	1	Total	C	N	O	S	0	0
			23	14	1	7	1		
3	A	1	Total	C	N	O	S	0	0
			23	14	1	7	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	214	Total	O	0	0
			214	214		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldose reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.39Å 66.64Å 47.14Å 90.00° 91.88° 90.00°	Depositor
Resolution (Å)	30.00 – 1.55 30.00 – 1.55	Depositor EDS
% Data completeness (in resolution range)	80.6 (30.00-1.55) 93.9 (30.00-1.55)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.10 (at 1.55Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.130 , 0.200 0.133 , 0.184	Depositor DCC
R_{free} test set	2087 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	13.0	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -l,k,h 0.027 for -h,-k,l 0.018 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2906	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, ITB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2688	1.49	30/3649 (0.8%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	182	ASN	CA-CB-CG	9.24	121.84	112.60
1	A	209	TYR	CA-CB-CG	9.16	130.40	113.90
1	A	283	GLN	CG-CD-OE1	8.86	138.52	120.80
1	A	187	HIS	CG-CD2-NE2	8.81	116.01	107.20
1	A	284	ASP	CA-CB-CG	8.27	120.87	112.60
1	A	307	LYS	CA-C-N	8.13	136.54	122.56
1	A	307	LYS	C-N-CA	8.13	136.54	122.56
1	A	283	GLN	CB-CG-CD	8.08	126.33	112.60
1	A	283	GLN	OE1-CD-NE2	-7.94	114.66	122.60
1	A	79	TRP	CA-C-N	7.84	131.56	120.28
1	A	79	TRP	C-N-CA	7.84	131.56	120.28
1	A	120	GLU	CA-C-N	7.67	133.73	120.68
1	A	120	GLU	C-N-CA	7.67	133.73	120.68
1	A	187	HIS	ND1-CG-CD2	-7.39	98.71	106.10
1	A	245	ALA	CA-C-N	7.02	130.26	120.29
1	A	245	ALA	C-N-CA	7.02	130.26	120.29
1	A	125	ASP	CA-CB-CG	6.92	119.52	112.60
1	A	245	ALA	O-C-N	-6.86	114.85	122.12
1	A	120	GLU	O-C-N	-6.79	115.10	123.04
1	A	283	GLN	CG-CD-NE2	-6.46	106.72	116.40
1	A	187	HIS	CE1-NE2-CD2	-6.37	102.63	109.00
1	A	132	PRO	CA-C-N	5.72	131.18	122.65
1	A	132	PRO	C-N-CA	5.72	131.18	122.65
1	A	7	ASN	CA-CB-CG	5.56	118.16	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	HIS	CB-CG-CD2	5.54	138.40	131.20
1	A	102	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	50	ASN	OD1-CG-ND2	-5.48	117.12	122.60
1	A	187	HIS	CB-CG-ND1	5.12	130.39	122.70
1	A	137	ILE	CA-C-O	-5.10	113.53	119.85
1	A	110	HIS	O-C-N	5.04	127.47	122.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2575	0	2583	25	0
2	A	48	0	25	3	0
3	A	69	0	21	2	0
4	A	214	0	0	4	0
All	All	2906	0	2629	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:GLU:HB3	3:A:800:ITB:O22	1.92	0.69
1:A:194:LYS:HD3	4:A:1217:HOH:O	1.94	0.68
1:A:221:LYS:HB3	1:A:223:GLU:OE1	2.00	0.60
1:A:185:GLU:OE2	1:A:187:HIS:HE1	1.86	0.59
1:A:244[B]:THR:HG22	4:A:1082:HOH:O	2.09	0.52
1:A:181:VAL:HA	1:A:205:VAL:O	2.10	0.52
1:A:68:LYS:HE2	1:A:71:GLU:OE2	2.11	0.51
1:A:84[B]:GLU:OE2	1:A:133:SER:OG	2.29	0.50
1:A:70:GLU:HG3	4:A:1186:HOH:O	2.11	0.50
1:A:219:TRP:O	1:A:221:LYS:HE3	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:GLU:OE2	1:A:229:GLU:HA	2.12	0.48
1:A:223:GLU:H	1:A:223:GLU:CD	2.22	0.47
1:A:136[A]:ASN:HD21	1:A:139:ASP:CG	2.25	0.45
1:A:222:PRO:HG2	1:A:223:GLU:OE2	2.17	0.44
1:A:301:LEU:O	1:A:304:THR:HG23	2.18	0.43
1:A:183:GLN:OE1	2:A:500:NAP:H2N	2.18	0.43
1:A:225:PRO:HG2	1:A:295:TRP:CE2	2.54	0.42
1:A:262:LYS:O	2:A:500:NAP:H8A	2.19	0.42
1:A:32:LYS:HE2	1:A:60:GLU:OE1	2.20	0.42
1:A:163[A]:HIS:ND1	1:A:315:PHE:O	2.49	0.42
1:A:164[B]:LEU:HD12	4:A:1175:HOH:O	2.19	0.42
2:A:500:NAP:C5N	3:A:600:ITB:H7	2.50	0.42
1:A:7:ASN:C	1:A:7:ASN:HD22	2.29	0.41
1:A:219:TRP:O	1:A:221:LYS:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/316 (104%)	326 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	294/281 (105%)	282 (96%)	12 (4%)	27 4

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	21	LYS
1	A	50	ASN
1	A	127	SER
1	A	136[A]	ASN
1	A	136[B]	ASN
1	A	138	LEU
1	A	194	LYS
1	A	209	TYR
1	A	221	LYS
1	A	300	LEU
1	A	313	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	26	GLN
1	A	50	ASN
1	A	129	ASN
1	A	182	ASN
1	A	187	HIS
1	A	200	GLN
1	A	283	GLN
1	A	292	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ITB	A	800	-	23,25,25	2.89	8 (34%)	31,39,39	2.91	8 (25%)
3	ITB	A	600	-	23,25,25	2.66	7 (30%)	31,39,39	4.73	18 (58%)
3	ITB	A	700	-	23,25,25	2.80	6 (26%)	31,39,39	2.42	9 (29%)
2	NAP	A	500	-	50,52,52	1.44	9 (18%)	71,80,80	1.75	16 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ITB	A	800	-	-	2/8/28/28	0/3/3/3
3	ITB	A	600	-	-	0/8/28/28	0/3/3/3
3	ITB	A	700	-	-	0/8/28/28	0/3/3/3
2	NAP	A	500	-	-	4/35/67/67	0/5/5/5

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	800	ITB	S16-N15	-11.34	1.52	1.67
3	A	700	ITB	S16-N15	-11.20	1.52	1.67
3	A	600	ITB	S16-N15	-9.69	1.54	1.67
3	A	600	ITB	O13-C11	3.98	1.34	1.22
3	A	800	ITB	O13-C11	3.64	1.33	1.22
3	A	700	ITB	O13-C11	3.48	1.32	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAP	P2B-O2B	3.32	1.65	1.59
3	A	600	ITB	O12-C11	-3.20	1.21	1.30
2	A	500	NAP	C6N-N1N	2.93	1.42	1.35
3	A	800	ITB	O18-S16	2.92	1.46	1.43
2	A	500	NAP	O4D-C1D	2.91	1.44	1.40
3	A	700	ITB	C10-C1	-2.75	1.38	1.43
3	A	600	ITB	C14-N15	-2.73	1.35	1.40
3	A	700	ITB	C14-N15	-2.67	1.35	1.40
2	A	500	NAP	C2A-N1A	2.52	1.38	1.33
3	A	700	ITB	O17-S16	2.49	1.46	1.43
3	A	600	ITB	O18-S16	2.48	1.46	1.43
3	A	800	ITB	C14-N15	-2.47	1.35	1.40
2	A	500	NAP	PA-O3	2.44	1.62	1.59
2	A	500	NAP	C4N-C3N	2.43	1.43	1.39
3	A	600	ITB	C10-C1	-2.42	1.39	1.43
3	A	800	ITB	O12-C11	-2.33	1.23	1.30
3	A	700	ITB	O12-C11	-2.29	1.23	1.30
3	A	600	ITB	C7-C6	-2.21	1.37	1.42
2	A	500	NAP	C3B-C2B	2.19	1.57	1.53
3	A	800	ITB	O17-S16	2.19	1.45	1.43
2	A	500	NAP	C2N-N1N	2.14	1.37	1.35
3	A	800	ITB	C7-C8	2.07	1.41	1.37
2	A	500	NAP	C2A-N3A	2.03	1.37	1.33
3	A	800	ITB	C10-C1	-2.02	1.40	1.43

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	600	ITB	C20-N15-S16	-11.37	114.24	121.49
3	A	600	ITB	C20-N15-C14	-10.12	115.51	122.66
3	A	600	ITB	O19-C14-C10	-9.70	117.28	129.21
3	A	600	ITB	C9-C10-C14	-9.66	105.42	111.68
3	A	800	ITB	C20-N15-C14	-8.75	116.48	122.66
3	A	800	ITB	C20-N15-S16	7.87	126.50	121.49
3	A	700	ITB	O19-C14-C10	-7.33	120.19	129.21
3	A	800	ITB	C9-C10-C14	-6.95	107.18	111.68
3	A	700	ITB	C9-C10-C14	-6.74	107.31	111.68
3	A	600	ITB	C8-C7-C6	6.41	132.68	121.12
2	A	500	NAP	C6A-C5A-C4A	5.92	125.27	117.18
3	A	600	ITB	O17-S16-N15	5.61	114.75	109.98
3	A	800	ITB	O19-C14-C10	-5.39	122.58	129.21
3	A	600	ITB	O12-C11-C8	5.32	130.42	115.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	NAP	C3N-C7N-N7N	-4.93	111.67	117.74
3	A	600	ITB	C14-N15-S16	4.89	118.08	115.13
3	A	600	ITB	O13-C11-C8	-4.71	110.72	121.97
3	A	600	ITB	C5-C6-C1	4.70	125.18	119.12
3	A	600	ITB	O19-C14-N15	4.40	130.38	123.45
3	A	600	ITB	C4-C5-C6	-4.13	114.74	120.48
3	A	600	ITB	C7-C6-C1	-3.96	114.09	119.26
3	A	600	ITB	O18-S16-N15	-3.93	106.64	109.98
2	A	500	NAP	C6N-N1N-C2N	3.48	124.84	121.88
2	A	500	NAP	C6A-C5A-N7A	-3.48	125.39	132.09
2	A	500	NAP	C2N-C3N-C4N	3.43	122.25	118.26
3	A	700	ITB	C20-N15-C14	-3.28	120.35	122.66
3	A	700	ITB	C2-C1-C6	3.10	121.93	117.92
3	A	700	ITB	C14-N15-S16	3.09	117.00	115.13
2	A	500	NAP	C5N-C4N-C3N	-2.71	117.70	120.36
3	A	700	ITB	C3-C2-C1	-2.66	117.33	120.91
3	A	800	ITB	C9-S16-N15	2.62	94.37	92.54
2	A	500	NAP	C5A-C4A-N3A	-2.59	123.14	126.72
3	A	700	ITB	C9-S16-N15	2.51	94.30	92.54
2	A	500	NAP	C5N-C6N-N1N	-2.51	116.96	120.38
3	A	600	ITB	O23-C21-O22	-2.46	117.00	123.33
3	A	800	ITB	O18-S16-N15	2.41	112.03	109.98
3	A	600	ITB	O23-C21-C20	2.34	122.61	113.38
2	A	500	NAP	C5A-C4A-N9A	2.32	108.34	105.81
3	A	700	ITB	O17-S16-O18	2.25	119.95	116.29
2	A	500	NAP	O3D-C3D-C4D	2.18	117.36	111.08
3	A	800	ITB	C14-N15-S16	2.17	116.44	115.13
2	A	500	NAP	O2A-PA-O3	2.17	113.14	107.27
3	A	800	ITB	O17-S16-O18	2.14	119.76	116.29
3	A	600	ITB	O17-S16-O18	2.12	119.73	116.29
2	A	500	NAP	C5A-C6A-N1A	-2.11	112.16	117.51
3	A	600	ITB	O12-C11-O13	-2.09	118.85	123.35
2	A	500	NAP	O3X-P2B-O2X	2.05	115.50	107.80
2	A	500	NAP	O2A-PA-O1A	2.04	121.91	112.44
2	A	500	NAP	O3X-P2B-O2B	-2.03	97.94	105.85
3	A	700	ITB	O19-C14-N15	2.01	126.61	123.45
2	A	500	NAP	O3-PA-O1A	-2.01	104.67	110.70

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	800	ITB	N15-C20-C21-O23
3	A	800	ITB	N15-C20-C21-O22
2	A	500	NAP	PA-O3-PN-O5D
2	A	500	NAP	C4D-C5D-O5D-PN
2	A	500	NAP	C2B-O2B-P2B-O1X
2	A	500	NAP	PN-O3-PA-O1A


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	800	ITB	1	0
3	A	600	ITB	1	0
2	A	500	NAP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand ITB A 800




Bond lengths



Bond angles




Torsions




Rings


Ligand ITB A 600




Bond lengths



Bond angles




Torsions




Rings


Ligand ITB A 700




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	315/316 (99%)	-0.28	8 (2%) 58 66	7, 12, 34, 67	15 (4%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	PHE	3.6
1	A	314	GLU	3.1
1	A	219	TRP	3.0
1	A	222	PRO	2.6
1	A	1	ALA	2.6
1	A	194	LYS	2.3
1	A	223	GLU	2.2
1	A	313	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ITB	A	800	23/23	0.89	0.11	24,34,41,63	0
3	ITB	A	600	23/23	0.96	0.07	11,15,30,32	0
3	ITB	A	700	23/23	0.97	0.06	18,25,32,41	0
2	NAP	A	500	48/48	0.98	0.04	6,9,13,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ITB A 800:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ITB A 600:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ITB A 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.