



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:04 PM UTC

PDB ID : 4NVQ / pdb_00004nvq
Title : Human G9a in Complex with Inhibitor A-366
Authors : Sweis, R.F.; Pliushchev, M.; Brown, P.J.; Guo, J.; Li, F.; Maag, D.; Petros, A.M.; Soni, N.B.; Tse, C.; Vedadi, M.; Michaelides, M.R.; Chiang, G.G.; Pappano, W.N.
Deposited on : 2013-12-05
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	
1	B	285	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histone-lysine N-methyltransferase EHMT2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2048	1276	360	389	23			
1	B	263	Total	C	N	O	S	0	0	0
			2025	1263	353	386	23			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	909	GLY	-	expression tag	UNP Q96KQ7
A	910	SER	-	expression tag	UNP Q96KQ7
A	911	HIS	-	expression tag	UNP Q96KQ7
A	912	MET	-	expression tag	UNP Q96KQ7
B	909	GLY	-	expression tag	UNP Q96KQ7
B	910	SER	-	expression tag	UNP Q96KQ7
B	911	HIS	-	expression tag	UNP Q96KQ7
B	912	MET	-	expression tag	UNP Q96KQ7

- Molecule 2 is 5'-methoxy-6'-[3-(pyrrolidin-1-yl)propoxy]spiro[cyclobutane-1,3'-indol]-2'-amine (CCD ID: 2OD) (formula: C₁₉H₂₇N₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			24	19	3	2		
2	B	1	Total	C	N	O	0	0
			24	19	3	2		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total 4	Zn 4	0	0
4	B	4	Total 4	Zn 4	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	139	Total 139	O 139	0	0
5	B	104	Total 104	O 104	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-lysine N-methyltransferase EHMT2



- Molecule 1: Histone-lysine N-methyltransferase EHMT2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.14Å 78.25Å 71.44Å 90.00° 92.61° 90.00°	Depositor
Resolution (Å)	45.59 – 2.03 45.59 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.3 (45.59-2.03) 99.3 (45.59-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.03Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.11.2, BUSTER 2.11.2	Depositor
R, R_{free}	0.202 , 0.239 0.209 , 0.253	Depositor DCC
R_{free} test set	1994 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.641	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 65.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4424	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.85 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7143e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, ZN, 2OD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/2091 (0.0%)	1.18	7/2837 (0.2%)
1	B	0.85	0/2069	1.19	9/2804 (0.3%)
All	All	0.87	1/4160 (0.0%)	1.18	16/5641 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	962	MET	SD-CE	-6.19	1.64	1.79

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1178	SER	N-CA-C	12.21	126.11	111.33
1	B	1177	HIS	CA-C-N	6.65	129.49	120.38
1	B	1177	HIS	C-N-CA	6.65	129.49	120.38
1	A	978	ASP	CA-CB-CG	5.83	118.43	112.60
1	A	1162	LYS	N-CA-C	5.74	118.75	111.69
1	B	1178	SER	CA-C-N	5.69	132.40	121.54
1	B	1178	SER	C-N-CA	5.69	132.40	121.54
1	A	1179	ALA	CA-C-N	5.64	127.84	120.28
1	A	1179	ALA	C-N-CA	5.64	127.84	120.28
1	A	972	GLN	N-CA-C	-5.50	101.01	109.76
1	B	1099	ILE	N-CA-C	-5.47	100.52	108.17
1	A	1031	VAL	N-CA-C	5.35	115.49	110.30
1	A	997	ASP	CA-CB-CG	5.30	117.90	112.60
1	B	997	ASP	CA-CB-CG	5.15	117.75	112.60
1	B	1162	LYS	N-CA-C	5.15	118.72	112.23
1	B	1131	LEU	N-CA-C	5.06	118.94	112.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2048	0	1827	16	0
1	B	2025	0	1796	4	0
2	A	24	0	27	1	0
2	B	24	0	27	0	0
3	A	26	0	19	1	0
3	B	26	0	19	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	139	0	0	14	0
5	B	104	0	0	2	0
All	All	4424	0	3715	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:980:CYS:SG	5:A:1425:HOH:O	2.11	1.07
1:A:1027:CYS:HB2	5:A:1425:HOH:O	1.56	1.06
1:A:1115:CYS:HB2	5:A:1398:HOH:O	1.75	0.85
1:A:1170:CYS:SG	5:A:1398:HOH:O	2.40	0.80
1:A:1172:SER:HB3	5:A:1398:HOH:O	1.83	0.76
1:A:1159:TRP:O	5:A:1399:HOH:O	2.04	0.75
1:A:1162:LYS:N	5:A:1399:HOH:O	2.19	0.73
1:A:1159:TRP:C	5:A:1399:HOH:O	2.44	0.61
1:B:1096:VAL:N	5:B:1399:HOH:O	2.34	0.60
2:A:1201:2OD:N3	5:A:1433:HOH:O	2.16	0.59
1:B:923:CYS:HB2	1:B:946:CYS:SG	2.43	0.58
1:A:1077:ALA:HB3	5:A:1433:HOH:O	2.07	0.55
1:B:955:GLU:HG2	5:B:1383:HOH:O	2.07	0.55
1:A:980:CYS:CB	5:A:1425:HOH:O	2.46	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1027:CYS:CB	5:A:1425:HOH:O	2.34	0.49
1:A:979:ASP:C	5:A:1425:HOH:O	2.58	0.47
1:A:1048:MET:O	3:A:1202:SAH:N	2.48	0.46
1:A:1163:SER:O	5:A:1436:HOH:O	2.21	0.45
1:A:1021:CYS:O	1:B:951:LYS:NZ	2.48	0.44
1:A:1162:LYS:HD2	1:A:1166:PHE:CE2	2.52	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/285 (92%)	251 (96%)	8 (3%)	2 (1%)	16	8
1	B	257/285 (90%)	247 (96%)	9 (4%)	1 (0%)	30	22
All	All	518/570 (91%)	498 (96%)	17 (3%)	3 (1%)	21	12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1179	ALA
1	A	1095	GLU
1	A	992	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/255 (82%)	208 (100%)	1 (0%)	81	81
1	B	206/255 (81%)	200 (97%)	6 (3%)	37	34
All	All	415/510 (81%)	408 (98%)	7 (2%)	53	53

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	993	ARG
1	B	946	CYS
1	B	961	THR
1	B	964	ILE
1	B	993	ARG
1	B	1090	ASP
1	B	1165	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1169	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2OD	A	1201	-	26,27,27	1.53	5 (19%)	31,39,39	1.08	3 (9%)
2	2OD	B	1202	-	26,27,27	1.21	2 (7%)	31,39,39	1.19	4 (12%)
3	SAH	A	1202	-	27,28,28	0.79	2 (7%)	36,40,40	0.63	1 (2%)
3	SAH	B	1201	-	27,28,28	0.36	0	36,40,40	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2OD	A	1201	-	-	0/9/40/40	0/4/4/4
2	2OD	B	1202	-	-	3/9/40/40	0/4/4/4
3	SAH	A	1202	-	-	1/15/31/31	0/3/3/3
3	SAH	B	1201	-	-	2/15/31/31	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	2OD	C5-C4	3.31	1.44	1.39
2	A	1201	2OD	C6-C7	3.20	1.44	1.38
2	A	1201	2OD	C2-C7	3.03	1.47	1.40
3	A	1202	SAH	O-C	2.88	1.30	1.22
2	A	1201	2OD	C3-C2	2.84	1.43	1.38
3	A	1202	SAH	OXT-C	-2.61	1.22	1.30
2	B	1202	2OD	C15-N3	2.19	1.36	1.33
2	B	1202	2OD	O1-C2	2.16	1.40	1.37
2	A	1201	2OD	C14-N1	2.13	1.51	1.47

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1202	2OD	O2-C7-C6	-3.06	116.89	123.49
2	A	1201	2OD	C1-O1-C2	2.64	121.38	117.51
2	B	1202	2OD	O2-C7-C2	2.64	121.20	115.75
3	A	1202	SAH	CB-CA-N	2.40	116.37	110.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	2OD	O2-C7-C2	2.31	120.52	115.75
2	A	1201	2OD	O2-C7-C6	-2.24	118.65	123.49
2	B	1202	2OD	C1-O1-C2	2.17	120.70	117.51
2	B	1202	2OD	C9-C10-N1	-2.00	108.30	113.93

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	B	1202	2OD	N1-C10-C9-C8
2	B	1202	2OD	C2-C7-O2-C8
2	B	1202	2OD	C6-C7-O2-C8
3	B	1201	SAH	CB-CG-SD-C5'
3	A	1202	SAH	CB-CG-SD-C5'
3	B	1201	SAH	C-CA-CB-CG


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1201	2OD	1	0
3	A	1202	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/285 (93%)	0.32	6 (2%) 62 62	38, 55, 80, 105	0
1	B	263/285 (92%)	0.42	16 (6%) 27 26	39, 57, 88, 109	0
All	All	530/570 (92%)	0.37	22 (4%) 40 40	38, 56, 85, 109	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1094	GLY	4.5
1	B	1190	ALA	3.2
1	A	916	ILE	3.2
1	B	918	THR	3.1
1	B	921	ILE	2.8
1	B	922	ILE	2.7
1	B	995	TRP	2.7
1	B	1006	PHE	2.6
1	B	1013	LEU	2.5
1	B	946	CYS	2.5
1	B	1090	ASP	2.5
1	B	1072	ILE	2.5
1	B	987	CYS	2.4
1	A	1187	SER	2.4
1	B	1165	TYR	2.3
1	B	977	VAL	2.3
1	B	986	LEU	2.3
1	A	984	ASN	2.2
1	A	1005	GLU	2.2
1	B	948	GLU	2.2
1	B	1182	ILE	2.1
1	A	1058	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	2OD	B	1202	24/24	0.91	0.10	46,62,65,68	0
2	2OD	A	1201	24/24	0.93	0.08	49,56,63,65	0
3	SAH	A	1202	26/26	0.93	0.07	44,52,56,57	0
3	SAH	B	1201	26/26	0.93	0.08	48,56,59,61	0
4	ZN	A	1203	1/1	0.97	0.04	55,55,55,55	0
4	ZN	A	1204	1/1	0.97	0.07	58,58,58,58	0
4	ZN	A	1205	1/1	0.98	0.06	59,59,59,59	0
4	ZN	A	1206	1/1	0.98	0.04	61,61,61,61	0
4	ZN	B	1205	1/1	0.99	0.04	48,48,48,48	0
4	ZN	B	1206	1/1	0.99	0.03	60,60,60,60	0
4	ZN	B	1203	1/1	1.00	0.02	44,44,44,44	0
4	ZN	B	1204	1/1	1.00	0.03	48,48,48,48	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2OD B 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2OD A 1201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SAH A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAH B 1201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.