



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:49 AM UTC

PDB ID : 1NWH / pdb_00001nwh
Title : Crystal Structure of Aspartate Semialdehyde Dehydrogenase from Haemophilus influenzae as a Tetrahedral Hemithioacetal Reaction Intermediate at 2.0 Å
Authors : Blanco, J.; Moore, R.A.; Viola, R.E.
Deposited on : 2003-02-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	371	84% 12% .
1	B	371	84% 11% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5847 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate-semialdehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	0	0
			2757	1759	463	520	15			
1	B	356	Total	C	N	O	S	0	0	0
			2756	1758	463	520	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	136	HTI	CYS	modified residue	UNP P44801
B	136	HTI	CYS	modified residue	UNP P44801


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	176	Total	O	0	0
			176	176		
2	B	158	Total	O	0	0
			158	158		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate-semialdehyde dehydrogenase



- Molecule 1: Aspartate-semialdehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.56Å 111.04Å 113.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.58 – 2.00 29.58 – 2.00	Depositor EDS
% Data completeness (in resolution range)	82.2 (29.58-2.00) 82.2 (29.58-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.43 (at 2.00Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.202 , 0.262 0.206 , 0.261	Depositor DCC
R_{free} test set	3919 reflections (9.79%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5847	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.93 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2298e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HTI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/2791	1.17	22/3773 (0.6%)
1	B	0.71	1/2789 (0.0%)	1.16	21/3769 (0.6%)
All	All	0.71	1/5580 (0.0%)	1.16	43/7542 (0.6%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	324	VAL	CA-CB	5.02	1.60	1.54

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	111	VAL	N-CA-C	8.10	119.95	108.36
1	A	111	VAL	N-CA-C	7.83	119.56	108.36
1	B	261	PRO	N-CA-C	7.82	123.33	111.14
1	B	217	ASP	N-CA-C	6.70	119.42	111.71
1	B	274	LEU	N-CA-C	6.41	119.22	111.40
1	B	249	ALA	N-CA-C	6.35	118.00	111.14
1	A	318	GLU	N-CA-C	6.29	121.48	113.55
1	B	324	VAL	N-CA-C	6.26	121.58	113.07
1	B	312	LYS	N-CA-C	6.17	118.55	111.02
1	A	217	ASP	N-CA-C	6.12	117.61	111.07
1	B	359	ALA	N-CA-C	6.10	119.92	112.23
1	A	113	ASP	N-CA-C	6.10	123.28	109.81
1	B	100	SER	N-CA-C	6.07	119.95	112.54
1	A	82	VAL	N-CA-C	6.06	117.60	111.00
1	B	145	ILE	N-CA-C	6.01	117.88	112.17
1	B	113	ASP	N-CA-C	5.97	123.00	109.81
1	A	241	THR	N-CA-C	-5.94	101.98	110.59
1	B	163	GLN	N-CA-C	5.93	119.93	109.96

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	PRO	N-CA-C	5.88	120.56	111.21
1	A	163	GLN	N-CA-C	5.87	119.82	109.96
1	B	4	VAL	N-CA-C	5.86	116.38	108.17
1	A	359	ALA	N-CA-C	5.77	119.50	112.23
1	B	159	VAL	N-CA-C	5.76	116.88	108.53
1	B	241	THR	N-CA-C	-5.68	101.61	110.42
1	B	318	GLU	N-CA-C	5.68	120.70	113.55
1	B	223	LEU	N-CA-C	-5.63	106.33	113.43
1	A	250	GLU	N-CA-C	5.60	118.58	111.69
1	B	301	HIS	N-CA-C	5.55	117.00	111.07
1	A	301	HIS	N-CA-C	5.52	116.98	111.07
1	B	101	ALA	N-CA-C	5.50	120.44	111.37
1	A	103	ARG	N-CA-C	5.38	117.56	111.11
1	A	249	ALA	N-CA-C	5.32	116.77	111.07
1	A	275	ARG	N-CA-C	5.31	120.56	112.54
1	A	116	ASN	N-CA-C	5.25	118.81	111.30
1	A	324	VAL	N-CA-C	5.24	120.25	113.07
1	A	101	ALA	N-CA-C	5.17	119.90	111.37
1	B	103	ARG	N-CA-C	5.13	117.54	111.33
1	A	299	ALA	N-CA-C	5.13	116.87	111.28
1	B	276	CYS	N-CA-C	5.08	117.25	109.07
1	A	169	GLY	N-CA-C	5.04	116.96	110.96
1	A	4	VAL	N-CA-C	5.02	115.20	108.17
1	A	282	THR	N-CA-C	-5.01	97.22	107.70
1	A	312	LYS	N-CA-C	5.01	116.43	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2757	0	2814	9	0
1	B	2756	0	2809	11	0
2	A	176	0	0	0	0
2	B	158	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5847	0	5623	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:CA	1:B:1:MET:CG	2.53	0.86
1:A:135:ASN:HD22	1:A:138:VAL:HG23	1.54	0.73
1:A:20:ARG:HH11	1:A:23:GLN:NE2	1.97	0.62
1:A:183:LEU:HD22	1:A:205:VAL:HG23	1.85	0.58
1:B:135:ASN:HD22	1:B:138:VAL:HG23	1.68	0.58
1:A:20:ARG:HH11	1:A:23:GLN:HE21	1.56	0.53
1:B:336:ARG:NH1	1:B:338:LEU:HD23	2.24	0.53
1:B:336:ARG:HH12	1:B:338:LEU:HD23	1.75	0.51
1:A:162:TYR:HB2	1:A:278:SER:HB2	1.93	0.50
1:A:136:HTI:OAO	1:A:270:ARG:NH2	2.41	0.49
1:B:25:ASN:O	1:B:28:GLU:HG2	2.13	0.48
1:B:293:GLU:O	1:B:297:ILE:HG12	2.15	0.47
1:B:162:TYR:HB2	1:B:278:SER:HB2	2.00	0.43
1:B:166:SER:HB2	1:B:274:LEU:O	2.19	0.43
1:A:320:THR:O	1:A:324:VAL:HG13	2.19	0.43
1:B:159:VAL:O	1:B:266:GLY:HA3	2.20	0.42
1:A:20:ARG:NH1	1:A:23:GLN:HE21	2.17	0.41
1:B:111:VAL:HG11	1:B:365:ILE:HB	2.02	0.41
1:A:97:ASP:HB3	1:A:132:VAL:HG13	2.02	0.41
1:B:148:LEU:HD21	1:B:294:ILE:HG23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/371 (95%)	342 (97%)	9 (3%)	0	100	100
1	B	351/371 (95%)	340 (97%)	11 (3%)	0	100	100
All	All	702/742 (95%)	682 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/304 (97%)	284 (96%)	12 (4%)	27	26
1	B	295/304 (97%)	285 (97%)	10 (3%)	32	33
All	All	591/608 (97%)	569 (96%)	22 (4%)	30	30

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	137	THR
1	A	153	LEU
1	A	192	LYS
1	A	202	GLU
1	A	205	VAL
1	A	234	LEU
1	A	235	LEU
1	A	287	LYS
1	A	291	LEU
1	A	321	PRO
1	A	324	VAL
1	A	363	ARG
1	B	137	THR
1	B	145	ILE
1	B	182	LEU
1	B	202	GLU
1	B	235	LEU
1	B	289	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	321	PRO
1	B	324	VAL
1	B	363	ARG
1	B	366	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	23	GLN
1	A	117	GLN
1	A	135	ASN
1	A	279	GLN
1	A	296	GLN
1	A	368	GLN
1	B	29	ASN
1	B	74	GLN
1	B	117	GLN
1	B	135	ASN
1	B	172	ASN
1	B	279	GLN
1	B	353	GLN
1	B	368	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	HTI	B	136	1	7,13,14	1.11	0	5,16,18	0.75	0
1	HTI	A	136	1	7,13,14	1.17	0	5,16,18	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HTI	B	136	1	-	3/11/15/17	-
1	HTI	A	136	1	-	2/11/15/17	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	136	HTI	CAH-CAJ-CAK-NAL
1	B	136	HTI	CAH-CAJ-CAK-NAL
1	B	136	HTI	CAH-CAJ-CAK-CAM
1	A	136	HTI	CAH-CAJ-CAK-CAM
1	B	136	HTI	NAL-CAK-CAM-OAO

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	136	HTI	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/371 (95%)	0.02	2 (0%) 85 85	8, 18, 30, 40	0
1	B	355/371 (95%)	0.16	9 (2%) 58 58	9, 19, 33, 46	0
All	All	710/742 (95%)	0.09	11 (1%) 72 71	8, 19, 32, 46	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	260	ASN	3.1
1	A	92	ASP	3.0
1	B	248	TYR	2.9
1	B	224	GLY	2.9
1	B	258	SER	2.9
1	B	298	ILE	2.4
1	B	193	ASP	2.3
1	B	261	PRO	2.3
1	B	55	LEU	2.2
1	B	185	GLN	2.1
1	A	55	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	HTI	B	136	14/15	0.93	0.10	16,24,28,28	0
1	HTI	A	136	14/15	0.94	0.09	14,23,25,26	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.