



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 06:14 PM UTC

PDB ID : 3NWA / pdb_00003nwa
Title : Glycoprotein B from Herpes simplex virus type 1, W174R mutant, low-pH
Authors : Stampfer, S.D.; Lou, H.; Cohen, G.H.; Eisenberg, R.J.; Heldwein, E.E.
Deposited on : 2010-07-09
Resolution : 2.26 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	703	
1	B	703	
1	C	703	
1	D	703	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 21541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	607	Total	C	N	O	S	0	1	0
			4896	3086	867	920	23			
1	A	608	Total	C	N	O	S	0	1	0
			4899	3088	867	922	22			
1	C	608	Total	C	N	O	S	0	0	0
			4883	3080	864	917	22			
1	D	607	Total	C	N	O	S	0	1	0
			4900	3087	867	923	23			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	28	ASP	-	expression tag	UNP P06437
B	29	PRO	-	expression tag	UNP P06437
B	58	ALA	PRO	SEE REMARK 999	UNP P06437
B	174	ARG	TRP	engineered mutation	UNP P06437
B	313	SER	THR	SEE REMARK 999	UNP P06437
B	443	LEU	GLN	SEE REMARK 999	UNP P06437
A	28	ASP	-	expression tag	UNP P06437
A	29	PRO	-	expression tag	UNP P06437
A	58	ALA	PRO	SEE REMARK 999	UNP P06437
A	174	ARG	TRP	engineered mutation	UNP P06437
A	313	SER	THR	SEE REMARK 999	UNP P06437
A	443	LEU	GLN	SEE REMARK 999	UNP P06437
C	28	ASP	-	expression tag	UNP P06437
C	29	PRO	-	expression tag	UNP P06437
C	58	ALA	PRO	SEE REMARK 999	UNP P06437
C	174	ARG	TRP	engineered mutation	UNP P06437
C	313	SER	THR	SEE REMARK 999	UNP P06437
C	443	LEU	GLN	SEE REMARK 999	UNP P06437
D	28	ASP	-	expression tag	UNP P06437
D	29	PRO	-	expression tag	UNP P06437
D	58	ALA	PRO	SEE REMARK 999	UNP P06437

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	174	ARG	TRP	engineered mutation	UNP P06437
D	313	SER	THR	SEE REMARK 999	UNP P06437
D	443	LEU	GLN	SEE REMARK 999	UNP P06437

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is MESO-ERYTHRITOL (CCD ID: MRY) (formula: $C_4H_{10}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			8	4	4		
3	B	1	Total	C	O	0	0
			8	4	4		
3	A	1	Total	C	O	0	0
			8	4	4		
3	C	1	Total	C	O	0	0
			8	4	4		
3	D	1	Total	C	O	0	0
			8	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	444	Total	O	0	0
			444	444		
4	A	478	Total	O	0	0
			478	478		
4	C	462	Total	O	0	0
			462	462		
4	D	483	Total	O	0	0
			483	483		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Envelope glycoprotein B



• Molecule 1: Envelope glycoprotein B





4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	117.09Å 117.09Å 321.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.32 – 2.26 47.32 – 2.26	Depositor EDS
% Data completeness (in resolution range)	87.1 (47.32-2.26) 87.1 (47.32-2.26)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.27Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.172 , 0.227 0.171 , 0.227	Depositor DCC
R_{free} test set	11385 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.493	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for -h,-k,l 0.467 for h,-h-k,-l 0.021 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	21541	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4096e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MRY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/5021	0.79	2/6822 (0.0%)
1	B	0.45	0/5017	0.82	4/6814 (0.1%)
1	C	0.45	0/5002	0.79	4/6798 (0.1%)
1	D	0.46	0/5020	0.80	6/6820 (0.1%)
All	All	0.46	0/20060	0.80	16/27254 (0.1%)

There are no bond length outliers.

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	357	VAL	CA-C-N	6.44	126.39	119.76
1	B	357	VAL	C-N-CA	6.44	126.39	119.76
1	D	116	CYS	CA-C-N	6.43	127.00	120.38
1	D	116	CYS	C-N-CA	6.43	127.00	120.38
1	A	104	ILE	N-CA-C	6.29	117.55	107.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4899	0	4713	141	0
1	B	4896	0	4717	190	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	4883	0	4691	147	0
1	D	4900	0	4708	151	0
2	A	14	0	13	0	0
2	B	14	0	13	1	0
2	C	14	0	13	1	0
2	D	14	0	13	1	0
3	A	8	0	10	0	0
3	B	16	0	20	0	0
3	C	8	0	10	0	0
3	D	8	0	10	0	0
4	A	478	0	0	13	0
4	B	444	0	0	12	0
4	C	462	0	0	19	0
4	D	483	0	0	14	0
All	All	21541	0	18931	625	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 625 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:515:ARG:HG3	1:D:515:ARG:HH11	1.27	0.98
1:C:543:ARG:HB3	1:C:550:ILE:HG21	1.55	0.88
1:D:515:ARG:HG3	1:D:515:ARG:NH1	1.82	0.88
1:B:189:ARG:HB2	1:B:349:LYS:HE2	1.59	0.85
1:D:403:PRO:HG3	1:D:476:PRO:HB3	1.60	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	605/703 (86%)	558 (92%)	43 (7%)	4 (1%)	18	17
1	B	604/703 (86%)	555 (92%)	46 (8%)	3 (0%)	24	24
1	C	604/703 (86%)	543 (90%)	55 (9%)	6 (1%)	12	10
1	D	604/703 (86%)	571 (94%)	29 (5%)	4 (1%)	18	17
All	All	2417/2812 (86%)	2227 (92%)	173 (7%)	17 (1%)	18	17

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	259	VAL
1	A	260	GLU
1	D	177	HIS
1	C	118	PRO
1	C	655	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	526/593 (89%)	502 (95%)	24 (5%)	24	28
1	B	526/593 (89%)	498 (95%)	28 (5%)	20	22
1	C	522/593 (88%)	503 (96%)	19 (4%)	31	39
1	D	527/593 (89%)	501 (95%)	26 (5%)	22	24
All	All	2101/2372 (89%)	2004 (95%)	97 (5%)	24	28

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	363	VAL
1	D	180	SER
1	C	432	THR
1	C	529	CYS
1	D	305	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	440	GLN
1	D	681	HIS
1	D	706	GLN
1	D	674	ASN
1	C	709	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MRY	D	2000	-	7,7,7	0.50	0	8,8,8	0.84	0
2	NAG	C	1141	1	14,14,15	0.57	0	17,19,21	1.10	1 (5%)
3	MRY	A	2000	-	7,7,7	0.52	0	8,8,8	0.69	0
3	MRY	B	3000	-	7,7,7	0.93	0	8,8,8	1.43	0
2	NAG	D	1398	1	14,14,15	0.57	0	17,19,21	0.96	1 (5%)
3	MRY	B	2000	-	7,7,7	0.38	0	8,8,8	0.79	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1398	1	14,14,15	0.54	0	17,19,21	1.41	1 (5%)
2	NAG	B	1398	1	14,14,15	0.47	0	17,19,21	0.99	1 (5%)
3	MRY	C	2000	-	7,7,7	0.22	0	8,8,8	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MRY	D	2000	-	-	0/8/8/8	-
2	NAG	C	1141	1	-	2/6/23/26	0/1/1/1
3	MRY	A	2000	-	-	0/8/8/8	-
3	MRY	B	3000	-	-	1/8/8/8	-
2	NAG	D	1398	1	-	2/6/23/26	0/1/1/1
3	MRY	B	2000	-	-	0/8/8/8	-
2	NAG	A	1398	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1398	1	-	2/6/23/26	0/1/1/1
3	MRY	C	2000	-	-	0/8/8/8	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1398	NAG	C1-O5-C5	4.77	118.58	112.19
2	B	1398	NAG	C1-O5-C5	3.17	116.44	112.19
2	C	1141	NAG	O5-C1-C2	2.62	115.34	111.29
2	D	1398	NAG	C1-O5-C5	2.08	114.97	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1141	NAG	O5-C5-C6-O6
2	B	1398	NAG	O5-C5-C6-O6
2	B	1398	NAG	C4-C5-C6-O6
2	D	1398	NAG	C8-C7-N2-C2
2	D	1398	NAG	O7-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1141	NAG	1	0
2	D	1398	NAG	1	0
2	B	1398	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	608/703 (86%)	-1.63	0 100 100	21, 33, 55, 75	1 (0%)
1	B	607/703 (86%)	-1.57	0 100 100	19, 36, 63, 90	1 (0%)
1	C	608/703 (86%)	-1.59	0 100 100	22, 35, 64, 88	0
1	D	607/703 (86%)	-1.61	0 100 100	15, 34, 55, 88	1 (0%)
All	All	2430/2812 (86%)	-1.60	0 100 100	15, 34, 60, 90	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1398	14/15	0.97	0.06	49,65,76,81	0
2	NAG	C	1141	14/15	0.97	0.05	84,89,92,95	0
2	NAG	D	1398	14/15	0.98	0.04	40,58,64,64	0
2	NAG	B	1398	14/15	0.99	0.03	46,59,65,66	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MRY	B	2000	8/8	0.99	0.03	29,35,39,44	0
3	MRY	B	3000	8/8	0.99	0.03	39,41,43,47	0
3	MRY	A	2000	8/8	0.99	0.04	27,33,36,37	0
3	MRY	C	2000	8/8	0.99	0.04	27,29,33,38	0
3	MRY	D	2000	8/8	1.00	0.03	27,27,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.