



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:55 PM UTC

PDB ID : 4NWG / pdb_00004nwg
Title : Crystal structure of the tyrosine phosphatase SHP-2 with E139D mutation
Authors : Qiu, W.; Lin, A.; Hutchinson, A.; Romanov, V.; Ruzanov, M.; Thompson, C.;
Lam, K.; Kisselman, G.; Battalie, K.; Chirgadze, N.Y.
Deposited on : 2013-12-06
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	539	19% 84% 8% 8%
1	B	539	15% 85% 6% 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	498	Total	C	N	O	S	0	2	0
			4044	2541	724	760	19			
1	B	492	Total	C	N	O	S	0	1	0
			3994	2510	712	753	19			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	ASP	GLU	engineered mutation	UNP Q06124
A	?	-	GLN	deletion	UNP Q06124
A	?	-	ALA	deletion	UNP Q06124
A	?	-	LEU	deletion	UNP Q06124
A	?	-	LEU	deletion	UNP Q06124
B	139	ASP	GLU	engineered mutation	UNP Q06124
B	?	-	GLN	deletion	UNP Q06124
B	?	-	ALA	deletion	UNP Q06124
B	?	-	LEU	deletion	UNP Q06124
B	?	-	LEU	deletion	UNP Q06124

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	181	Total 181	O 181	0	0
5	B	181	Total 181	O 181	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.32Å 212.42Å 92.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.64 – 2.45 38.64 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.64-2.45) 99.9 (38.64-2.45)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.45Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.280 , 0.326 0.275 , 0.320	Depositor DCC
R_{free} test set	1070 reflections (2.58%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.410	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.40$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8456	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.09 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.7478e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/4131	1.24	6/5569 (0.1%)
1	B	0.68	0/4074	1.22	2/5495 (0.0%)
All	All	0.68	0/8205	1.23	8/11064 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	HIS	N-CA-C	6.26	124.14	110.80
1	A	411	THR	N-CA-C	-6.23	105.36	114.39
1	A	506	GLN	N-CA-C	5.81	118.48	111.40
1	B	506	GLN	N-CA-C	5.76	118.43	111.40
1	A	85	HIS	N-CA-C	5.60	117.47	111.36
1	A	117	LEU	N-CA-C	-5.54	102.28	110.48
1	B	130	GLY	N-CA-C	5.18	118.03	112.33
1	A	130	GLY	N-CA-C	5.09	117.93	112.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4044	0	3973	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3994	0	3908	12	0
2	A	28	0	42	5	0
2	B	16	0	24	2	0
3	A	7	0	10	1	0
4	B	5	0	0	0	0
5	A	181	0	0	0	0
5	B	181	0	0	0	0
All	All	8456	0	7957	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:405:LYS:H	1:A:411:THR:CG2	2.07	0.67
1:A:73:THR:HG21	2:A:604:EDO:H11	1.78	0.66
1:A:405:LYS:H	1:A:411:THR:HG21	1.68	0.57
1:A:133:GLY:HA3	1:A:213:LYS:HG2	1.88	0.55
1:B:69:GLU:HB3	2:B:605:EDO:H21	1.88	0.55
1:B:280:LYS:HG2	2:B:605:EDO:H22	1.89	0.54
1:B:409:GLY:C	1:B:411:THR:H	2.15	0.53
1:A:280:LYS:HG2	2:A:607:EDO:H11	1.91	0.53
1:A:69:GLU:HB3	2:A:607:EDO:H12	1.90	0.53
1:A:25:VAL:HG11	1:B:175:GLN:HB3	1.91	0.52
1:A:175:GLN:HB3	1:B:25:VAL:HG11	1.92	0.51
1:A:65:LEU:HD23	1:A:68:GLY:HA3	1.94	0.50
1:B:65:LEU:HD23	1:B:68:GLY:HA3	1.95	0.49
1:B:43:LEU:HB2	1:B:56:ILE:HD11	1.95	0.48
1:A:43:LEU:HB2	1:A:56:ILE:HD11	1.95	0.47
1:A:325:LYS:HA	3:A:608:PEG:H41	1.95	0.47
1:A:335:GLN:HG2	1:A:368:VAL:HG11	1.97	0.47
1:B:80:TYR:CE1	1:B:85:HIS:HB2	2.50	0.47
1:A:70:LYS:H	2:A:607:EDO:H12	1.81	0.46
1:A:386[A]:ARG:HB3	1:A:402:LYS:HB2	1.97	0.45
1:B:56:ILE:HG12	1:B:65:LEU:HD12	2.01	0.42
1:A:4:ARG:HD3	1:A:256:GLN:HA	2.00	0.42
1:B:526:GLN:O	1:B:530:GLU:HG2	2.20	0.42
1:A:56:ILE:HG12	1:A:65:LEU:HD12	2.01	0.42
1:B:138:ARG:HH12	1:B:141:GLN:HE22	1.66	0.41
1:A:290:VAL:HG22	1:A:309:ILE:HG13	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:VAL:HG11	1:B:344:MET:HG3	2.03	0.41
1:A:261:LEU:O	2:A:604:EDO:H21	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/539 (90%)	473 (97%)	12 (2%)	1 (0%)	43	54
1	B	477/539 (88%)	462 (97%)	14 (3%)	1 (0%)	43	54
All	All	963/1078 (89%)	935 (97%)	26 (3%)	2 (0%)	43	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	505	VAL
1	B	505	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/480 (93%)	426 (96%)	19 (4%)	26	38
1	B	440/480 (92%)	424 (96%)	16 (4%)	31	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	885/960 (92%)	850 (96%)	35 (4%)	28	41

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	VAL
1	A	83	GLU
1	A	89	LYS
1	A	91	LYS
1	A	92	ASN
1	A	117	LEU
1	A	141	GLN
1	A	171	MET
1	A	185	GLU
1	A	190	LEU
1	A	211	GLN
1	A	220	ARG
1	A	226	ILE
1	A	249	GLU
1	A	264	SER
1	A	359	GLU
1	A	411	THR
1	A	476	ILE
1	A	508	GLU
1	B	84	HIS
1	B	96	ILE
1	B	116	HIS
1	B	141	GLN
1	B	171	MET
1	B	185	GLU
1	B	190	LEU
1	B	205	THR
1	B	226	ILE
1	B	249	GLU
1	B	264	SER
1	B	335	GLN
1	B	359	GLU
1	B	476	ILE
1	B	508	GLU
1	B	528	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	143	HIS
1	A	200	ASN
1	A	217	ASN
1	A	256	GLN
1	A	257	GLN
1	A	281	ASN
1	A	339	ASN
1	A	446	GLN
1	B	37	ASN
1	B	48	ASN
1	B	57	GLN
1	B	87	GLN
1	B	141	GLN
1	B	200	ASN
1	B	217	ASN
1	B	257	GLN
1	B	271	GLN
1	B	281	ASN
1	B	339	ASN
1	B	520	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	607	-	3,3,3	0.48	0	2,2,2	0.34	0
2	EDO	A	606	-	3,3,3	0.50	0	2,2,2	0.39	0
2	EDO	A	602	-	3,3,3	0.46	0	2,2,2	0.41	0
2	EDO	B	604	-	3,3,3	0.51	0	2,2,2	0.34	0
2	EDO	B	602	-	3,3,3	0.53	0	2,2,2	0.28	0
2	EDO	A	603	-	3,3,3	0.50	0	2,2,2	0.39	0
2	EDO	A	605	-	3,3,3	0.50	0	2,2,2	0.34	0
3	PEG	A	608	-	6,6,6	0.07	0	5,5,5	0.03	0
2	EDO	B	603	-	3,3,3	0.45	0	2,2,2	0.44	0
2	EDO	B	605	-	3,3,3	0.48	0	2,2,2	0.35	0
4	SO4	B	601	-	4,4,4	0.22	0	6,6,6	0.10	0
2	EDO	A	604	-	3,3,3	0.41	0	2,2,2	0.55	0
2	EDO	A	601	-	3,3,3	0.48	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	607	-	-	0/1/1/1	-
2	EDO	A	606	-	-	0/1/1/1	-
2	EDO	A	602	-	-	0/1/1/1	-
2	EDO	B	604	-	-	0/1/1/1	-
2	EDO	B	602	-	-	0/1/1/1	-
2	EDO	A	603	-	-	0/1/1/1	-
2	EDO	A	605	-	-	0/1/1/1	-
3	PEG	A	608	-	-	0/4/4/4	-
2	EDO	B	603	-	-	1/1/1/1	-
2	EDO	B	605	-	-	0/1/1/1	-
2	EDO	A	604	-	-	0/1/1/1	-
2	EDO	A	601	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	603	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	607	EDO	3	0
3	A	608	PEG	1	0
2	B	605	EDO	2	0
2	A	604	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	498/539 (92%)	1.19	101 (20%) 3 2	21, 47, 78, 115	2 (0%)
1	B	492/539 (91%)	1.19	83 (16%) 4 3	20, 47, 78, 109	1 (0%)
All	All	990/1078 (91%)	1.19	184 (18%) 3 2	20, 47, 78, 115	3 (0%)

All (184) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	533	GLN	6.0
1	A	119	GLY	5.9
1	A	122	ALA	5.4
1	B	88	LEU	5.2
1	A	409	GLY	4.6
1	B	85	HIS	4.5
1	A	534	LYS	4.4
1	A	109	SER	4.3
1	A	85	HIS	4.0
1	B	86	GLY	3.9
1	A	11	ILE	3.7
1	B	66	TYR	3.7
1	A	165	SER	3.6
1	A	125	LEU	3.6
1	A	169	HIS	3.6
1	B	111	ARG	3.6
1	B	84	HIS	3.5
1	B	298	ASN	3.5
1	A	171	MET	3.5
1	B	167	VAL	3.5
1	B	87	GLN	3.5
1	A	437	ASP	3.4
1	A	248	TRP	3.4
1	B	95	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	476	ILE	3.4
1	A	324	LYS	3.3
1	A	522	ILE	3.3
1	B	224	ALA	3.3
1	B	96	ILE	3.3
1	A	210	LEU	3.2
1	A	526	GLN	3.2
1	A	234	SER	3.2
1	A	252	GLU	3.2
1	B	68	GLY	3.2
1	B	469	PHE	3.2
1	B	409	GLY	3.2
1	A	86	GLY	3.1
1	A	221	ILE	3.1
1	B	35	LYS	3.1
1	A	311	MET	3.1
1	B	485	ASP	3.1
1	B	48	ASN	3.1
1	B	301	VAL	3.1
1	A	34	SER	3.0
1	A	91	LYS	3.0
1	A	226	ILE	3.0
1	A	488	ILE	3.0
1	B	275	ASN	3.0
1	A	224	ALA	3.0
1	B	119	GLY	3.0
1	B	2	THR	3.0
1	A	256	GLN	3.0
1	A	167	VAL	2.9
1	A	223	ALA	2.9
1	A	89	LYS	2.9
1	B	116	HIS	2.9
1	B	166	LYS	2.8
1	B	194	VAL	2.8
1	B	340	ASP	2.8
1	A	7	PHE	2.8
1	B	529	ILE	2.8
1	A	524	THR	2.8
1	A	216	LEU	2.7
1	A	108	THR	2.7
1	A	485	ASP	2.7
1	B	295	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	116	HIS	2.7
1	A	246	GLY	2.7
1	B	463	ILE	2.7
1	A	471	VAL	2.7
1	A	472	ILE	2.6
1	A	10	ASN	2.6
1	A	313	GLU	2.6
1	B	234	SER	2.6
1	A	168	THR	2.6
1	A	525	LEU	2.6
1	B	203	VAL	2.6
1	B	114	HIS	2.6
1	B	40	ASP	2.6
1	A	126	LEU	2.6
1	B	204	GLU	2.6
1	B	335	GLN	2.6
1	A	150	SER	2.6
1	A	124	LYS	2.6
1	A	521	TYR	2.6
1	B	360	VAL	2.5
1	A	469	PHE	2.5
1	B	117	LEU	2.5
1	A	123	GLU	2.5
1	B	484	VAL	2.5
1	B	4	ARG	2.5
1	B	215	PRO	2.5
1	B	62	TYR	2.5
1	A	479	ILE	2.5
1	B	216	LEU	2.5
1	A	484	VAL	2.5
1	B	165	SER	2.5
1	B	205	THR	2.4
1	A	67	GLY	2.4
1	A	451	ASP	2.4
1	A	117	LEU	2.4
1	A	115	GLY	2.4
1	B	332	GLY	2.4
1	B	206	LEU	2.4
1	B	211	GLN	2.4
1	A	4	ARG	2.4
1	A	199	LYS	2.4
1	A	96	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	437	ASP	2.4
1	B	171	MET	2.4
1	A	37	ASN	2.4
1	A	410	ASN	2.4
1	A	293	HIS	2.3
1	B	113	PHE	2.3
1	A	230	VAL	2.3
1	A	298	ASN	2.3
1	B	50	ALA	2.3
1	B	218	THR	2.3
1	B	3	SER	2.3
1	B	472	ILE	2.3
1	A	229	ARG	2.3
1	A	520	HIS	2.3
1	B	54	ILE	2.3
1	A	170	VAL	2.3
1	A	18	ASN	2.2
1	B	192	ASP	2.2
1	A	481	GLU	2.2
1	A	147	PHE	2.2
1	A	247	PHE	2.2
1	A	9	PRO	2.2
1	B	270	ARG	2.2
1	B	52	THR	2.2
1	A	39	GLY	2.2
1	A	206	LEU	2.2
1	B	155	ASP	2.2
1	B	121	GLU	2.2
1	A	275	ASN	2.2
1	A	87	GLN	2.2
1	B	208	THR	2.2
1	B	271	GLN	2.2
1	A	483	GLY	2.2
1	B	532	GLU	2.2
1	B	402	LYS	2.2
1	A	450	MET	2.2
1	B	450	MET	2.2
1	A	204	GLU	2.2
1	A	480	ARG	2.2
1	A	300	PRO	2.2
1	A	208	THR	2.1
1	A	476	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	432	PRO	2.1
1	B	190	LEU	2.1
1	A	325	LYS	2.1
1	B	213	LYS	2.1
1	A	148	VAL	2.1
1	B	5	ARG	2.1
1	A	90	GLU	2.1
1	B	449	ILE	2.1
1	B	366	LYS	2.1
1	A	84	HIS	2.1
1	A	257	GLN	2.1
1	B	187	PHE	2.1
1	A	92	ASN	2.1
1	B	65	LEU	2.1
1	A	295	GLY	2.1
1	A	463	ILE	2.1
1	A	477	ASP	2.1
1	A	326	SER	2.1
1	A	312	PRO	2.1
1	B	143	HIS	2.1
1	A	222	ASN	2.0
1	B	43	LEU	2.0
1	B	49	GLY	2.0
1	B	523	GLU	2.0
1	B	266	LYS	2.0
1	A	294	ASP	2.0
1	B	294	ASP	2.0
1	B	341	PHE	2.0
1	A	88	LEU	2.0
1	A	102	LEU	2.0
1	B	153	THR	2.0
1	A	38	PRO	2.0
1	A	3	SER	2.0
1	B	293	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	B	601	5/5	0.61	0.27	107,107,107,107	0
2	EDO	A	604	4/4	0.69	0.30	52,53,53,53	0
2	EDO	B	603	4/4	0.78	0.21	36,36,37,38	0
2	EDO	A	606	4/4	0.78	0.19	65,65,65,65	0
2	EDO	B	604	4/4	0.79	0.14	48,48,49,49	0
3	PEG	A	608	7/7	0.79	0.19	69,69,69,69	0
2	EDO	A	603	4/4	0.79	0.13	47,48,48,49	0
2	EDO	A	601	4/4	0.80	0.15	46,46,47,47	0
2	EDO	B	602	4/4	0.81	0.17	52,52,52,52	0
2	EDO	A	605	4/4	0.82	0.16	51,51,51,51	0
2	EDO	A	607	4/4	0.86	0.15	42,43,43,43	0
2	EDO	A	602	4/4	0.88	0.19	44,44,44,44	0
2	EDO	B	605	4/4	0.88	0.13	41,42,42,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.