



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 07:25 PM UTC

PDB ID : 7NWQ / pdb_00007nwq
Title : A carbohydrate binding module family 9 (CBM9) from Caldicellulosiruptor kristjanssonii in complex with cellotriose
Authors : Krska, D.; Mazurkewich, S.; Navarro Poulsen, J.; Larsbrink, J.; Lo Leggio, L.
Deposited on : 2021-03-17
Resolution : 2.23 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	215	5% 75% 15% 9%
2	BBB	3	67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BGC	BBB	2	X	-	-	-
3	PEG	AAA	303	-	-	X	-
3	PEG	AAA	305	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 1877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-xylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	195	Total	C	N	O	S	0	4	0
			1608	1017	272	316	3			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	1	MET	-	initiating methionine	UNP E4S6E9
AAA	2	GLY	-	expression tag	UNP E4S6E9
AAA	3	SER	-	expression tag	UNP E4S6E9
AAA	4	SER	-	expression tag	UNP E4S6E9
AAA	5	HIS	-	expression tag	UNP E4S6E9
AAA	6	HIS	-	expression tag	UNP E4S6E9
AAA	7	HIS	-	expression tag	UNP E4S6E9
AAA	8	HIS	-	expression tag	UNP E4S6E9
AAA	9	HIS	-	expression tag	UNP E4S6E9
AAA	10	HIS	-	expression tag	UNP E4S6E9
AAA	11	SER	-	expression tag	UNP E4S6E9
AAA	12	SER	-	expression tag	UNP E4S6E9
AAA	13	GLU	-	expression tag	UNP E4S6E9
AAA	14	ASN	-	expression tag	UNP E4S6E9
AAA	15	LEU	-	expression tag	UNP E4S6E9
AAA	16	TYR	-	expression tag	UNP E4S6E9
AAA	17	PHE	-	expression tag	UNP E4S6E9
AAA	18	GLN	-	expression tag	UNP E4S6E9
AAA	19	GLY	-	expression tag	UNP E4S6E9
AAA	20	HIS	-	expression tag	UNP E4S6E9

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	BBB	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			7	4	3		
3	AAA	1	Total	C	O	0	0
			7	4	3		
3	AAA	1	Total	C	O	0	0
			7	4	3		
3	AAA	1	Total	C	O	0	0
			7	4	3		
3	AAA	1	Total	C	O	0	0
			7	4	3		
3	AAA	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	AAA	1	Total	C	O	0	0
			4	2	2		
4	AAA	1	Total	C	O	0	0
			4	2	2		
4	AAA	1	Total	C	O	0	0
			4	2	2		
4	AAA	1	Total	C	O	0	0
			4	2	2		
4	AAA	1	Total	C	O	0	0
			4	2	2		
4	AAA	1	Total	C	O	0	0
			4	2	2		
4	AAA	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	3	Total	Ca	0	0
			3	3		

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	C	O	0	0
			10	6	4		
6	AAA	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	AAA	1	Total	C	O	0	0
			16	10	6		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	AAA	111	Total 111	O 111	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-xylanase



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	172.49Å 172.49Å 172.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.52 – 2.23 70.52 – 2.23	Depositor EDS
% Data completeness (in resolution range)	99.7 (70.52-2.23) 99.7 (70.52-2.23)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.94 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.185 , 0.226 0.185 , 0.226	Depositor DCC
R_{free} test set	2000 reflections (9.25%)	wwPDB-VP
Wilson B-factor (Å ²)	41.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1877	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CA, BGC, PEG, 1PE, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.68	0/1645	1.10	2/2237 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	99	ASP	CA-CB-CG	5.55	118.15	112.60
1	AAA	133	PHE	CA-CB-CG	5.46	119.26	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1608	0	1540	35	0
2	BBB	34	0	30	0	1
3	AAA	49	0	70	21	0
4	AAA	36	0	54	5	0
5	AAA	3	0	0	0	0
6	AAA	20	0	28	1	0
7	AAA	16	0	22	2	0
8	AAA	111	0	0	9	1
All	All	1877	0	1744	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:38:ASP:H	3:AAA:305:PEG:H42	1.03	1.15
1:AAA:38:ASP:N	3:AAA:305:PEG:H42	1.74	1.01
1:AAA:38:ASP:H	3:AAA:305:PEG:C4	1.82	0.92
7:AAA:321:1PE:H142	8:AAA:423:HOH:O	1.73	0.88
4:AAA:310:EDO:H21	8:AAA:503:HOH:O	1.78	0.83

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:BBB:3:BGC:O2	8:AAA:511:HOH:O[18_655]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	197/215 (92%)	189 (96%)	7 (4%)	1 (0%)	24 23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	214	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	176/190 (93%)	168 (96%)	8 (4%)	24	26

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	208	ASN
1	AAA	206	SER
1	AAA	173	SER
1	AAA	56[B]	GLN
1	AAA	176	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	BBB	1	2	12,12,12	0.60	0	17,17,17	1.78	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	BBB	2	2	11,11,12	1.03	1 (9%)	15,15,17	2.43	6 (40%)
2	BGC	BBB	3	2	11,11,12	0.50	0	15,15,17	1.78	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	BBB	1	2	-	2/2/22/22	0/1/1/1
2	BGC	BBB	2	2	1/1/4/5	2/2/19/22	0/1/1/1
2	BGC	BBB	3	2	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	BBB	2	BGC	O4-C4	-2.82	1.36	1.43

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	BBB	2	BGC	O4-C4-C5	4.95	121.52	109.32
2	BBB	3	BGC	C2-C3-C4	-3.86	104.06	110.86
2	BBB	1	BGC	O5-C5-C4	-3.71	103.01	109.70
2	BBB	2	BGC	O4-C4-C3	3.57	118.80	110.38
2	BBB	2	BGC	C1-O5-C5	3.30	116.61	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	BBB	2	BGC	C4

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	BBB	1	BGC	C4-C5-C6-O6
2	BBB	3	BGC	O5-C5-C6-O6
2	BBB	1	BGC	O5-C5-C6-O6
2	BBB	2	BGC	C4-C5-C6-O6
2	BBB	3	BGC	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	3	BGC	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 3 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	AAA	315	-	3,3,3	0.21	0	2,2,2	0.43	0
3	PEG	AAA	302	-	6,6,6	0.26	0	5,5,5	0.15	0
4	EDO	AAA	312	-	3,3,3	0.26	0	2,2,2	0.51	0
4	EDO	AAA	314	-	3,3,3	0.17	0	2,2,2	0.38	0
3	PEG	AAA	304	-	6,6,6	0.26	0	5,5,5	0.27	0
3	PEG	AAA	306	-	6,6,6	0.12	0	5,5,5	0.22	0
4	EDO	AAA	316	-	3,3,3	0.19	0	2,2,2	0.26	0
6	PGE	AAA	320	-	9,9,9	0.42	0	8,8,8	0.30	0
4	EDO	AAA	311	-	3,3,3	0.14	0	2,2,2	0.43	0
7	1PE	AAA	321	-	15,15,15	0.31	0	14,14,14	0.24	0
3	PEG	AAA	307	-	6,6,6	0.71	0	5,5,5	0.35	0
6	PGE	AAA	322	-	9,9,9	0.28	0	8,8,8	0.22	0
3	PEG	AAA	303	-	6,6,6	0.28	0	5,5,5	0.23	0
3	PEG	AAA	305	-	6,6,6	0.51	0	5,5,5	0.98	0
3	PEG	AAA	301	-	6,6,6	0.15	0	5,5,5	0.09	0
4	EDO	AAA	309	-	3,3,3	0.11	0	2,2,2	0.38	0
4	EDO	AAA	313	-	3,3,3	0.24	0	2,2,2	0.78	0
4	EDO	AAA	308	-	3,3,3	0.46	0	2,2,2	0.87	0
4	EDO	AAA	310	-	3,3,3	0.06	0	2,2,2	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	AAA	315	-	-	1/1/1/1	-
3	PEG	AAA	302	-	-	3/4/4/4	-
4	EDO	AAA	312	-	-	1/1/1/1	-
4	EDO	AAA	314	-	-	0/1/1/1	-
3	PEG	AAA	304	-	-	3/4/4/4	-
3	PEG	AAA	306	-	-	1/4/4/4	-
4	EDO	AAA	316	-	-	1/1/1/1	-
6	PGE	AAA	320	-	-	2/7/7/7	-
4	EDO	AAA	311	-	-	1/1/1/1	-
7	1PE	AAA	321	-	-	6/13/13/13	-
3	PEG	AAA	307	-	-	3/4/4/4	-
6	PGE	AAA	322	-	-	3/7/7/7	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	AAA	303	-	-	3/4/4/4	-
3	PEG	AAA	305	-	-	1/4/4/4	-
3	PEG	AAA	301	-	-	1/4/4/4	-
4	EDO	AAA	309	-	-	1/1/1/1	-
4	EDO	AAA	313	-	-	1/1/1/1	-
4	EDO	AAA	308	-	-	0/1/1/1	-
4	EDO	AAA	310	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	AAA	320	PGE	C3-C4-O3-C5
3	AAA	307	PEG	C1-C2-O2-C3
6	AAA	322	PGE	O3-C5-C6-O4
3	AAA	305	PEG	O2-C3-C4-O4
7	AAA	321	1PE	OH5-C14-C24-OH4

There are no ring outliers.

12 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	302	PEG	1	0
4	AAA	314	EDO	1	0
3	AAA	304	PEG	2	0
4	AAA	316	EDO	1	0
6	AAA	320	PGE	1	0
4	AAA	311	EDO	1	0
7	AAA	321	1PE	2	0
3	AAA	307	PEG	3	0
3	AAA	303	PEG	4	0
3	AAA	305	PEG	11	0
4	AAA	308	EDO	1	0
4	AAA	310	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	195/215 (90%)	-0.10	10 (5%) 33 32	13, 34, 71, 167	4 (2%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	213	ILE	10.1
1	AAA	209	ALA	10.1
1	AAA	210	VAL	6.3
1	AAA	211	VAL	5.9
1	AAA	212	PRO	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	BBB	1	12/12	-	-	23,30,36,40	12
2	BGC	BBB	2	11/12	-	-	37,41,44,45	11
2	BGC	BBB	3	11/12	-	-	31,35,38,42	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	PGE	AAA	320	10/10	0.73	0.36	72,96,122,140	0
4	EDO	AAA	313	4/4	0.75	0.27	73,76,82,88	0
4	EDO	AAA	312	4/4	0.75	0.34	73,79,82,94	0
3	PEG	AAA	307	7/7	0.81	0.22	20,22,26,30	7
4	EDO	AAA	316	4/4	0.83	0.26	85,85,98,98	0
6	PGE	AAA	322	10/10	0.83	0.24	61,80,105,108	0
3	PEG	AAA	306	7/7	0.85	0.22	74,76,85,88	0
4	EDO	AAA	308	4/4	0.85	0.23	65,67,70,80	0
4	EDO	AAA	315	4/4	0.85	0.21	61,82,86,89	0
7	1PE	AAA	321	16/16	0.85	0.31	56,108,144,158	0
4	EDO	AAA	311	4/4	0.86	0.17	55,57,64,69	0
3	PEG	AAA	301	7/7	0.86	0.25	74,84,98,101	0
4	EDO	AAA	309	4/4	0.86	0.19	70,79,84,86	0
4	EDO	AAA	310	4/4	0.86	0.28	64,65,71,76	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	AAA	303	7/7	0.91	0.14	38,46,52,55	0
3	PEG	AAA	305	7/7	0.92	0.17	13,16,20,22	7
3	PEG	AAA	302	7/7	0.93	0.13	56,63,83,85	0
3	PEG	AAA	304	7/7	0.93	0.14	69,72,79,87	0
4	EDO	AAA	314	4/4	0.94	0.18	51,55,57,58	0
5	CA	AAA	319	1/1	0.99	0.05	44,44,44,44	1
5	CA	AAA	317	1/1	0.99	0.10	49,49,49,49	0
5	CA	AAA	318	1/1	1.00	0.02	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.