



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 02:34 AM UTC

PDB ID : 4NXU / pdb_00004nxu
Title : Crystal structure of the cytosolic domain of human MiD51
Authors : Richter, V.; Kvansakul, M.; Ryan, M.T.
Deposited on : 2013-12-09
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	 4% 82% 12% • 6%
1	B	347	 3% 84% 10% • 5%
1	C	347	 3% 86% 8% • •
1	D	347	 14% 78% 15% • 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	C	504	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20858 atoms, of which 10206 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitochondrial dynamic protein MID51.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	327	Total	C	H	N	O	S	0	1	0
			5085	1634	2543	425	471	12			
1	B	328	Total	C	H	N	O	S	0	0	0
			5146	1646	2576	435	478	11			
1	C	332	Total	C	H	N	O	S	0	0	0
			5145	1652	2572	436	474	11			
1	D	326	Total	C	H	N	O	S	0	0	0
			4937	1606	2451	412	457	11			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	117	GLY	-	expression tag	UNP Q9NQG6
A	118	SER	-	expression tag	UNP Q9NQG6
B	117	GLY	-	expression tag	UNP Q9NQG6
B	118	SER	-	expression tag	UNP Q9NQG6
C	117	GLY	-	expression tag	UNP Q9NQG6
C	118	SER	-	expression tag	UNP Q9NQG6
D	117	GLY	-	expression tag	UNP Q9NQG6
D	118	SER	-	expression tag	UNP Q9NQG6

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	B	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	C	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	D	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0


- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C H O 14 3 8 3	0	0
4	C	1	Total C H O 14 3 8 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	113	Total O 113 113	0	0
5	B	99	Total O 99 99	0	0
5	C	85	Total O 85 85	0	0
5	D	44	Total O 44 44	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	72.57Å 79.27Å 79.38Å 65.44° 84.18° 63.35°	Depositor
Resolution (Å)	39.19 – 2.30 39.19 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.8 (39.19-2.30) 93.4 (39.19-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.216 , 0.248 0.220 , 0.251	Depositor DCC
R_{free} test set	3091 reflections (3.17%)	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	20858	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	1/2600 (0.0%)	0.74	4/3547 (0.1%)
1	B	0.41	1/2625 (0.0%)	0.75	0/3578
1	C	0.38	0/2632	0.72	0/3595
1	D	0.41	0/2542	0.77	1/3477 (0.0%)
All	All	0.41	2/10399 (0.0%)	0.75	5/14197 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	237	ASN	CA-C	7.28	1.62	1.52
1	B	276	TRP	NE1-CE2	-5.10	1.31	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	ASN	O-C-N	7.91	130.41	121.32
1	A	237	ASN	CA-C-N	-6.28	113.22	119.56
1	A	237	ASN	C-N-CA	-6.28	113.22	119.56
1	A	459	VAL	N-CA-C	-5.69	105.55	111.58
1	D	201	HIS	N-CA-C	5.32	116.93	108.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2542	2543	2543	20	2
1	B	2570	2576	2577	24	0
1	C	2573	2572	2569	19	1
1	D	2486	2451	2451	30	1
2	A	27	12	12	0	0
2	B	27	12	12	0	0
2	C	27	12	12	0	0
2	D	27	12	12	0	0
3	A	5	0	0	0	0
3	C	15	0	0	0	2
4	B	6	8	8	3	0
4	C	6	8	8	0	0
5	A	113	0	0	0	0
5	B	99	0	0	0	0
5	C	85	0	0	0	0
5	D	44	0	0	2	0
All	All	10652	10206	10204	86	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:376:ALA:HB1	1:D:449:THR:HG21	1.66	0.78
1:B:267:GLU:HG3	1:B:295:ALA:HB1	1.67	0.75
1:A:299:GLU:OE2	1:B:288:ARG:NH1	2.21	0.74
1:A:458:GLU:HG3	1:D:331:ALA:HB3	1.72	0.70
1:C:448:TYR:HA	1:C:451:TYR:HB2	1.74	0.69

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ASN:HD21	3:C:504:SO4:O2[1_645]	1.55	0.05
1:A:213:ASN:ND2	3:C:504:SO4:O2[1_645]	2.15	0.05
1:C:144:ARG:O	1:D:212:GLN:NE2[1_556]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/347 (93%)	315 (97%)	9 (3%)	0	100	100
1	B	324/347 (93%)	316 (98%)	8 (2%)	0	100	100
1	C	330/347 (95%)	324 (98%)	6 (2%)	0	100	100
1	D	320/347 (92%)	304 (95%)	16 (5%)	0	100	100
All	All	1298/1388 (94%)	1259 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/298 (92%)	257 (94%)	16 (6%)	18	26
1	B	278/298 (93%)	270 (97%)	8 (3%)	37	55
1	C	275/298 (92%)	266 (97%)	9 (3%)	33	50
1	D	260/298 (87%)	242 (93%)	18 (7%)	14	20
All	All	1086/1192 (91%)	1035 (95%)	51 (5%)	23	35

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	431	LYS
1	D	158	LYS
1	D	458	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	448	TYR
1	C	456	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	226	ASN
1	C	354	GLN
1	B	410	GLN
1	B	275	ASN
1	B	462	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	C	502	-	5,5,5	0.91	0	5,5,5	0.91	0
3	SO4	A	502	-	4,4,4	0.34	0	6,6,6	0.16	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	C	504	-	4,4,4	0.30	0	6,6,6	0.23	0
4	GOL	B	502	-	5,5,5	0.95	0	5,5,5	0.93	0
2	ADP	D	501	-	28,29,29	1.48	5 (17%)	43,45,45	1.70	11 (25%)
2	ADP	C	501	-	28,29,29	1.48	4 (14%)	43,45,45	1.73	9 (20%)
2	ADP	A	501	-	28,29,29	1.54	5 (17%)	43,45,45	1.74	9 (20%)
2	ADP	B	501	-	28,29,29	1.53	5 (17%)	43,45,45	1.73	9 (20%)
3	SO4	C	503	-	4,4,4	0.30	0	6,6,6	0.13	0
3	SO4	C	505	-	4,4,4	0.34	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	502	-	-	2/4/4/4	-
4	GOL	B	502	-	-	4/4/4/4	-
2	ADP	D	501	-	-	2/16/32/32	0/3/3/3
2	ADP	C	501	-	-	3/16/32/32	0/3/3/3
2	ADP	A	501	-	-	2/16/32/32	0/3/3/3
2	ADP	B	501	-	-	2/16/32/32	0/3/3/3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	ADP	C5-C4	5.08	1.48	1.39
2	A	501	ADP	C5-C4	5.04	1.48	1.39
2	C	501	ADP	C5-C4	4.96	1.47	1.39
2	D	501	ADP	C5-C4	4.96	1.47	1.39
2	A	501	ADP	C5-N7	-2.79	1.34	1.39

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ADP	C5-C4-N3	-5.42	119.25	126.72
2	C	501	ADP	C5-C4-N3	-5.06	119.75	126.72
2	B	501	ADP	C5-C4-N3	-4.89	119.98	126.72
2	D	501	ADP	C5-C4-N3	-4.30	120.79	126.72
2	D	501	ADP	C4-C5-N7	-4.18	105.81	110.58

There are no chirality outliers.

5 of 15 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	B	502	GOL	O1-C1-C2-C3
4	C	502	GOL	C1-C2-C3-O3
4	B	502	GOL	C1-C2-C3-O3
2	C	501	ADP	C3'-C4'-C5'-O5'
2	C	501	ADP	O4'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	504	SO4	0	2
4	B	502	GOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/347 (94%)	0.23	15 (4%) 37 39	21, 41, 69, 103	1 (0%)
1	B	328/347 (94%)	0.21	10 (3%) 52 54	25, 40, 65, 102	0
1	C	332/347 (95%)	0.38	12 (3%) 46 48	23, 48, 81, 113	0
1	D	326/347 (93%)	1.02	47 (14%) 6 7	39, 64, 93, 142	0
All	All	1313/1388 (94%)	0.46	84 (6%) 25 27	21, 48, 83, 142	1 (0%)

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	462	GLN	6.9
1	D	294	GLU	5.3
1	C	451	TYR	5.1
1	A	462	GLN	4.9
1	D	353	ASP	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	502	6/6	0.70	0.24	35,42,46,46	0
4	GOL	C	502	6/6	0.74	0.23	32,39,41,42	0
3	SO4	C	505	5/5	0.81	0.30	44,45,46,47	0
3	SO4	A	502	5/5	0.89	0.23	46,47,48,49	0
3	SO4	C	503	5/5	0.90	0.19	44,45,46,46	0
3	SO4	C	504	5/5	0.94	0.15	42,43,44,45	0
2	ADP	D	501	27/27	0.94	0.09	42,44,53,71	0
2	ADP	B	501	27/27	0.95	0.08	23,34,42,43	0
2	ADP	C	501	27/27	0.96	0.07	36,40,50,88	0
2	ADP	A	501	27/27	0.96	0.07	24,31,39,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around ADP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.