



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 02:24 PM UTC

PDB ID : 7NYW / pdb_00007nyw
EMDB ID : EMD-12656
Title : Cryo-EM structure of the MukBEF-MatP-DNA head module
Authors : Buermann, F.; Lowe, J.
Deposited on : 2021-03-23
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Q-score	-	25397	14724 (2.60 - 3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1482	
1	B	1482	
2	C	440	
2	D	440	
3	E	240	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	240	
4	G	78	
4	H	78	
5	I	151	
5	J	151	
6	K	80	
7	L	80	
8	M	30	
8	N	30	

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 48634 atoms, of which 23846 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chromosome partition protein MukB.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	685	Total	C	H	N	O	S	0	0
			11041	3411	5563	1025	1020	22		
1	B	858	Total	C	H	N	O	S	0	0
			13847	4273	6953	1289	1306	26		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1407	GLN	GLU	engineered mutation	UNP A0A0F7LRY2
B	1407	GLN	GLU	engineered mutation	UNP A0A0F7LRY2

- Molecule 2 is a protein called Chromosome partition protein MukF.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	C	335	Total	C	H	N	O	S	0	0
			5330	1702	2626	470	524	8		
2	D	94	Total	C	H	N	O	S	0	0
			1469	461	736	125	143	4		

- Molecule 3 is a protein called Chromosome partition protein MukE.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	E	212	Total	C	H	N	O	S	0	0
			3441	1090	1719	301	322	9		
3	F	198	Total	C	H	N	O	S	0	0
			3246	1029	1627	284	298	8		

- Molecule 4 is a protein called Acyl carrier protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	G	72	Total	C	H	N	O	S	0	0
			1105	349	543	85	127	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						AltConf	Trace
4	H	72	Total	C	H	N	O	S	0	0
			1105	349	543	85	127	1		

- Molecule 5 is a protein called Macrodmain Ter protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	I	134	Total	C	H	N	O	S	0	0
			2272	714	1137	215	202	4		
5	J	135	Total	C	H	N	O	S	0	0
			2288	719	1144	217	204	4		

- Molecule 6 is a DNA chain called matS2 DNA 80 b, oligo FBA769.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	K	22	Total	C	H	N	O	P	0	0
			694	214	249	80	130	21		

- Molecule 7 is a DNA chain called matS2 DNA 80 b, oligo FBA770.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	L	22	Total	C	H	N	O	P	0	0
			702	216	248	84	132	22		

- Molecule 8 is a DNA chain called DNA 80 b.

Mol	Chain	Residues	Atoms						AltConf	Trace
8	M	30	Total	C	H	N	O	P	0	0
			961	300	346	105	180	30		
8	N	30	Total	C	H	N	O	P	0	0
			961	300	346	105	180	30		

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
9	A	1	Total	Mg	0
			1	1	
9	B	1	Total	Mg	0
			1	1	

- Molecule 10 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms						AltConf
10	A	1	Total	C	H	N	O	P	0
			43	10	12	5	13	3	
10	B	1	Total	C	H	N	O	P	0
			43	10	12	5	13	3	

- Molecule 11 is 4'-PHOSPHOPANTETHEINE (CCD ID: PNS) (formula: C₁₁H₂₃N₂O₇PS).





Mol	Chain	Residues	Atoms							AltConf
11	G	1	Total	C	H	N	O	P	S	0
			42	11	21	2	6	1	1	
11	H	1	Total	C	H	N	O	P	S	0
			42	11	21	2	6	1	1	

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Chromosome partition protein MukB





● Molecule 1: Chromosome partition protein MukB



LEU
GLY
VAL
ALA
GLU
GLU
ASP
PHE
SER
GLY
LEU
PRO
ALA
GLU
TRP
LEU
ALA
TLE
ASN
ASP
TYR
GLY
ALA
ALA
LYS
VAL
GLN
ALA
HIS
VAL
TLE
ASP
THR
TYR

• Molecule 3: Chromosome partition protein MukE

Chain E:



MET
S2
E7
D39
E52
Q53
I66
E70
R76
L82
L91
D92
M93
M94
L99
L102
Q112
E123
V137
R140
S141
S142
G143
S144
D147
K150
K154
T158
R161
L171
K177
T180
T181
V184
V191

D207
A210
M211
P212
V213
GLY
SER
LEU
SER
LEU
ASP
ASP
SER
SER
ASN
GLU
THR
PRO
ASN
ASN
SER
ALA
GLU
GLY
ALA
GLY
ASP
GLN
PRO

• Molecule 3: Chromosome partition protein MukE

Chain F:



MET
SER
SER
THR
HIS
TLE
GLU
GLN
F9
E25
D42
N43
E52
Q53
V63
R78
I83
L104
S105
P106
E107
R108
L109
G113
Y121
I125
D129
K132
L133
M134
V137
N138
Q139
R140
S141
S142
G143
S144
D145
L146
L151
V155
N174
N175

I180
T181
A182
A183
V184
F187
G188
A189
D190
V191
R192
S193
G194
D195
D196
P197
R198
E199
R203
M204
I205
R206
ASP
GLY
GLU
ALA
MET
PRO
VAL
GLU
GLY
SER
SER
SER
LEU
ASP
SER
GLU
ASN
ASP
GLU
THR
PRO
ASP
ASN
SER
ALA
GLU
ALA
GLY
ASP
GLU
GLN
PRO

• Molecule 4: Acyl carrier protein

Chain G:



MET
SER
T2
I3
V7
I11
G12
E20
T23
N24
N25
F28
M44
A45
L46
E47
E57
I62
T63
T64
V65
Q66
A67
I72
N73
GLY
HIS
GLN
ALA

• Molecule 4: Acyl carrier protein

Chain H:



MET
SER
T2
L15
G16
E20
N24
V29
E30
A34
S35
S36
V40
V43
T64
V65
Q66
N73
GLY
HIS
GLN
ALA

• Molecule 5: Macrodomain Ter protein


Chain I:



V1
E10
T29
I32
A40
L47
P51
V52
R53
V54
I58
M62
L66
M70
K71
Q72
R78
N83
A84
T89
R90
V100
V120
K129
Y132
A133
M134
GLN
MET
SER
SER
LEU
LYS
HIS
ASP
LEU
GLU
ALA
TLE
ILE
LEU
GLY
LYS
ASN

GLU

• Molecule 5: Macrodomain Ter protein



- [illegible]



- | Category | Item | Value |
|----------|--------|-------|
| T1 | T1.1 | 0.00 |
| | T1.2 | 0.00 |
| | T1.3 | 0.00 |
| | T1.4 | 0.00 |
| G5 | G5.1 | 0.00 |
| | G5.2 | 0.00 |
| | G5.3 | 0.00 |
| | G5.4 | 0.00 |
| T1.4 | T1.4.1 | 0.00 |
| | T1.4.2 | 0.00 |
| | T1.4.3 | 0.00 |
| | T1.4.4 | 0.00 |
| G22 | G22.1 | 0.00 |
| | G22.2 | 0.00 |
| | G22.3 | 0.00 |
| | G22.4 | 0.00 |
| D1 | D1.1 | 0.00 |
| | D1.2 | 0.00 |
| | D1.3 | 0.00 |
| | D1.4 | 0.00 |
| D2 | D2.1 | 0.00 |
| | D2.2 | 0.00 |
| | D2.3 | 0.00 |
| | D2.4 | 0.00 |
| D3 | D3.1 | 0.00 |
| | D3.2 | 0.00 |
| | D3.3 | 0.00 |
| | D3.4 | 0.00 |
| D4 | D4.1 | 0.00 |
| | D4.2 | 0.00 |
| | D4.3 | 0.00 |
| | D4.4 | 0.00 |
| D5 | D5.1 | 0.00 |
| | D5.2 | 0.00 |
| | D5.3 | 0.00 |
| | D5.4 | 0.00 |
| D6 | D6.1 | 0.00 |
| | D6.2 | 0.00 |
| | D6.3 | 0.00 |
| | D6.4 | 0.00 |
| D7 | D7.1 | 0.00 |
| | D7.2 | 0.00 |
| | D7.3 | 0.00 |
| | D7.4 | 0.00 |
| D8 | D8.1 | 0.00 |
| | D8.2 | 0.00 |
| | D8.3 | 0.00 |
| | D8.4 | 0.00 |
| D9 | D9.1 | 0.00 |
| | D9.2 | 0.00 |
| | D9.3 | 0.00 |
| | D9.4 | 0.00 |
| D10 | D10.1 | 0.00 |
| | D10.2 | 0.00 |
| | D10.3 | 0.00 |
| | D10.4 | 0.00 |
| D11 | D11.1 | 0.00 |
| | D11.2 | 0.00 |
| | D11.3 | 0.00 |
| | D11.4 | 0.00 |
| D12 | D12.1 | 0.00 |
| | D12.2 | 0.00 |
| | D12.3 | 0.00 |
| | D12.4 | 0.00 |
| D13 | D13.1 | 0.00 |
| | D13.2 | 0.00 |
| | D13.3 | 0.00 |
| | D13.4 | 0.00 |
| D14 | D14.1 | 0.00 |
| | D14.2 | 0.00 |
| | D14.3 | 0.00 |
| | D14.4 | 0.00 |
| D15 | D15.1 | 0.00 |
| | D15.2 | 0.00 |
| | D15.3 | 0.00 |
| | D15.4 | 0.00 |
| D16 | D16.1 | 0.00 |
| | D16.2 | 0.00 |
| | D16.3 | 0.00 |
| | D16.4 | 0.00 |
| D17 | D17.1 | 0.00 |
| | D17.2 | 0.00 |
| | D17.3 | 0.00 |
| | D17.4 | 0.00 |
| D18 | D18.1 | 0.00 |
| | D18.2 | 0.00 |
| | D18.3 | 0.00 |
| | D18.4 | 0.00 |
| D19 | D19.1 | 0.00 |
| | D19.2 | 0.00 |
| | D19.3 | 0.00 |
| | D19.4 | 0.00 |
| D20 | D20.1 | 0.00 |
| | D20.2 | 0.00 |
| | D20.3 | 0.00 |
| | D20.4 | 0.00 |
| D21 | D21.1 | 0.00 |
| | D21.2 | 0.00 |
| | D21.3 | 0.00 |
| | D21.4 | 0.00 |
| D22 | D22.1 | 0.00 |
| | D22.2 | 0.00 |
| | D22.3 | 0.00 |
| | D22.4 | 0.00 |



-

- 
- A3
T4
A5
T6
A7
T12
A13
T16
A23
T24
T32

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	200438	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.131	Depositor
Minimum map value	-0.037	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.017	Depositor
Map size (Å)	462.23843, 462.23843, 462.23843	wwPDB
Map dimensions	318, 318, 318	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.45358, 1.45358, 1.45358	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PNS, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/5553	0.27	0/7469
1	B	0.13	0/6981	0.26	0/9385
2	C	0.12	0/2753	0.27	0/3727
2	D	0.09	0/741	0.23	0/1000
3	E	0.12	0/1753	0.25	0/2361
3	F	0.12	0/1648	0.28	0/2218
4	G	0.11	0/565	0.24	0/765
4	H	0.11	0/565	0.25	0/765
5	I	0.12	0/1160	0.25	0/1560
5	J	0.13	0/1169	0.26	0/1572
6	K	0.21	0/498	0.42	0/766
7	L	0.21	0/509	0.40	0/784
8	M	0.22	0/689	0.49	0/1061
8	N	0.18	0/689	0.43	0/1061
All	All	0.14	0/25273	0.29	0/34494

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5478	5563	5559	59	0
1	B	6894	6953	6947	90	0
2	C	2704	2626	2623	47	0
2	D	733	736	735	20	0
3	E	1722	1719	1718	30	0
3	F	1619	1627	1626	20	0
4	G	562	543	542	12	0
4	H	562	543	542	6	0
5	I	1135	1137	1136	18	0
5	J	1144	1144	1144	18	0
6	K	445	249	250	2	0
7	L	454	248	249	2	0
8	M	615	346	346	1	0
8	N	615	346	346	5	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	A	31	12	12	3	0
10	B	31	12	12	1	0
11	G	21	21	21	1	0
11	H	21	21	21	0	0
All	All	24788	23846	23829	295	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 295 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:223:PRO:O	1:B:1340:TYR:OH	1.89	0.90
1:B:186:LYS:NZ	1:B:1388:GLU:OE2	2.11	0.83
1:A:1251:GLU:OE1	1:A:1254:ARG:NH1	2.12	0.82
1:B:255:ASP:OD2	1:B:1199:ARG:NH2	2.13	0.81
2:C:262:ARG:NH2	2:D:42:ARG:O	2.15	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	681/1482 (46%)	651 (96%)	30 (4%)	0	100	100
1	B	836/1482 (56%)	806 (96%)	30 (4%)	0	100	100
2	C	331/440 (75%)	316 (96%)	15 (4%)	0	100	100
2	D	92/440 (21%)	89 (97%)	3 (3%)	0	100	100
3	E	210/240 (88%)	206 (98%)	4 (2%)	0	100	100
3	F	196/240 (82%)	186 (95%)	10 (5%)	0	100	100
4	G	70/78 (90%)	69 (99%)	1 (1%)	0	100	100
4	H	70/78 (90%)	67 (96%)	3 (4%)	0	100	100
5	I	132/151 (87%)	129 (98%)	3 (2%)	0	100	100
5	J	133/151 (88%)	131 (98%)	2 (2%)	0	100	100
All	All	2751/4782 (58%)	2650 (96%)	101 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	PNS	H	101	4	14,20,21	0.23	0	18,26,29	0.47	0
10	ATP	B	1502	9	32,33,33	0.51	0	48,52,52	0.72	2 (4%)
11	PNS	G	101	4	14,20,21	0.23	0	18,26,29	0.36	0
10	ATP	A	1502	9	32,33,33	0.55	0	48,52,52	0.72	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	PNS	H	101	4	-	4/24/26/27	-
10	ATP	B	1502	9	-	3/22/38/38	0/3/3/3
11	PNS	G	101	4	-	3/24/26/27	-
10	ATP	A	1502	9	-	2/22/38/38	0/3/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	1502	ATP	O2'-C2'-C3'	-2.10	105.09	111.82
10	A	1502	ATP	O3'-C3'-C2'	-2.05	105.23	111.82
10	B	1502	ATP	O3'-C3'-C2'	-2.05	105.24	111.82
10	B	1502	ATP	O2'-C2'-C3'	-2.00	105.40	111.82

There are no chirality outliers.

5 of 12 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
11	H	101	PNS	C37-C38-C39-N41
11	G	101	PNS	O33-C32-C34-O35
11	H	101	PNS	O33-C32-C34-O35
11	H	101	PNS	C37-C38-C39-O40
10	B	1502	ATP	PB-O3B-PG-O1G


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	B	1502	ATP	1	0
11	G	101	PNS	1	0
10	A	1502	ATP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12656. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 159




Y Index: 159



Z Index: 159


6.2.2 Raw map



X Index: 159



Y Index: 159




Z Index: 159


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 141



Y Index: 145




Z Index: 216


6.3.2 Raw map



X Index: 141



Y Index: 145



Z Index: 214


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.017. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_12656_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 279 nm^3 ; this corresponds to an approximate mass of 252 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.79	6.25	3.88

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.79 differs from the reported value 3.1 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-12656 and PDB model 7NYW. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.017 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.017).

9.4 Atom inclusion ⓘ



At the recommended contour level, 88% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.017) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8790	 0.4520
A	 0.9150	 0.4850
B	 0.8890	 0.4640
C	 0.8360	 0.4130
D	 0.8030	 0.3790
E	 0.8980	 0.4680
F	 0.8360	 0.4260
G	 0.7980	 0.3330
H	 0.8020	 0.3290
I	 0.9170	 0.4930
J	 0.8920	 0.4410
K	 0.9870	 0.4900
L	 0.9800	 0.4890
M	 0.9840	 0.4390
N	 0.9810	 0.4470

