



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:05 AM UTC

PDB ID : 4NZ0 / pdb_00004nz0
Title : The EMCV 3Dpol structure at 2.8Å resolution
Authors : Vives-adrian, L.; Ferrer-orta, C.; Verdaguer, N.
Deposited on : 2013-12-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	
1	B	460	
1	C	460	
1	D	460	
1	E	460	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	460	3% 95% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 22240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	460	Total	C	N	O	S	0	1	0
			3689	2362	628	684	15			
1	B	460	Total	C	N	O	S	0	0	0
			3681	2357	625	684	15			
1	C	460	Total	C	N	O	S	0	0	0
			3681	2357	625	684	15			
1	D	460	Total	C	N	O	S	0	0	0
			3681	2357	625	684	15			
1	E	460	Total	C	N	O	S	0	1	0
			3692	2363	629	685	15			
1	F	460	Total	C	N	O	S	0	0	0
			3681	2357	625	684	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	MET	VAL	SEE REMARK 999	UNP P12296
B	247	MET	VAL	SEE REMARK 999	UNP P12296
C	247	MET	VAL	SEE REMARK 999	UNP P12296
D	247	MET	VAL	SEE REMARK 999	UNP P12296
E	247	MET	VAL	SEE REMARK 999	UNP P12296
F	247	MET	VAL	SEE REMARK 999	UNP P12296

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	29	Total 29	O 29	0	0
4	B	6	Total 6	O 6	0	0
4	C	4	Total 4	O 4	0	0
4	D	8	Total 8	O 8	0	0
4	E	25	Total 25	O 25	0	0
4	F	14	Total 14	O 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Genome polyprotein




- Molecule 1: Genome polyprotein




- Molecule 1: Genome polyprotein




- Molecule 1: Genome polyprotein





- Molecule 1: Genome polyprotein



- Molecule 1: Genome polyprotein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	230.40Å 140.79Å 170.58Å 90.00° 125.86° 90.00°	Depositor
Resolution (Å)	138.20 – 2.80 138.20 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.8 (138.20-2.80) 98.8 (138.20-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.246 , 0.268 0.244 , 0.270	Depositor DCC
R_{free} test set	6006 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	22240	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3780	0.64	0/5121
1	B	0.40	0/3769	0.65	1/5107 (0.0%)
1	C	0.40	0/3769	0.65	0/5107
1	D	0.39	0/3769	0.64	1/5107 (0.0%)
1	E	0.39	0/3780	0.65	1/5121 (0.0%)
1	F	0.39	0/3769	0.65	1/5107 (0.0%)
All	All	0.39	0/22636	0.65	4/30670 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	97	ASP	N-CA-C	5.69	117.35	111.03
1	F	97	ASP	N-CA-C	5.52	116.99	110.97
1	D	97	ASP	N-CA-C	5.36	116.81	110.97
1	B	97	ASP	N-CA-C	5.21	116.64	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3689	0	3671	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3681	0	3658	10	0
1	C	3681	0	3658	11	0
1	D	3681	0	3660	9	0
1	E	3692	0	3670	11	0
1	F	3681	0	3658	9	0
2	D	18	0	24	0	0
2	E	18	0	24	0	0
2	F	12	0	16	0	0
3	D	1	0	0	0	0
4	A	29	0	0	0	0
4	B	6	0	0	0	0
4	C	4	0	0	0	0
4	D	8	0	0	0	0
4	E	25	0	0	0	0
4	F	14	0	0	0	0
All	All	22240	0	22039	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:16[B]:ARG:HG2	1:E:16[B]:ARG:O	1.99	0.63
1:A:285:TYR:CE2	1:A:287:ILE:HD11	2.35	0.62
1:F:71:VAL:HG13	1:F:308:ILE:HG23	1.82	0.62
1:C:181:PHE:CE2	1:C:185:ILE:HD11	2.40	0.56
1:A:214:PRO:O	1:A:218:TRP:HB2	2.06	0.56
1:A:116:THR:HG21	1:A:155:GLN:HE22	1.72	0.55
1:A:71:VAL:HG13	1:A:308:ILE:HG23	1.91	0.53
1:B:52:VAL:O	1:B:53:ALA:HB3	2.09	0.52
1:E:214:PRO:O	1:E:218:TRP:HB2	2.11	0.51
1:B:181:PHE:CE2	1:B:185:ILE:HD11	2.46	0.51
1:C:71:VAL:HG13	1:C:308:ILE:HG23	1.92	0.51
1:D:326:VAL:HG13	1:D:337:VAL:HG13	1.93	0.51
1:B:251:LEU:CD1	1:B:305:MET:HE2	2.42	0.49
1:D:116:THR:HG21	1:D:155:GLN:HE22	1.77	0.49
1:F:60:ASN:HD22	1:F:61:GLN:N	2.10	0.49
1:D:99:LEU:O	1:D:102:MET:N	2.45	0.49
1:F:214:PRO:O	1:F:218:TRP:HB2	2.14	0.48
1:A:459:PHE:HB3	1:B:365:THR:HB	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:295:CYS:O	1:B:297:ALA:N	2.47	0.47
1:F:137:ALA:O	1:F:138:ALA:HB3	2.15	0.47
1:C:14:VAL:HG13	1:C:157:PHE:HB2	1.97	0.46
1:E:420:THR:HG21	1:E:446:PRO:HD2	1.98	0.46
1:A:285:TYR:CD2	1:A:287:ILE:HD11	2.52	0.46
1:A:12:ILE:HD12	1:A:279:ALA:HB1	1.98	0.45
1:D:51:GLU:O	1:D:55:SER:N	2.49	0.45
1:B:116:THR:OG1	1:B:155:GLN:NE2	2.47	0.45
1:D:181:PHE:CE2	1:D:185:ILE:HD11	2.51	0.45
1:D:115:TYR:HA	1:D:120:MET:HE3	1.97	0.45
1:A:144:MET:HE1	1:A:183:HIS:HA	1.99	0.44
1:E:16[B]:ARG:O	1:E:16[B]:ARG:CG	2.65	0.44
1:E:56:LYS:HB2	1:E:57:HIS:HA	1.99	0.44
1:C:251:LEU:CD1	1:C:305:MET:HE2	2.48	0.44
1:B:339:THR:HG21	1:B:343:LEU:HD21	2.00	0.44
1:C:137:ALA:O	1:C:138:ALA:HB3	2.18	0.44
1:A:245:VAL:HG23	1:A:274:ALA:HB1	2.00	0.44
1:D:118:LEU:HB2	1:D:120:MET:HE2	2.00	0.43
1:A:406:TYR:O	1:A:415:LYS:NZ	2.51	0.43
1:B:79:VAL:HG12	1:B:206:LEU:HD12	1.99	0.43
1:E:404:LEU:HD21	1:E:419:ILE:HG21	2.00	0.43
1:A:12:ILE:HD13	1:A:113:LEU:HD11	2.01	0.43
1:F:100:GLU:HA	1:F:101:GLY:HA2	1.78	0.43
1:C:441:VAL:O	1:C:441:VAL:HG12	2.17	0.43
1:A:278:HIS:CD2	1:A:287:ILE:HD13	2.54	0.43
1:C:330:SER:HB3	1:C:335:LEU:HD22	2.01	0.42
1:F:116:THR:HG21	1:F:155:GLN:HE22	1.85	0.42
1:E:181:PHE:CE2	1:E:185:ILE:HD11	2.54	0.42
1:D:424:VAL:CG2	1:D:451:VAL:HG13	2.50	0.42
1:E:307:ASN:HB3	1:E:328:VAL:HG11	2.01	0.42
1:D:295:CYS:HA	1:D:296:ALA:HA	1.76	0.42
1:E:242:THR:HG21	1:E:357:TYR:HA	2.01	0.41
1:E:100:GLU:HB3	1:E:101:GLY:HA3	2.02	0.41
1:C:214:PRO:O	1:C:218:TRP:HB2	2.20	0.41
1:E:102:MET:O	1:E:103:ASP:CG	2.64	0.41
1:C:294:GLY:HA2	1:C:295:CYS:HA	1.89	0.41
1:B:97:ASP:N	1:B:98:GLY:HA3	2.36	0.41
1:F:129:GLU:O	1:F:130:THR:OG1	2.34	0.40
1:C:363:ASN:N	1:C:363:ASN:HD22	2.19	0.40
1:F:436:ALA:N	1:F:437:PRO:CD	2.85	0.40
1:B:436:ALA:N	1:B:437:PRO:CD	2.84	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:116:THR:HG21	1:C:155:GLN:HE22	1.87	0.40
1:F:296:ALA:O	1:F:297:ALA:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	459/460 (100%)	439 (96%)	20 (4%)	0	100	100
1	B	458/460 (100%)	414 (90%)	37 (8%)	7 (2%)	8	28
1	C	458/460 (100%)	427 (93%)	29 (6%)	2 (0%)	30	60
1	D	458/460 (100%)	437 (95%)	20 (4%)	1 (0%)	43	72
1	E	459/460 (100%)	435 (95%)	22 (5%)	2 (0%)	30	60
1	F	458/460 (100%)	431 (94%)	26 (6%)	1 (0%)	43	72
All	All	2750/2760 (100%)	2583 (94%)	154 (6%)	13 (0%)	24	55

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	296	ALA
1	B	442	GLY
1	B	144	MET
1	B	366	SER
1	F	366	SER
1	C	243	HIS
1	E	296	ALA
1	E	229	GLU
1	B	371	GLU
1	B	428	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	98	GLY
1	C	98	GLY
1	D	98	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/399 (100%)	400 (100%)	0	100	100
1	B	399/399 (100%)	398 (100%)	1 (0%)	86	95
1	C	399/399 (100%)	397 (100%)	2 (0%)	81	93
1	D	399/399 (100%)	398 (100%)	1 (0%)	86	95
1	E	400/399 (100%)	397 (99%)	3 (1%)	73	90
1	F	399/399 (100%)	395 (99%)	4 (1%)	68	88
All	All	2396/2394 (100%)	2385 (100%)	11 (0%)	84	93

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	142	GLU
1	C	307	ASN
1	C	319	LYS
1	D	21	ARG
1	E	16[A]	ARG
1	E	16[B]	ARG
1	E	103	ASP
1	F	26	ARG
1	F	58	THR
1	F	335	LEU
1	F	395	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	HIS
1	A	146	ASN
1	A	155	GLN
1	A	217	HIS
1	B	155	GLN
1	B	169	GLN
1	B	199	GLN
1	B	425	HIS
1	C	146	ASN
1	C	155	GLN
1	C	217	HIS
1	C	226	GLN
1	C	344	ASN
1	C	363	ASN
1	C	425	HIS
1	D	13	HIS
1	D	145	ASN
1	D	155	GLN
1	D	217	HIS
1	D	226	GLN
1	D	425	HIS
1	E	146	ASN
1	E	155	GLN
1	F	30	GLN
1	F	155	GLN
1	F	278	HIS
1	F	342	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	E	502	-	5,5,5	0.28	0	5,5,5	0.21	0
2	GOL	D	503	-	5,5,5	0.28	0	5,5,5	0.27	0
2	GOL	F	502	-	5,5,5	0.28	0	5,5,5	0.24	0
2	GOL	F	501	-	5,5,5	0.30	0	5,5,5	0.15	0
2	GOL	D	501	-	5,5,5	0.30	0	5,5,5	0.19	0
2	GOL	E	501	-	5,5,5	0.25	0	5,5,5	0.33	0
2	GOL	E	503	-	5,5,5	0.31	0	5,5,5	0.19	0
2	GOL	D	502	-	5,5,5	0.26	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	E	502	-	-	0/4/4/4	-
2	GOL	D	503	-	-	1/4/4/4	-
2	GOL	F	502	-	-	0/4/4/4	-
2	GOL	F	501	-	-	1/4/4/4	-
2	GOL	D	501	-	-	2/4/4/4	-
2	GOL	E	501	-	-	0/4/4/4	-
2	GOL	E	503	-	-	1/4/4/4	-
2	GOL	D	502	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	501	GOL	C1-C2-C3-O3
2	D	501	GOL	O2-C2-C3-O3
2	F	501	GOL	C1-C2-C3-O3
2	D	503	GOL	O1-C1-C2-C3
2	E	503	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	460/460 (100%)	0.36	13 (2%)	55	45	41, 73, 103, 121	1 (0%)
1	B	460/460 (100%)	1.08	58 (12%)	8	6	78, 126, 151, 162	0
1	C	460/460 (100%)	0.80	38 (8%)	17	12	75, 107, 134, 146	0
1	D	460/460 (100%)	0.53	21 (4%)	37	29	59, 92, 120, 136	0
1	E	460/460 (100%)	0.28	11 (2%)	59	49	36, 78, 102, 139	1 (0%)
1	F	460/460 (100%)	0.35	14 (3%)	52	42	59, 76, 103, 113	0
All	All	2760/2760 (100%)	0.57	155 (5%)	30	23	36, 89, 137, 162	2 (0%)

All (155) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	TYR	6.5
1	A	205	GLU	5.1
1	B	343	LEU	4.8
1	E	2	ALA	4.4
1	B	275	ILE	4.0
1	B	350	THR	3.9
1	A	296	ALA	3.8
1	D	102	MET	3.7
1	C	141	LEU	3.7
1	D	296	ALA	3.5
1	B	99	LEU	3.4
1	D	365	THR	3.4
1	B	1	GLY	3.3
1	E	296	ALA	3.3
1	B	370	LEU	3.3
1	E	59	SER	3.3
1	E	101	GLY	3.3
1	D	460	TRP	3.3
1	B	444	ILE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	294	GLY	3.2
1	F	98	GLY	3.2
1	A	56	LYS	3.2
1	B	316	LEU	3.2
1	D	199	GLN	3.2
1	F	363	ASN	3.2
1	B	297	ALA	3.2
1	B	366	SER	3.1
1	B	149	PHE	3.1
1	C	362	ALA	3.0
1	A	1	GLY	3.0
1	B	53	ALA	3.0
1	F	296	ALA	3.0
1	B	445	VAL	3.0
1	B	365	THR	3.0
1	D	362	ALA	3.0
1	C	145	ASN	3.0
1	B	412	LEU	3.0
1	C	389	PRO	3.0
1	A	366	SER	2.9
1	B	82	LEU	2.9
1	B	364	LYS	2.9
1	A	58	THR	2.9
1	F	362	ALA	2.9
1	B	372	SER	2.9
1	C	98	GLY	2.8
1	D	350	THR	2.8
1	B	245	VAL	2.8
1	B	252	ALA	2.8
1	B	296	ALA	2.8
1	D	128	TRP	2.8
1	D	98	GLY	2.8
1	E	62	GLU	2.7
1	B	2	ALA	2.7
1	C	12	ILE	2.7
1	B	363	ASN	2.7
1	C	365	THR	2.7
1	D	1	GLY	2.7
1	C	297	ALA	2.7
1	B	285	TYR	2.7
1	E	56	LYS	2.7
1	B	341	TYR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	317	THR	2.7
1	B	144	MET	2.6
1	B	300	MET	2.6
1	B	184	CYS	2.6
1	E	57	HIS	2.6
1	B	88	GLY	2.6
1	C	245	VAL	2.6
1	B	211	GLY	2.6
1	D	285	TYR	2.6
1	F	295	CYS	2.6
1	D	2	ALA	2.6
1	C	135	PRO	2.5
1	E	58	THR	2.5
1	A	102	MET	2.5
1	C	364	LYS	2.5
1	B	233	ASP	2.5
1	C	356	GLY	2.5
1	C	102	MET	2.4
1	B	345	PHE	2.4
1	C	152	ILE	2.4
1	F	152	ILE	2.4
1	B	98	GLY	2.4
1	D	88	GLY	2.4
1	D	366	SER	2.4
1	B	81	ALA	2.4
1	C	296	ALA	2.4
1	A	199	GLN	2.4
1	A	57	HIS	2.4
1	B	384	PHE	2.4
1	C	205	GLU	2.4
1	F	58	THR	2.4
1	C	64	LEU	2.4
1	B	145	ASN	2.3
1	B	360	THR	2.3
1	C	99	LEU	2.3
1	B	319	LYS	2.3
1	B	354	LYS	2.3
1	C	118	LEU	2.3
1	F	166	GLU	2.3
1	C	286	LEU	2.3
1	C	295	CYS	2.3
1	C	458	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	460	TRP	2.3
1	F	360	THR	2.3
1	B	56	LYS	2.3
1	C	57	HIS	2.3
1	D	152	ILE	2.3
1	B	369	PRO	2.3
1	C	267	LYS	2.2
1	B	148	ASP	2.2
1	D	63	THR	2.2
1	C	147	LYS	2.2
1	D	307	ASN	2.2
1	F	99	LEU	2.2
1	B	92	VAL	2.2
1	D	65	PRO	2.2
1	A	363	ASN	2.2
1	B	419	ILE	2.2
1	A	365	THR	2.2
1	B	339	THR	2.2
1	A	362	ALA	2.2
1	F	86	ASP	2.2
1	B	117	THR	2.2
1	C	58	THR	2.2
1	C	291	LEU	2.2
1	D	237	SER	2.2
1	B	344	ASN	2.1
1	C	18	THR	2.1
1	C	143	LYS	2.1
1	E	99	LEU	2.1
1	C	1	GLY	2.1
1	A	53	ALA	2.1
1	F	199	GLN	2.1
1	F	57	HIS	2.1
1	B	298	THR	2.1
1	B	287	ILE	2.1
1	B	65	PRO	2.1
1	B	446	PRO	2.1
1	C	114	PRO	2.1
1	B	3	LEU	2.1
1	C	239	PHE	2.1
1	B	411	THR	2.1
1	D	58	THR	2.1
1	C	312	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	105	MET	2.0
1	B	57	HIS	2.0
1	B	266	VAL	2.0
1	C	117	THR	2.0
1	C	359	ILE	2.0
1	B	362	ALA	2.0
1	D	56	LYS	2.0
1	E	152	ILE	2.0
1	F	175	ILE	2.0
1	E	102	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	D	502	6/6	0.85	0.27	88,89,89,90	0
2	GOL	F	502	6/6	0.85	0.19	82,82,82,82	0
2	GOL	E	501	6/6	0.86	0.20	92,94,95,96	0
2	GOL	D	503	6/6	0.87	0.20	91,92,93,93	0
2	GOL	E	503	6/6	0.88	0.19	102,103,104,104	0
2	GOL	F	501	6/6	0.88	0.18	106,107,108,108	0
2	GOL	E	502	6/6	0.88	0.15	126,127,127,128	0
2	GOL	D	501	6/6	0.89	0.15	89,90,90,90	0
3	CL	D	504	1/1	0.95	0.08	70,70,70,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.