



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:10 PM UTC

PDB ID : 5NZL / pdb_00005nzl
Title : Crystal structure of UDP-glucose pyrophosphorylase from Leishmania major in complex with resveratrol
Authors : Cramer, J.T.; Fuehring, J.I.; Baruch, P.; Bruetting, C.; Hesse, R.; Knoelker, H.-J.; Gerardy-Schahn, R.; Fedorov, R.
Deposited on : 2017-05-14
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	9% 79% 13% • •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-glucose pyrophosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	483	Total	C	N	O	S	0	0	0
			3730	2351	636	717	26			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	495	MET	-	expression tag	UNP Q4QDU3
A	496	ARG	-	expression tag	UNP Q4QDU3
A	497	PRO	-	expression tag	UNP Q4QDU3
A	498	LEU	-	expression tag	UNP Q4QDU3
A	499	GLU	-	expression tag	UNP Q4QDU3
A	500	HIS	-	expression tag	UNP Q4QDU3
A	501	HIS	-	expression tag	UNP Q4QDU3
A	502	HIS	-	expression tag	UNP Q4QDU3
A	503	HIS	-	expression tag	UNP Q4QDU3
A	504	HIS	-	expression tag	UNP Q4QDU3
A	505	HIS	-	expression tag	UNP Q4QDU3

- Molecule 2 is RESVERATROL (CCD ID: STL) (formula: C₁₄H₁₂O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			17	14	3		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	72	Total O 72 72	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UDP-glucose pyrophosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.52Å 108.45Å 153.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.36 – 2.40 47.36 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.36-2.40) 99.9 (47.36-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.39Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.224 , 0.249 0.234 , 0.251	Depositor DCC
R_{free} test set	1197 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	50.0	Xtriage
Anisotropy	0.428	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3851	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, STL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	4/3800 (0.1%)	0.55	6/5147 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	419	ALA	CA-C	-8.91	1.41	1.52
1	A	421	TYR	C-O	-7.36	1.14	1.24
1	A	417	ASP	C-O	-6.49	1.16	1.23
1	A	57	SER	C-O	-5.13	1.18	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	421	TYR	N-CA-C	16.10	132.72	112.92
1	A	472	ALA	N-CA-C	11.59	123.47	111.07
1	A	473	PHE	N-CA-C	7.84	122.20	111.39
1	A	57	SER	N-CA-C	6.22	119.70	108.69
1	A	55	VAL	N-CA-C	6.17	116.96	110.72
1	A	417	ASP	N-CA-C	5.24	117.83	110.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3730	0	3720	50	4
2	A	17	0	10	5	0
3	A	32	0	48	3	0
4	A	72	0	0	5	2
All	All	3851	0	3778	50	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:420:HIS:NE2	1:A:449:LEU:HB2	1.21	1.48
1:A:420:HIS:CD2	1:A:449:LEU:HA	1.46	1.48
1:A:420:HIS:CD2	1:A:449:LEU:CA	2.13	1.30
1:A:420:HIS:NE2	1:A:449:LEU:CB	1.95	1.30
1:A:419:ALA:HB1	4:A:731:HOH:O	1.01	1.16
1:A:420:HIS:NE2	1:A:449:LEU:CA	2.21	1.02
1:A:420:HIS:CD2	1:A:449:LEU:CB	2.45	0.95
1:A:417:ASP:OD1	1:A:448:GLY:N	2.00	0.94
1:A:420:HIS:CE1	1:A:449:LEU:HB2	2.07	0.90
1:A:420:HIS:HD2	1:A:449:LEU:HA	0.78	0.88
1:A:420:HIS:CD2	1:A:449:LEU:HB2	2.08	0.83
1:A:419:ALA:CB	4:A:731:HOH:O	1.80	0.70
1:A:420:HIS:NE2	1:A:449:LEU:N	2.40	0.69
1:A:69:ASP:OD2	4:A:701:HOH:O	2.11	0.68
1:A:405:ASP:OD1	4:A:702:HOH:O	2.13	0.66
1:A:420:HIS:CE1	4:A:731:HOH:O	2.51	0.64
1:A:374:SER:HB3	2:A:601:STL:H11	1.80	0.62
1:A:248:ARG:HB2	2:A:601:STL:H2	1.81	0.62
1:A:439:VAL:HG13	1:A:440:GLU:HG2	1.80	0.62
1:A:467:THR:HG23	1:A:467:THR:O	2.01	0.60
1:A:52:ILE:HD11	1:A:281:LEU:HD11	1.84	0.59
1:A:372:PRO:HA	2:A:601:STL:H6	1.86	0.57
1:A:420:HIS:CD2	1:A:449:LEU:N	2.71	0.57
1:A:473:PHE:C	1:A:473:PHE:CD1	2.85	0.54
1:A:149:TRP:CD1	1:A:149:TRP:H	2.27	0.53
1:A:373:ARG:H	2:A:601:STL:C7	2.21	0.53
1:A:474:VAL:HG12	1:A:475:ILE:N	2.24	0.52
1:A:290:LYS:HE2	1:A:290:LYS:H	1.77	0.49
1:A:126:ARG:HD2	1:A:212:TYR:CZ	2.47	0.49
1:A:339:VAL:HG12	1:A:346:SER:HB2	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:421:TYR:CD1	1:A:421:TYR:N	2.81	0.48
1:A:6:LYS:C	1:A:8:LEU:H	2.22	0.47
1:A:416:LEU:HA	1:A:416:LEU:HD12	1.70	0.47
1:A:373:ARG:H	2:A:601:STL:H7	1.80	0.47
1:A:404:ASP:OD1	1:A:405:ASP:N	2.47	0.47
1:A:91:LEU:HD21	1:A:98:LEU:HD13	1.99	0.45
1:A:370:VAL:HG12	3:A:604:EDO:H11	1.98	0.45
1:A:19:MET:HE2	1:A:29:ILE:HG12	1.99	0.44
1:A:80:LYS:HG3	1:A:110:THR:HG21	1.99	0.44
1:A:484:THR:OG1	1:A:485:THR:N	2.50	0.44
1:A:245:GLU:HB2	1:A:309:ASN:HB2	1.98	0.43
1:A:161:MET:H	3:A:603:EDO:H22	1.84	0.42
1:A:46:SER:OG	1:A:289:PRO:HG3	2.19	0.42
1:A:100:VAL:O	1:A:389:ARG:HG2	2.19	0.42
1:A:148:PRO:O	1:A:152:GLN:HG2	2.19	0.42
1:A:415:ASP:O	1:A:416:LEU:HD12	2.20	0.41
1:A:91:LEU:HD12	1:A:91:LEU:HA	1.83	0.41
1:A:396:THR:OG1	1:A:398:ASP:OD1	2.36	0.41
1:A:118:ARG:HA	1:A:122:SER:O	2.20	0.41
1:A:201:SER:HB3	3:A:603:EDO:H21	2.02	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418:SER:CB	1:A:418:SER:CB[3_554]	1.37	0.83
1:A:92:CYS:CB	1:A:92:CYS:SG[3_554]	1.81	0.39
1:A:49:ASP:O	4:A:701:HOH:O[8_455]	2.14	0.06
1:A:52:ILE:O	4:A:701:HOH:O[8_455]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/505 (95%)	468 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/432 (95%)	373 (91%)	37 (9%)	9	15

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	49	ASP
1	A	52	ILE
1	A	55	VAL
1	A	61	LEU
1	A	87	THR
1	A	91	LEU
1	A	92	CYS
1	A	101	LYS
1	A	126	ARG
1	A	129	LEU
1	A	138	SER
1	A	194	ILE
1	A	213	ARG
1	A	238	GLU
1	A	254	LYS
1	A	255	LYS
1	A	277	LYS
1	A	278	ARG
1	A	280	LEU
1	A	284	GLU
1	A	290	LYS
1	A	310	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	318	LEU
1	A	335	ASN
1	A	348	LYS
1	A	406	ARG
1	A	416	LEU
1	A	418	SER
1	A	420	HIS
1	A	421	TYR
1	A	423	MET
1	A	444	VAL
1	A	471	SER
1	A	480	LYS
1	A	483	ASP
1	A	484	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	163	ASN
1	A	306	ASN
1	A	351	GLN
1	A	482	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	603	-	3,3,3	0.33	0	2,2,2	0.27	0
3	EDO	A	608	-	3,3,3	0.28	0	2,2,2	0.99	0
3	EDO	A	607	-	3,3,3	0.25	0	2,2,2	0.96	0
3	EDO	A	609	-	3,3,3	0.28	0	2,2,2	0.29	0
3	EDO	A	606	-	3,3,3	0.29	0	2,2,2	0.19	0
3	EDO	A	605	-	3,3,3	0.27	0	2,2,2	0.69	0
2	STL	A	601	-	18,18,18	1.87	4 (22%)	24,24,24	1.38	2 (8%)
3	EDO	A	604	-	3,3,3	0.24	0	2,2,2	0.52	0
3	EDO	A	602	-	3,3,3	0.32	0	2,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	603	-	-	0/1/1/1	-
3	EDO	A	608	-	-	0/1/1/1	-
3	EDO	A	607	-	-	1/1/1/1	-
3	EDO	A	609	-	-	1/1/1/1	-
3	EDO	A	606	-	-	0/1/1/1	-
3	EDO	A	605	-	-	1/1/1/1	-
2	STL	A	601	-	-	3/5/5/5	0/2/2/2
3	EDO	A	604	-	-	1/1/1/1	-
3	EDO	A	602	-	-	1/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	STL	O2-C3	4.57	1.47	1.37
2	A	601	STL	C4-C5	-2.95	1.34	1.39
2	A	601	STL	C6-C5	-2.84	1.35	1.39
2	A	601	STL	C5-C7	2.43	1.54	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	STL	C9-C8-C7	-4.18	108.75	125.89
2	A	601	STL	C5-C6-C1	2.33	122.21	120.26

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	STL	C7-C8-C9-C10
2	A	601	STL	C7-C8-C9-C14
3	A	609	EDO	O1-C1-C2-O2
3	A	607	EDO	O1-C1-C2-O2
3	A	604	EDO	O1-C1-C2-O2
3	A	605	EDO	O1-C1-C2-O2
3	A	602	EDO	O1-C1-C2-O2
2	A	601	STL	C4-C5-C7-C8

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	EDO	2	0
2	A	601	STL	5	0
3	A	604	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	483/505 (95%)	0.50	45 (9%) 14 11	30, 53, 88, 122	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	419	ALA	6.4
1	A	420	HIS	5.3
1	A	457	VAL	4.5
1	A	472	ALA	4.3
1	A	448	GLY	4.1
1	A	452	PHE	3.9
1	A	481	LEU	3.8
1	A	470	ALA	3.8
1	A	471	SER	3.7
1	A	458	LEU	3.5
1	A	488	PRO	3.5
1	A	418	SER	3.5
1	A	475	ILE	3.4
1	A	474	VAL	3.3
1	A	464	ILE	3.3
1	A	6	LYS	3.0
1	A	449	LEU	3.0
1	A	468	ASP	3.0
1	A	450	VAL	3.0
1	A	473	PHE	2.9
1	A	479	ALA	2.9
1	A	446	VAL	2.8
1	A	273	PRO	2.8
1	A	413	VAL	2.8
1	A	438	LEU	2.8
1	A	486	ALA	2.7
1	A	462	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	429	LYS	2.6
1	A	269	LYS	2.5
1	A	466	ASN	2.4
1	A	487	SER	2.4
1	A	434	GLY	2.4
1	A	433	HIS	2.4
1	A	456	ASN	2.4
1	A	444	VAL	2.2
1	A	467	THR	2.2
1	A	485	THR	2.2
1	A	469	SER	2.2
1	A	149	TRP	2.2
1	A	454	ALA	2.2
1	A	459	THR	2.1
1	A	439	VAL	2.1
1	A	430	LEU	2.1
1	A	267	LYS	2.0
1	A	9	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	STL	A	601	17/17	0.55	0.23	66,72,83,85	0
3	EDO	A	602	4/4	0.70	0.20	70,71,76,81	0
3	EDO	A	603	4/4	0.71	0.20	60,63,67,72	0
3	EDO	A	609	4/4	0.73	0.20	62,65,65,81	0
3	EDO	A	607	4/4	0.85	0.17	58,60,65,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	608	4/4	0.86	0.18	46,56,57,64	0
3	EDO	A	604	4/4	0.87	0.16	56,58,67,71	0
3	EDO	A	606	4/4	0.89	0.14	55,60,66,76	0
3	EDO	A	605	4/4	0.91	0.16	45,50,51,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.