



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 09:05 PM UTC

PDB ID : 7NZF / pdb_00007nzf
Title : Crystal structure of HLA-DR4 in complex with a mutated human collagen type II peptide
Authors : Ge, C.; Dobritsch, D.; Holmdahl, R.
Deposited on : 2021-03-24
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	180	3% 88% 12%
2	BBB	191	12% 70% 20% 7%
3	CCC	13	15% 100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6129 atoms, of which 2908 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	180	Total	C	H	N	O	S	36	0	0
			2906	956	1430	241	274	5			

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DR beta chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	BBB	177	Total	C	H	N	O	S	46	0	0
			2841	928	1378	255	275	5			

- Molecule 3 is a protein called mutant human collagen type II,259-273.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	CCC	13	Total	C	H	N	O		0	0	0
			155	51	72	14	18				

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	AAA	1	Total	C	H	N	O	3	0
			28	8	14	1	5		
4	AAA	1	Total	C	H	N	O	3	0
			28	8	14	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	94	Total	O	0	0
			94	94		
5	BBB	74	Total	O	0	0
			74	74		
5	CCC	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain AAA: 




- Molecule 2: HLA class II histocompatibility antigen, DR beta chain

Chain BBB: 



- Molecule 3: mutant human collagen type II,259-273

Chain CCC: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.44Å 71.44Å 138.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.95 – 1.90 36.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.4 (36.95-1.90) 98.4 (36.95-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.87 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.221 , 0.267 0.222 , 0.267	Depositor DCC
R_{free} test set	1564 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6129	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.02	0/1520	1.26	4/2072 (0.2%)
2	BBB	1.01	0/1501	1.34	5/2038 (0.2%)
3	CCC	0.93	0/85	1.40	0/114
All	All	1.01	0/3106	1.30	9/4224 (0.2%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	93	THR	CA-C-N	7.87	130.83	120.28
1	AAA	93	THR	C-N-CA	7.87	130.83	120.28
2	BBB	145	THR	N-CA-CB	-7.47	98.70	110.28
1	AAA	180	PHE	CA-C-O	-7.01	108.88	120.80
2	BBB	93	ARG	CG-CD-NE	-6.06	98.67	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1476	1430	1424	12	1
2	BBB	1463	1378	1365	30	1
3	CCC	83	72	69	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	AAA	28	28	26	0	0
5	AAA	94	0	0	2	0
5	BBB	74	0	0	5	0
5	CCC	3	0	0	0	0
All	All	3221	2908	2884	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:BBB:104:ALA:O	5:BBB:201:HOH:O	1.87	0.92
2:BBB:38:VAL:HG13	2:BBB:54:GLY:HA3	1.61	0.81
2:BBB:113:ASN:CB	5:BBB:257:HOH:O	2.43	0.66
2:BBB:133:ARG:NH2	5:BBB:203:HOH:O	2.29	0.64
2:BBB:138:GLU:HA	2:BBB:138:GLU:OE1	2.02	0.59

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:50:ARG:HE	2:BBB:69:GLU:OE1[6_654]	1.57	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	178/180 (99%)	177 (99%)	1 (1%)	0	100 100
2	BBB	171/191 (90%)	161 (94%)	8 (5%)	2 (1%)	10 4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	CCC	11/13 (85%)	11 (100%)	0	0	100	100
All	All	360/384 (94%)	349 (97%)	9 (2%)	2 (1%)	21	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	BBB	52	GLU
2	BBB	50	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	164/164 (100%)	160 (98%)	4 (2%)	43	38
2	BBB	157/172 (91%)	150 (96%)	7 (4%)	24	17
3	CCC	6/6 (100%)	6 (100%)	0	100	100
All	All	327/342 (96%)	316 (97%)	11 (3%)	32	25

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	BBB	88	SER
2	BBB	136	GLN
2	BBB	172	THR
2	BBB	145	THR
2	BBB	28	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	AAA	202	1	14,14,15	0.58	0	17,19,21	1.78	3 (17%)
4	NAG	AAA	201	1	14,14,15	0.69	0	17,19,21	2.26	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	AAA	202	1	-	0/6/23/26	0/1/1/1
4	NAG	AAA	201	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	AAA	202	NAG	C1-O5-C5	5.64	119.74	112.19
4	AAA	201	NAG	C2-N2-C7	5.29	129.99	122.90
4	AAA	201	NAG	C1-O5-C5	4.67	118.44	112.19
4	AAA	201	NAG	C1-C2-N2	3.93	116.62	110.43
4	AAA	201	NAG	O5-C5-C6	3.05	113.60	107.66

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	AAA	201	NAG	O5-C5-C6-O6
4	AAA	201	NAG	C4-C5-C6-O6
4	AAA	201	NAG	C1-C2-N2-C7
4	AAA	201	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	180/180 (100%)	0.12	5 (2%) 55 59	16, 28, 52, 103	0
2	BBB	177/191 (92%)	0.83	23 (12%) 7 7	15, 37, 75, 90	4 (2%)
3	CCC	13/13 (100%)	1.42	2 (15%) 5 5	36, 44, 69, 72	0
All	All	370/384 (96%)	0.50	30 (8%) 18 19	15, 32, 71, 103	4 (1%)

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	BBB	23	ARG	8.5
2	BBB	189	ARG	8.0
1	AAA	1	ILE	7.3
2	BBB	22	GLU	6.9
2	BBB	19	ASN	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	AAA	201	14/15	0.66	0.19	30,80,83,88	3
4	NAG	AAA	202	14/15	0.87	0.10	30,42,49,50	3

6.5 Other polymers [i](#)

There are no such residues in this entry.