



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:10 PM UTC

PDB ID : 5O0Q / pdb_00005o0q
Title : Deglycosylated Nogo Receptor with native disulfide structure
Authors : Pronker, M.F.; Janssen, B.J.C.
Deposited on : 2017-05-16
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	3% 79% 8% 13%
1	B	334	3% 78% 9% 13%
1	C	334	4% 76% 10% 13%
1	D	334	3% 82% 5% 13%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reticulon-4 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2274	1436	426	397	15			
1	B	291	Total	C	N	O	S	0	0	0
			2282	1442	427	398	15			
1	C	290	Total	C	N	O	S	0	0	0
			2274	1436	426	397	15			
1	D	292	Total	C	N	O	S	0	0	0
			2289	1447	428	399	15			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	24	GLY	-	expression tag	UNP Q99PI8
A	25	SER	-	expression tag	UNP Q99PI8
A	349	ALA	-	expression tag	UNP Q99PI8
A	350	ALA	-	expression tag	UNP Q99PI8
A	351	ALA	-	expression tag	UNP Q99PI8
A	352	HIS	-	expression tag	UNP Q99PI8
A	353	HIS	-	expression tag	UNP Q99PI8
A	354	HIS	-	expression tag	UNP Q99PI8
A	355	HIS	-	expression tag	UNP Q99PI8
A	356	HIS	-	expression tag	UNP Q99PI8
A	357	HIS	-	expression tag	UNP Q99PI8
B	24	GLY	-	expression tag	UNP Q99PI8
B	25	SER	-	expression tag	UNP Q99PI8
B	349	ALA	-	expression tag	UNP Q99PI8
B	350	ALA	-	expression tag	UNP Q99PI8
B	351	ALA	-	expression tag	UNP Q99PI8
B	352	HIS	-	expression tag	UNP Q99PI8
B	353	HIS	-	expression tag	UNP Q99PI8
B	354	HIS	-	expression tag	UNP Q99PI8
B	355	HIS	-	expression tag	UNP Q99PI8
B	356	HIS	-	expression tag	UNP Q99PI8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	357	HIS	-	expression tag	UNP Q99PI8
C	24	GLY	-	expression tag	UNP Q99PI8
C	25	SER	-	expression tag	UNP Q99PI8
C	349	ALA	-	expression tag	UNP Q99PI8
C	350	ALA	-	expression tag	UNP Q99PI8
C	351	ALA	-	expression tag	UNP Q99PI8
C	352	HIS	-	expression tag	UNP Q99PI8
C	353	HIS	-	expression tag	UNP Q99PI8
C	354	HIS	-	expression tag	UNP Q99PI8
C	355	HIS	-	expression tag	UNP Q99PI8
C	356	HIS	-	expression tag	UNP Q99PI8
C	357	HIS	-	expression tag	UNP Q99PI8
D	24	GLY	-	expression tag	UNP Q99PI8
D	25	SER	-	expression tag	UNP Q99PI8
D	349	ALA	-	expression tag	UNP Q99PI8
D	350	ALA	-	expression tag	UNP Q99PI8
D	351	ALA	-	expression tag	UNP Q99PI8
D	352	HIS	-	expression tag	UNP Q99PI8
D	353	HIS	-	expression tag	UNP Q99PI8
D	354	HIS	-	expression tag	UNP Q99PI8
D	355	HIS	-	expression tag	UNP Q99PI8
D	356	HIS	-	expression tag	UNP Q99PI8
D	357	HIS	-	expression tag	UNP Q99PI8

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	131	Total	O	0	0
			131	131		
6	B	35	Total	O	0	0
			35	35		
6	C	92	Total	O	0	0
			92	92		
6	D	75	Total	O	0	0
			75	75		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Reticulon-4 receptor




• Molecule 1: Reticulon-4 receptor




• Molecule 1: Reticulon-4 receptor



• Molecule 1: Reticulon-4 receptor





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	132.12Å 46.33Å 132.23Å 90.00° 91.93° 90.00°	Depositor
Resolution (Å)	45.94 – 2.50 45.94 – 2.50	Depositor EDS
% Data completeness (in resolution range)	94.3 (45.94-2.50) 94.3 (45.94-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.211 , 0.261 0.212 , 0.262	Depositor DCC
R_{free} test set	2775 reflections (3.81%)	wwPDB-VP
Wilson B-factor (Å ²)	40.6	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.088 for l,k,-h 0.026 for h,-k,-l 0.024 for l,-k,h	Xtriage
F_o , F_c correlation	0.93	EDS
Total number of atoms	9680	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.10 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.1439e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/2332	0.35	0/3175
1	B	0.13	0/2340	0.32	0/3187
1	C	0.14	0/2331	0.33	0/3174
1	D	0.14	0/2347	0.34	0/3197
All	All	0.14	0/9350	0.33	0/12733

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2274	0	2245	13	0
1	B	2282	0	2256	13	0
1	C	2274	0	2246	20	0
1	D	2289	0	2264	7	0
2	A	42	0	39	0	0
2	B	28	0	26	0	0
2	C	42	0	39	0	0
2	D	42	0	39	1	0
3	A	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	0	0	0
3	C	10	0	0	0	0
3	D	15	0	0	0	0
4	A	6	0	8	0	0
4	C	12	0	16	1	0
5	A	1	0	0	0	0
6	A	131	0	0	2	0
6	B	35	0	0	0	0
6	C	92	0	0	3	0
6	D	75	0	0	0	0
All	All	9680	0	9178	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:282:SER:H	4:C:406:GOL:H11	1.56	0.69
1:A:95:ARG:NH1	6:A:501:HOH:O	2.29	0.65
1:A:38:LYS:HB2	1:A:61:ARG:HD3	1.87	0.56
2:D:401:NAG:H83	2:D:401:NAG:H3	1.87	0.55
1:A:275:LEU:HD13	1:A:299:LYS:HG2	1.88	0.55
1:D:298:LEU:HD11	1:D:306:LEU:HD21	1.91	0.53
1:B:201:LEU:HD13	1:B:204:LEU:HD22	1.90	0.52
1:C:206:ARG:NH2	6:C:505:HOH:O	2.42	0.49
1:B:61:ARG:HG2	1:B:85:ILE:HB	1.94	0.49
1:A:193:VAL:HB	1:A:217:VAL:HG13	1.95	0.48
1:A:206:ARG:NH1	6:A:508:HOH:O	2.46	0.48
1:B:38:LYS:HB2	1:B:61:ARG:HD3	1.96	0.48
1:B:138:ASP:OD1	1:B:138:ASP:N	2.44	0.48
1:C:287:CYS:HB2	1:C:293:LEU:O	2.14	0.48
1:D:272:TRP:O	1:D:276:GLN:HG2	2.13	0.47
1:D:165:ASN:OD1	1:D:189:ARG:NH2	2.48	0.47
1:C:208:LEU:HD22	1:C:232:TYR:CD2	2.50	0.47
1:C:272:TRP:O	1:C:276:GLN:HG2	2.16	0.46
1:B:275:LEU:HD13	1:B:299:LYS:HG2	1.98	0.45
1:C:285:VAL:HB	1:C:298:LEU:HB3	1.97	0.45
1:C:247:MET:HE3	1:C:274:TRP:HD1	1.82	0.45
1:B:124:THR:HA	1:B:127:HIS:CD2	2.52	0.45
1:B:235:ALA:H	1:B:259:ASP:HB3	1.82	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:222:PHE:HB3	1:D:249:LEU:HD21	1.99	0.44
1:C:64:LEU:HB2	1:C:88:LEU:HD23	2.00	0.44
1:B:287:CYS:HB2	1:B:293:LEU:O	2.18	0.43
1:B:332:LEU:HB2	1:B:333:PRO:HD3	1.99	0.43
1:A:134:THR:HG22	1:A:158:TYR:HB2	2.01	0.43
1:C:204:LEU:HD21	1:C:207:LEU:HB2	2.00	0.43
1:A:138:ASP:OD1	1:A:138:ASP:N	2.45	0.43
1:A:272:TRP:O	1:A:276:GLN:HG2	2.19	0.43
1:C:84:THR:HA	1:C:107:LEU:HA	1.99	0.43
1:C:111:ASP:HA	1:C:136:HIS:HB2	2.01	0.43
1:A:28:PRO:HG3	1:A:54:GLY:O	2.18	0.42
1:C:189:ARG:NH1	6:C:508:HOH:O	2.43	0.42
1:C:182:HIS:HB3	1:C:184:PHE:CE1	2.54	0.42
1:D:101:PHE:HB3	1:D:104:LEU:HD12	2.01	0.42
1:C:174:PHE:HB3	1:C:177:LEU:HD12	2.00	0.42
1:B:201:LEU:HB3	1:B:204:LEU:HB2	2.02	0.42
1:C:194:PRO:HG2	1:C:197:ALA:HB2	2.01	0.42
1:A:93:LEU:HD13	1:A:96:ILE:HD11	2.02	0.42
1:A:287:CYS:HB2	1:A:293:LEU:O	2.20	0.41
1:C:247:MET:HE3	1:C:274:TRP:CD1	2.55	0.41
1:C:257:LEU:HB3	1:C:262:TRP:HZ2	1.86	0.41
1:C:297:ASP:O	1:C:301:LEU:HG	2.20	0.41
1:C:49:GLN:NE2	6:C:504:HOH:O	2.39	0.41
1:B:257:LEU:HB3	1:B:262:TRP:HZ2	1.85	0.41
1:D:122:ASP:HA	1:D:123:PRO:HD3	1.94	0.41
1:A:235:ALA:H	1:A:259:ASP:HB3	1.85	0.41
1:A:269:ARG:HA	1:A:306:LEU:HD23	2.02	0.41
1:D:265:ASP:HB2	1:D:308:GLY:O	2.20	0.41
1:B:242:PRO:O	1:B:245:VAL:HG12	2.20	0.41
1:C:134:THR:HG22	1:C:158:TYR:HB2	2.03	0.40
1:B:198:PHE:HZ	1:B:209:LEU:HD11	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/334 (86%)	270 (94%)	16 (6%)	0	100	100
1	B	287/334 (86%)	274 (96%)	13 (4%)	0	100	100
1	C	286/334 (86%)	263 (92%)	23 (8%)	0	100	100
1	D	288/334 (86%)	270 (94%)	18 (6%)	0	100	100
All	All	1147/1336 (86%)	1077 (94%)	70 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/283 (87%)	242 (98%)	5 (2%)	48	75
1	B	248/283 (88%)	239 (96%)	9 (4%)	31	58
1	C	247/283 (87%)	244 (99%)	3 (1%)	63	83
1	D	249/283 (88%)	245 (98%)	4 (2%)	55	79
All	All	991/1132 (88%)	970 (98%)	21 (2%)	47	74

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	GLU
1	A	43	CYS
1	A	120	VAL
1	A	138	ASP
1	A	335	CYS
1	B	43	CYS
1	B	138	ASP
1	B	144	GLU
1	B	151	ARG
1	B	165	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	169	LEU
1	B	237	ASN
1	B	250	ARG
1	B	309	CYS
1	C	120	VAL
1	C	139	ARG
1	C	247	MET
1	D	43	CYS
1	D	138	ASP
1	D	139	ARG
1	D	299	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	119	HIS
1	B	65	HIS
1	B	202	HIS
1	B	288	ASN
1	C	196	HIS
1	C	220	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 1 is monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	405	-	4,4,4	0.95	0	6,6,6	0.45	0
3	PO4	D	406	-	4,4,4	0.95	0	6,6,6	0.45	0
4	GOL	A	408	-	5,5,5	0.36	0	5,5,5	0.34	0
2	NAG	B	402	1	14,14,15	0.26	0	17,19,21	0.43	0
3	PO4	C	405	-	4,4,4	0.95	0	6,6,6	0.46	0
2	NAG	D	401	1	14,14,15	0.51	0	17,19,21	1.31	2 (11%)
2	NAG	A	402	1	14,14,15	0.27	0	17,19,21	0.47	0
3	PO4	A	407	-	4,4,4	0.96	0	6,6,6	0.43	0
4	GOL	C	406	-	5,5,5	0.37	0	5,5,5	0.36	0
3	PO4	B	403	-	4,4,4	0.95	0	6,6,6	0.46	0
2	NAG	B	401	1	14,14,15	0.23	0	17,19,21	0.49	0
2	NAG	D	403	1	14,14,15	0.25	0	17,19,21	0.41	0
2	NAG	C	403	1	14,14,15	0.24	0	17,19,21	0.47	0
3	PO4	D	404	-	4,4,4	0.94	0	6,6,6	0.47	0
3	PO4	D	405	-	4,4,4	0.96	0	6,6,6	0.48	0
2	NAG	C	402	1	14,14,15	0.37	0	17,19,21	0.56	0
2	NAG	A	401	1	14,14,15	0.35	0	17,19,21	0.47	0
2	NAG	D	402	1	14,14,15	0.38	0	17,19,21	0.54	0
3	PO4	A	404	-	4,4,4	0.95	0	6,6,6	0.43	0
2	NAG	C	401	1	14,14,15	0.28	0	17,19,21	0.52	0
4	GOL	C	407	-	5,5,5	0.37	0	5,5,5	0.36	0
3	PO4	A	406	-	4,4,4	0.95	0	6,6,6	0.49	0
3	PO4	C	404	-	4,4,4	0.94	0	6,6,6	0.49	0
2	NAG	A	403	1	14,14,15	0.32	0	17,19,21	0.41	0
3	PO4	B	404	-	4,4,4	0.95	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	402	1	-	2/6/23/26	0/1/1/1
2	NAG	D	401	1	-	4/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	401	1	-	4/6/23/26	0/1/1/1
2	NAG	B	401	1	-	2/6/23/26	0/1/1/1
2	NAG	D	402	1	-	2/6/23/26	0/1/1/1
2	NAG	D	403	1	-	2/6/23/26	0/1/1/1
2	NAG	C	403	1	-	0/6/23/26	0/1/1/1
4	GOL	A	408	-	-	2/4/4/4	-
4	GOL	C	407	-	-	2/4/4/4	-
4	GOL	C	406	-	-	2/4/4/4	-
2	NAG	B	402	1	-	2/6/23/26	0/1/1/1
2	NAG	C	401	1	-	0/6/23/26	0/1/1/1
2	NAG	A	403	1	-	0/6/23/26	0/1/1/1
2	NAG	A	402	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	401	NAG	C2-N2-C7	4.50	128.93	122.90
2	D	401	NAG	C1-C2-N2	2.16	113.84	110.43

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	408	GOL	O1-C1-C2-C3
4	C	406	GOL	O1-C1-C2-C3
2	D	403	NAG	O5-C5-C6-O6
2	B	402	NAG	O5-C5-C6-O6
2	A	401	NAG	O5-C5-C6-O6
2	C	402	NAG	O5-C5-C6-O6
2	D	403	NAG	C4-C5-C6-O6
2	A	402	NAG	O5-C5-C6-O6
2	B	402	NAG	C4-C5-C6-O6
2	A	401	NAG	C4-C5-C6-O6
2	A	402	NAG	C4-C5-C6-O6
2	C	402	NAG	C4-C5-C6-O6
2	A	401	NAG	C8-C7-N2-C2
2	A	401	NAG	O7-C7-N2-C2
2	B	401	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	401	NAG	O7-C7-N2-C2
2	D	401	NAG	C8-C7-N2-C2
2	D	401	NAG	O7-C7-N2-C2
4	C	407	GOL	O1-C1-C2-O2
4	C	407	GOL	O1-C1-C2-C3
4	A	408	GOL	O1-C1-C2-O2
2	D	402	NAG	O5-C5-C6-O6
2	D	402	NAG	C4-C5-C6-O6
4	C	406	GOL	O1-C1-C2-O2
2	D	401	NAG	C1-C2-N2-C7
2	D	401	NAG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	NAG	1	0
4	C	406	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/334 (86%)	0.12	11 (3%) 44 39	25, 39, 63, 95	0
1	B	291/334 (87%)	0.58	11 (3%) 44 39	30, 54, 86, 122	0
1	C	290/334 (86%)	0.29	12 (4%) 41 37	28, 44, 70, 105	0
1	D	292/334 (87%)	0.25	9 (3%) 51 47	26, 43, 68, 98	0
All	All	1163/1336 (87%)	0.31	43 (3%) 45 40	25, 45, 75, 122	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	310	ALA	4.9
1	B	332	LEU	4.9
1	D	332	LEU	4.7
1	D	25	SER	4.5
1	C	25	SER	4.4
1	D	311	VAL	4.4
1	B	26	PRO	4.0
1	B	25	SER	3.7
1	A	333	PRO	3.3
1	B	237	ASN	3.3
1	A	25	SER	3.3
1	A	239	SER	3.3
1	A	240	MET	3.3
1	D	75	ALA	3.1
1	C	26	PRO	3.1
1	A	310	ALA	3.0
1	B	336	CYS	2.9
1	C	292	ARG	2.9
1	C	336	CYS	2.9
1	B	263	VAL	2.8
1	C	311	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	31	CYS	2.8
1	A	336	CYS	2.8
1	B	238	LEU	2.7
1	C	291	GLN	2.6
1	A	264	CYS	2.5
1	A	334	LYS	2.4
1	B	333	PRO	2.3
1	A	215	ALA	2.3
1	A	309	CYS	2.3
1	D	57	ALA	2.2
1	B	189	ARG	2.2
1	C	242	PRO	2.2
1	A	335	CYS	2.2
1	D	26	PRO	2.2
1	C	259	ASP	2.1
1	D	242	PRO	2.1
1	C	45	GLN	2.1
1	D	310	ALA	2.0
1	C	29	GLY	2.0
1	C	294	ALA	2.0
1	B	244	GLU	2.0
1	C	163	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	403	14/15	0.63	0.21	64,84,90,94	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	C	403	14/15	0.64	0.20	74,97,102,102	0
3	PO4	B	404	5/5	0.64	0.16	107,108,110,113	0
2	NAG	D	403	14/15	0.66	0.23	81,98,103,106	0
3	PO4	A	406	5/5	0.74	0.19	119,121,122,122	0
3	PO4	D	405	5/5	0.75	0.14	108,110,110,111	0
2	NAG	C	402	14/15	0.77	0.15	58,65,79,84	0
3	PO4	C	404	5/5	0.77	0.18	100,100,102,102	0
3	PO4	B	403	5/5	0.77	0.17	101,104,105,109	0
4	GOL	A	408	6/6	0.77	0.23	83,87,88,88	0
4	GOL	C	406	6/6	0.78	0.20	73,78,79,80	0
2	NAG	A	402	14/15	0.79	0.14	44,58,69,70	0
3	PO4	D	404	5/5	0.80	0.18	103,103,105,106	0
2	NAG	D	402	14/15	0.80	0.15	56,67,76,78	0
2	NAG	D	401	14/15	0.81	0.13	53,62,73,77	0
3	PO4	D	406	5/5	0.82	0.13	98,99,104,105	0
2	NAG	B	402	14/15	0.83	0.15	64,76,88,90	0
2	NAG	C	401	14/15	0.83	0.13	48,58,72,76	0
3	PO4	A	405	5/5	0.84	0.15	105,106,108,110	0
3	PO4	A	407	5/5	0.84	0.19	96,96,98,99	0
2	NAG	B	401	14/15	0.85	0.15	61,68,89,94	0
2	NAG	A	401	14/15	0.86	0.12	43,53,63,63	0
4	GOL	C	407	6/6	0.86	0.16	68,70,74,77	0
3	PO4	A	404	5/5	0.87	0.15	88,93,93,94	0
3	PO4	C	405	5/5	0.88	0.11	102,104,105,105	0
5	CL	A	409	1/1	0.95	0.10	68,68,68,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.