



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:33 PM UTC

PDB ID : 1O1H / pdb_00001o1h
Title : STRUCTURE OF GLUCOSE ISOMERASE DERIVATIZED WITH KR.
Authors : Nowak, E.; Panjikar, S.; Tucker, P.A.
Deposited on : 2002-11-07
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	
1	B	387	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	KR	A	1008	-	-	X	-
5	KR	B	1005	-	-	X	-
5	KR	B	1007	-	-	X	-
7	MPD	A	1015	X	-	-	-
7	MPD	B	1016	X	X	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7071 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	2	0
			3050	1917	551	574	8			
1	B	386	Total	C	N	O	S	0	2	0
			3046	1915	548	575	8			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		


- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		
4	B	2	Total	Cl	0	0
			2	2		

- Molecule 5 is KRYPTON (CCD ID: KR) (formula: Kr).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Kr	0	0
			2	2		
5	B	2	Total	Kr	0	0
			2	2		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		
6	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			8	6	2		
7	B	1	Total	C	O	0	0
			8	6	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	435	Total	O	0	0
			435	435		
8	B	496	Total	O	0	0
			496	496		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylose isomerase



- Molecule 1: Xylose isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.33Å 93.07Å 98.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.40 20.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.40) 99.9 (20.00-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.36 (at 1.40Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.156 , 0.161 0.154 , 0.160	Depositor DCC
R_{free} test set	1548 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	8.4	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7071	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.70 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5077e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, CA, MG, CL, TRS, KR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	4/3133 (0.1%)	1.07	12/4241 (0.3%)
1	B	0.43	0/3123	1.07	14/4230 (0.3%)
All	All	0.45	4/6256 (0.1%)	1.07	26/8471 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
All	All	0	5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	23	ASP	C-N	5.27	1.41	1.33
1	A	24[A]	PRO	N-CD	5.14	1.54	1.47
1	A	24[B]	PRO	N-CD	5.14	1.54	1.47
1	A	23	ASP	C-O	-5.12	1.17	1.24

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	23[A]	ASP	CA-CB-CG	7.40	120.00	112.60
1	B	23[B]	ASP	CA-CB-CG	7.40	120.00	112.60
1	A	186	PRO	N-CA-CB	7.14	110.74	103.25
1	A	185	GLU	CA-C-O	-7.10	110.43	120.16
1	A	244	ASP	CA-CB-CG	6.97	119.57	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	344	ASP	CA-CB-CG	6.94	119.54	112.60
1	B	185	GLU	CA-C-O	-6.72	110.96	120.16
1	A	23	ASP	CB-CA-C	6.70	123.37	110.17
1	B	244	ASP	CA-CB-CG	6.62	119.22	112.60
1	B	254	ASP	CA-CB-CG	6.35	118.95	112.60
1	A	186	PRO	CA-N-CD	-6.17	103.36	112.00
1	B	186	PRO	N-CA-CB	6.09	109.64	103.25
1	B	186	PRO	CA-N-CD	-5.96	103.66	112.00
1	B	185	GLU	CA-C-N	5.95	127.28	119.84
1	B	185	GLU	C-N-CA	5.95	127.28	119.84
1	A	344	ASP	CA-CB-CG	5.86	118.46	112.60
1	B	54	ASP	CA-CB-CG	5.68	118.28	112.60
1	B	356	PHE	CA-CB-CG	-5.58	108.22	113.80
1	B	335	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	23	ASP	CA-CB-CG	5.53	118.13	112.60
1	B	109	ASP	CA-CB-CG	5.48	118.08	112.60
1	A	254	ASP	CA-CB-CG	5.43	118.03	112.60
1	A	109	ASP	CA-CB-CG	5.42	118.03	112.60
1	A	23	ASP	CA-C-N	-5.12	114.55	119.87
1	A	23	ASP	C-N-CA	-5.12	114.55	119.87
1	A	54	ASP	CA-CB-CG	5.08	117.68	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	185	GLU	Mainchain,Peptide
1	B	185	GLU	Mainchain,Peptide
1	B	24[B]	PRO	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3050	0	2920	42	0
1	B	3046	0	2913	50	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	2	0	0	1	0
4	B	2	0	0	2	0
5	A	2	0	0	6	0
5	B	2	0	0	6	0
6	A	8	0	11	0	0
6	B	8	0	11	0	0
7	A	8	0	13	1	0
7	B	8	0	13	8	0
8	A	435	0	0	17	10
8	B	496	0	0	31	13
All	All	7071	0	5881	100	17

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:ARG:HD2	8:A:1379:HOH:O	1.29	1.26
1:B:296:ASP:HB2	8:B:1487:HOH:O	1.14	1.24
1:B:79:ASP:HB2	8:B:1505:HOH:O	1.06	1.19
1:A:373:ARG:HD3	8:A:1096:HOH:O	1.40	1.19
1:B:75:ARG:HD2	8:B:1370:HOH:O	1.46	1.14
1:B:204:ARG:NH1	8:B:1497:HOH:O	1.83	1.11
1:B:265:ARG:HD2	8:B:1459:HOH:O	1.50	1.11
7:B:1016:MPD:H52	8:B:1167:HOH:O	1.50	1.11
1:A:60:PHE:CE1	8:A:1369:HOH:O	2.05	1.09
7:B:1016:MPD:HM1	8:B:1479:HOH:O	1.52	1.09
1:A:204:ARG:NH1	8:A:1392:HOH:O	1.87	1.07
1:B:265:ARG:CD	8:B:1459:HOH:O	2.06	1.03
1:B:206:GLU:HG3	8:B:1493:HOH:O	1.58	1.02
7:B:1016:MPD:C5	8:B:1167:HOH:O	2.05	1.00
1:A:265:ARG:CD	8:A:1379:HOH:O	1.91	1.00
1:B:84:LYS:HE3	8:B:1280:HOH:O	1.62	0.99
1:B:84:LYS:HE2	8:B:1429:HOH:O	1.62	0.97
7:B:1016:MPD:CM	8:B:1452:HOH:O	2.16	0.93
1:A:95:HIS:HD2	1:A:97:VAL:H	1.17	0.89
1:B:95:HIS:HD2	1:B:97:VAL:H	1.19	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:79:ASP:CB	8:B:1505:HOH:O	1.82	0.87
1:A:204:ARG:HH21	5:A:1008:KR:KR	2.23	0.82
1:B:91:ASN:HD21	1:B:94:THR:H	1.27	0.81
1:A:91:ASN:HD21	1:A:94:THR:H	1.26	0.80
1:A:60:PHE:HE1	8:A:1369:HOH:O	1.51	0.79
5:A:1008:KR:KR	1:B:233:GLN:HE21	2.27	0.78
1:B:79:ASP:OD2	8:B:1129:HOH:O	2.00	0.77
1:B:373:ARG:HG2	8:B:1198:HOH:O	1.84	0.77
1:A:233:GLN:HE21	5:B:1007:KR:KR	2.29	0.75
1:A:3:GLN:NE2	8:A:1411:HOH:O	2.15	0.74
1:B:41:ARG:HD3	8:B:1448:HOH:O	1.89	0.71
4:A:1010:CL:CL	8:A:1256:HOH:O	2.46	0.70
1:A:22:ARG:HD3	8:A:1347:HOH:O	1.92	0.69
4:B:1009:CL:CL	8:B:1187:HOH:O	2.49	0.67
1:A:1:ASN:N	8:A:1389:HOH:O	1.89	0.67
1:B:204:ARG:HE	5:B:1007:KR:KR	2.37	0.67
1:A:91:ASN:HD22	1:A:91:ASN:C	2.03	0.66
1:B:135:ALA:HB2	5:B:1005:KR:KR	2.58	0.65
1:B:91:ASN:HD22	1:B:91:ASN:C	2.05	0.65
1:B:84:LYS:CE	8:B:1429:HOH:O	2.31	0.64
7:B:1016:MPD:HM1	8:B:1452:HOH:O	1.92	0.64
1:A:93:PHE:HE1	7:A:1015:MPD:H11	1.64	0.62
1:B:95:HIS:CD2	1:B:97:VAL:H	2.10	0.61
1:A:135:ALA:HB2	5:A:1006:KR:KR	2.60	0.61
1:A:95:HIS:CD2	1:A:97:VAL:H	2.09	0.61
1:A:91:ASN:ND2	1:A:94:THR:H	1.99	0.60
1:B:326:GLN:HG2	8:B:1341:HOH:O	2.01	0.60
1:B:265:ARG:NH2	4:B:1011:CL:CL	2.64	0.60
1:B:75:ARG:NH2	1:B:127:GLU:HG2	2.18	0.59
1:B:9:ARG:NH1	8:B:1180:HOH:O	2.36	0.59
1:A:64:ASP:O	1:A:68:GLU:HG2	2.05	0.57
1:B:127:GLU:OE1	8:B:1376:HOH:O	2.17	0.57
1:A:236:TRP:CZ3	1:B:204:ARG:HG3	2.39	0.56
1:A:5:THR:HB	1:A:6:PRO:HD2	1.86	0.56
1:A:249:ASN:ND2	8:A:1398:HOH:O	2.22	0.55
1:B:91:ASN:ND2	1:B:94:THR:H	2.00	0.55
7:B:1016:MPD:O4	8:B:1232:HOH:O	2.18	0.55
1:B:75:ARG:HH22	1:B:127:GLU:HG2	1.70	0.54
1:A:322:ASP:OD2	1:A:381:HIS:HE1	1.91	0.53
8:A:1191:HOH:O	1:B:95:HIS:HE1	1.92	0.53
1:B:322:ASP:OD2	1:B:381:HIS:HE1	1.92	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:ARG:HH21	5:B:1007:KR:KR	2.52	0.53
1:B:71:VAL:O	1:B:75:ARG:HG3	2.09	0.53
1:B:80:ASP:OD2	8:B:1346:HOH:O	2.19	0.53
1:B:5:THR:HB	1:B:6:PRO:HD2	1.90	0.53
7:B:1016:MPD:H51	8:B:1167:HOH:O	1.89	0.52
1:A:327:GLU:OE2	8:A:1252:HOH:O	2.19	0.52
1:A:95:HIS:HE1	8:B:1213:HOH:O	1.93	0.52
1:B:373:ARG:CG	8:B:1198:HOH:O	2.48	0.52
1:A:91:ASN:ND2	1:A:93:PHE:H	2.09	0.51
1:B:91:ASN:ND2	1:B:93:PHE:H	2.09	0.51
1:A:381:HIS:HD2	8:A:1173:HOH:O	1.92	0.51
1:B:91:ASN:HD22	1:B:93:PHE:H	1.59	0.50
1:B:381:HIS:HD2	8:B:1193:HOH:O	1.93	0.50
1:B:324:GLU:CD	1:B:386:ARG:HH22	2.19	0.50
1:A:23:ASP:HB2	1:A:24[A]:PRO:HD2	1.94	0.50
1:B:31:ARG:HG3	1:B:295:PHE:HE1	1.77	0.49
1:B:249:ASN:ND2	8:B:1228:HOH:O	2.22	0.49
1:A:204:ARG:HE	5:A:1008:KR:KR	2.56	0.49
1:A:249:ASN:ND2	1:A:262:GLY:HA2	2.28	0.48
7:B:1016:MPD:O2	8:B:1242:HOH:O	2.20	0.48
1:A:295:PHE:HE1	8:A:1134:HOH:O	1.95	0.48
1:A:60:PHE:CD1	8:A:1308:HOH:O	2.55	0.48
1:B:75:ARG:HH22	1:B:127:GLU:CG	2.28	0.47
1:A:91:ASN:HD22	1:A:93:PHE:H	1.63	0.47
1:A:31:ARG:HH12	1:A:33:LEU:HD23	1.78	0.47
1:B:164:LEU:HD13	5:B:1005:KR:KR	2.78	0.45
1:A:221:GLN:HE21	1:A:248:GLN:HB3	1.83	0.44
1:A:91:ASN:HD21	1:A:94:THR:N	2.04	0.43
1:B:221:GLN:HE21	1:B:248:GLN:HB3	1.83	0.43
1:A:96:PRO:HB2	1:B:365:ALA:HB1	2.00	0.43
1:B:91:ASN:C	1:B:91:ASN:ND2	2.76	0.43
1:A:315:ARG:HA	1:A:315:ARG:HD3	1.86	0.42
1:B:249:ASN:ND2	1:B:262:GLY:HA2	2.34	0.42
5:A:1008:KR:KR	1:B:237:ALA:HB2	2.81	0.41
1:A:60:PHE:CZ	8:A:1369:HOH:O	2.55	0.41
1:B:164:LEU:CD1	5:B:1005:KR:KR	3.30	0.40
1:A:204:ARG:NH2	5:A:1008:KR:KR	3.05	0.40
1:A:227:PHE:CZ	1:A:231:ILE:HD11	2.56	0.40

All (17) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1431:HOH:O	8:A:1445:HOH:O[2_775]	1.92	0.28
8:A:1429:HOH:O	8:B:1444:HOH:O[4_566]	1.95	0.25
8:A:1292:HOH:O	8:B:1472:HOH:O[1_655]	2.02	0.18
8:A:1337:HOH:O	8:A:1427:HOH:O[4_467]	2.07	0.13
8:B:1138:HOH:O	8:B:1210:HOH:O[4_566]	2.07	0.13
8:B:1139:HOH:O	8:B:1148:HOH:O[4_566]	2.07	0.13
8:A:1114:HOH:O	8:B:1129:HOH:O[4_566]	2.08	0.12
8:A:1255:HOH:O	8:A:1368:HOH:O[2_775]	2.09	0.11
8:A:1315:HOH:O	8:B:1472:HOH:O[1_655]	2.13	0.07
8:B:1127:HOH:O	8:B:1138:HOH:O[4_466]	2.13	0.07
8:B:1138:HOH:O	8:B:1293:HOH:O[4_566]	2.13	0.07
8:B:1138:HOH:O	8:B:1165:HOH:O[4_566]	2.13	0.07
8:A:1412:HOH:O	8:B:1472:HOH:O[1_655]	2.14	0.06
8:B:1148:HOH:O	8:B:1221:HOH:O[4_466]	2.14	0.06
8:A:1224:HOH:O	8:A:1337:HOH:O[4_567]	2.15	0.05
8:B:1295:HOH:O	8:B:1316:HOH:O[4_566]	2.16	0.04
8:A:1284:HOH:O	8:B:1504:HOH:O[1_655]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/387 (100%)	375 (97%)	9 (2%)	2 (0%)	24	7
1	B	386/387 (100%)	373 (97%)	11 (3%)	2 (0%)	24	7
All	All	772/774 (100%)	748 (97%)	20 (3%)	4 (0%)	24	7

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	185	GLU
1	B	185	GLU
1	A	186	PRO
1	B	186	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/303 (101%)	303 (99%)	2 (1%)	76	52
1	B	304/303 (100%)	302 (99%)	2 (1%)	76	52
All	All	609/606 (100%)	605 (99%)	4 (1%)	76	52

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	204	ARG
1	B	68	GLU
1	B	91	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	95	HIS
1	A	221	GLN
1	A	233	GLN
1	A	308	ASN
1	A	347	GLN
1	A	376	GLN
1	A	381	HIS
1	B	91	ASN
1	B	95	HIS
1	B	221	GLN
1	B	233	GLN
1	B	249	ASN
1	B	308	ASN
1	B	347	GLN
1	B	376	GLN
1	B	381	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	MPD	A	1015	-	7,7,7	1.57	1 (14%)	9,10,10	3.19	3 (33%)
7	MPD	B	1016	-	7,7,7	1.61	1 (14%)	9,10,10	3.24	3 (33%)
6	TRS	A	1013	3	7,7,7	1.65	2 (28%)	9,9,9	3.02	5 (55%)
6	TRS	B	1014	3	7,7,7	1.58	2 (28%)	9,9,9	3.11	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MPD	A	1015	-	1/1/2/2	4/5/5/5	-
7	MPD	B	1016	-	1/1/2/2	5/5/5/5	-
6	TRS	A	1013	3	-	2/9/9/9	-
6	TRS	B	1014	3	-	2/9/9/9	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	1016	MPD	O4-C4	-3.99	1.26	1.43
7	A	1015	MPD	O4-C4	-3.92	1.27	1.43
6	A	1013	TRS	O2-C2	-2.59	1.34	1.42
6	B	1014	TRS	C3-C	-2.52	1.46	1.53
6	A	1013	TRS	C3-C	-2.51	1.46	1.53
6	B	1014	TRS	O2-C2	-2.29	1.35	1.42

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	1016	MPD	C5-C4-C3	-7.03	79.03	111.67
7	A	1015	MPD	C5-C4-C3	-6.99	79.22	111.67
6	B	1014	TRS	O3-C3-C	5.57	126.40	110.88
7	B	1016	MPD	O4-C4-C3	-5.13	90.94	111.35
7	A	1015	MPD	O4-C4-C3	-5.11	91.03	111.35
6	A	1013	TRS	O3-C3-C	5.02	124.86	110.88
6	A	1013	TRS	O1-C1-C	4.27	122.76	110.88
6	B	1014	TRS	O1-C1-C	4.02	122.08	110.88
7	B	1016	MPD	O4-C4-C5	-3.76	93.27	109.45
7	A	1015	MPD	O4-C4-C5	-3.74	93.35	109.45
6	A	1013	TRS	C2-C-C1	3.63	120.31	110.66
6	B	1014	TRS	C2-C-C1	3.62	120.30	110.66
6	B	1014	TRS	C1-C-N	-3.23	99.92	108.17
6	A	1013	TRS	O2-C2-C	3.16	119.70	110.88
6	B	1014	TRS	O2-C2-C	2.87	118.89	110.88
6	A	1013	TRS	C1-C-N	-2.87	100.84	108.17

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	A	1015	MPD	C4
7	B	1016	MPD	C4

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	1015	MPD	C1-C2-C3-C4
7	A	1015	MPD	C2-C3-C4-O4
7	B	1016	MPD	C2-C3-C4-O4
7	B	1016	MPD	C2-C3-C4-C5
6	A	1013	TRS	C1-C-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	1014	TRS	C1-C-C3-O3
7	A	1015	MPD	C2-C3-C4-C5
7	B	1016	MPD	O2-C2-C3-C4
7	A	1015	MPD	CM-C2-C3-C4
7	B	1016	MPD	C1-C2-C3-C4
7	B	1016	MPD	CM-C2-C3-C4
6	A	1013	TRS	N-C-C3-O3
6	B	1014	TRS	N-C-C3-O3

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1015	MPD	1	0
7	B	1016	MPD	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/387 (99%)	-0.26	12 (3%) 51 54	5, 8, 22, 32	2 (0%)
1	B	386/387 (99%)	-0.43	9 (2%) 61 63	4, 8, 18, 27	2 (0%)
All	All	772/774 (99%)	-0.35	21 (2%) 56 58	4, 8, 20, 32	4 (0%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	385	ALA	3.6
1	A	81	THR	3.3
1	B	295	PHE	3.1
1	A	295	PHE	3.0
1	B	24[A]	PRO	2.8
1	A	63	SER	2.8
1	A	24[A]	PRO	2.7
1	A	174	ASP	2.7
1	B	75	ARG	2.6
1	B	296	ASP	2.6
1	A	65	SER	2.5
1	B	65	SER	2.4
1	A	300	ALA	2.4
1	A	80	ASP	2.3
1	A	61	GLY	2.3
1	B	25	PHE	2.3
1	A	62	SER	2.1
1	A	1	ASN	2.1
1	B	80	ASP	2.1
1	B	373	ARG	2.0
1	B	79	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	MPD	B	1016	8/8	0.63	0.26	35,35,35,35	0
7	MPD	A	1015	8/8	0.70	0.18	30,30,31,31	0
4	CL	B	1011	1/1	0.87	0.17	21,21,21,21	0
4	CL	B	1009	1/1	0.90	0.14	18,18,18,18	0
4	CL	A	1012	1/1	0.91	0.18	23,23,23,23	0
6	TRS	A	1013	8/8	0.92	0.11	11,13,13,14	0
6	TRS	B	1014	8/8	0.92	0.11	11,12,13,15	0
4	CL	A	1010	1/1	0.93	0.12	21,21,21,21	0
5	KR	B	1005	1/1	0.98	0.06	11,11,11,11	1
5	KR	A	1008	1/1	0.99	0.03	12,12,12,12	1
3	MG	A	1003	1/1	0.99	0.02	6,6,6,6	0
5	KR	A	1006	1/1	0.99	0.03	10,10,10,10	1
2	CA	A	1001	1/1	1.00	0.01	4,4,4,4	0
3	MG	B	1004	1/1	1.00	0.02	5,5,5,5	0
2	CA	B	1002	1/1	1.00	0.01	4,4,4,4	0
5	KR	B	1007	1/1	1.00	0.02	11,11,11,11	1

6.5 Other polymers [i](#)

There are no such residues in this entry.